HepciFer study: Hepcidin overexpression after hepatectomy : Does iron supplementation make sense ?

Benoliel A¹., Assouline B.², Legouis D.², Blaudzun G.² Kraiem A.², Favre M.², Schiffer E.²

¹Hôpital Cochin, Dept of Anaesthesiology & Intensive Care, Paris, France,

²Hôpitaux Universitaires Geneva (HUG), Dept of Anaesthesiology, Geneva, Switzerland

Background and goal of the study

Anemia is frequent in liver surgery with an estimated incidence of 32%⁽¹⁾. The causes of anemia in surgical patients are often the iron deficiency anemia and the functional iron deficiency (FID). FID is associated with an elevated level of hepcidin (synthesized by the liver), the central regulatory protein of iron metabolism. Our hypothesis was that Hepcidin is overexpressed after liver surgery and contributes to postoperative anemia. We aimed to assess the impact of intravenous iron administration in the immediate postoperative period on hemoglobin levels.

Material and methods

The HepciFer trial was a single-center, randomized, placebo-controlled, double-blinded clinical trial (RCT) in 50 patients undergoing liver surgery. Adult patients undergoing elective liver surgery were screened and included from September 2015 to August 2016. According to the randomization process, patients received either ferric carboxymaltose (15mg/Kg, maximum 1g) or placebo 4 hours after surgery. Primary outcome assessed the difference of plasma Hb levels between the two groups seven days after surgery. Secondary outcomes included plasma level of Hepcidin and iron status at different time points after surgery, length of hospitalization and incidence of blood transfusion.

Results

Both groups were comparable regarding patients' baseline characteristics, type of surgery, pathology analyses and postoperative complications.

Iron supplementation was associated with an increase of 1.2 g/dL of Hb level ($p=0.04$) and higher hematocrit level (34.7 (30.9-36.6) vs 30.4 (29.0-33.5); $P=0.02$) at POD 7. Hepcidin level was also higher after iron supplementation (50ng/l [30,4-134] vs 8,4ng/l [4,4-25] ; $P< 0.0001$) whereas IL-6 level did not differ across groups. At POD 7, the median ferritin level was 1307ug/l (1099-1671) vs 241ug/l (174-497) in the iron group ($p<0.001$). Finally, length of hospitalisation and rate of blood transfusion were comparable.

Conclusion

This study is to our knowledge the first RCT to evaluate the impact of IV iron administration in the postoperative period after liver surgery. Our findings suggest that IV iron supplementation increase Hb and Hct level at POD 7 by treating the FID. In addition, this trial adds new knowledge regarding the physiopathology of iron metabolism and Hepcidin kinetics and thus, provides a better understanding of mechanisms involved in FID during postoperative period after liver surgery.

(1) Tohme S, and al. HPB. mars 2016;18(3):255-61.