Spontaneous vestibular schwannoma regression: a case control study

T. Tikka, C.P. Yiannakis, E. Stapleton, R. Locke,
J.A. Crowther, W.A.S. Taylor, G. Kontorinis

ENT Department, Queen Elizabeth University Hospital Glasgow

Introduction

- Vestibular schwannomas (VS) account for 85% of tumours in the cerebellopontine angle.
- The incidence is rising with an average of 1.3-2 new cases per 100,000 persons per year.
- VS can grow in various patterns.
- The majority of cases have static appearances from the time of diagnosis, a significant number of cases show definite growth and interestingly a small number of tumours demonstrate regression without any treatment.
- Neutrophil to lymphocyte ration (NLR) can predict VS growth. $NLR > 3.05$ being associated with VS growth.
- To date no factors have been linked with spontaneous tumour shrinkage.

Aim

- To explore behaviours of regressing VS
- Identify possible predictors of VS regression

Methods

- Retrospective case-control study
- Cases of regressed VS were identified from a the centralised west of Scotland VS database over a 15 year period (2000 – 2015).
- Random selection of equal number of growing VS from the same database for direct comparison.

Results

- From the 540 patients included in the database: 28 patients had spontaneous regression * with a mean follow-up of 122 months (47 – 180)

* $\geq 2\text{mm}$ decrease per annum in VS largest diameter on MRI – AAOHNS guidelines)

- Mean absolute regression: 3.9mm (range, $2\text{mm} - 10\text{mm}$).
- Relative tumour regression: 25.7% (range, $10.3\% - 55.6\%$)

- Gradual regression was observed in 60.7% ($n=17$) with the remaining patients ($n=11$; 39.3%) having a rapid tumour regression between consecutive scans.
- 25% ($n=7$) showed growth followed by regression, 57.1% ($n=16$) showed regression initially followed by a period of stability with the remainder ($n=5$; 17.9%) showing regression after a period of stability.

Patient demographics and tumour characteristics at diagnosis	Regressing VS N=28	Growing VS N=28	p-value
Gender	15F: 12M	17F: 11M	0.589
Age at diagnosis (years)	62.1 (SD 12.8)	59.6 (SD 11.9)	0.456
Presenting symptoms*			0.685
- Unilateral hearing loss	13	11	
- Asymmetric hearing loss	8	6	
- Unilateral tinnitus	4	5	
- Vertigo/balance problems	2	4	
- Facial weakness	0	1	
- Incidental finding	4	6	
Tumour laterality			0.694
Right	15	16	
Left	13	12	
Size of tumour at diagnosis (mm)	15.8 (SD 6.5)	10.7 (SD 7.4)	0.014
Tumour position at diagnosis			0.491
Intacanalicular only	0	2	
Extracanalicular component	28	26	

- Regressing VS had a significantly larger size than the control group ($p=0.014$); while the regressing tumours were located further from the fundus than the controlled group (mean distance: 8.2mm vs 6.3mm ; $p=0.044$).

Distance (mm) at time of diagnosis	Growing VS N=28	Regressing VS N=28	p-value
VS to fundus	6.3 (SD 3.4)	8.2 (SD 3.4)	0.044
VS to brainstem	3.3 (SD 4.4)	2.4 (SD 2.8)	0.675
VS to clivus	29.9 (SD 8)	26.5 (SD 7.3)	0.102
VS to sigmoid	29.5 (SD 4.5)	27.6 (SD 4.8)	0.125
Clivus/sigmoid ratio	1.03 (SD 0.32)	0.99 (SD 0.36)	0.658

- The NLR was 2.30 (SD 1.3) in the regressing VS group and 2.04 (SD 0.66) in the control group ($p=0.371$) with available bloods from 24 and 20 patients respectively. 4 (20%) patients with regressed VS had an NLR above 3.05 compared to 2 (8.3%) in the control group ($p=0.384$).

Conclusions

- A link between tumour distance from fundus and VS regression probability is identified.
- This information could be used when counselling patients about future tumour behaviour; regression, whilst rare is still a realistic possibility.