Seasonality and Bipolar Disorder Hospitalizations

A cross sectional study in a portuguese hospital

Z. Gameiro⁽¹⁾, J. Valente⁽²⁾, G. Lima⁽¹⁾

1- Barreiro-Montijo Hospital Center, Department of Psychiatry and Mental Health, Barreiro, Portugal

2- Primary Health Clusters Center ACES Sintra, Public Health Unit, Sintra, Portugal

19TH WPA WORLD
CONGRESS OF PSYCHIATRY
LISBON, 21-24 AUGUST, 2019

Objectives

Characterize and investigate determinants of seasonality in hospitalization of patients with Bipolar Disorder (BD).

Background

Seasonal hospitalization patterns have been described in BD, with higher hospital admission rates in Spring and in Summer in temperate countries. We aimed to investigate possible determinants of seasonality in hospital admissions of patients with BD.

Materials and Methods

Data was obtained from all 111 psychiatry admissions of patients with BD as the primary diagnose in Barreiro-Montijo Hospital Center, located in Lisbon Metropolitan Area, in 2017. Data was analysed with SPSS® version 22. Association between admission's season, sex and BD polarity was investigated with chi-square test and association between admission's season and age was investigated with ANOVA test. The associations were considered significant if p value <0,05.

Results

The majority of BD admissions were females 75 (67,6%). The average age at admission was 46,3 years. Spring was the season with the highest number of BD admissions 31 (27,9%), followed by summer 30 (27,0%), autumn 28 (25,2%) and winter 22 (19,8%). The majority of BD admissions were due to maniac/hypomaniac episodes 52 (50,5%), with 23 (20,7%) admissions due to depressive episodes. No significant differences were found between the season of admission and sex (p=0,703), age (p=0,768) and BD characteristics (p=0,235).

BD admissions / Month

BD admissions / Sex

BD admissions / Age group

BD admissions / Episode type

Conclusions

Increased knowledge about the patterns of seasonality and its determinants in BD patients may lead to improved healthcare service planning and to the development of specific programs to BD patients according to the season.

References

- Fellinger M, Waldhoer T, König D et al. Seasonality in bipolar disorder: Effect of sex and age. J Affect Disord. 2019 Jan 15;243:322-326
- Geoffroy PA, Bellivier F, Scott J and Etain B. Seasonality and bipolar disorder: A systematic review, from admission rates to seasonality of symptoms. J Affect Disord. 2014 Oct;168:210-23
- Aguglia A, Borsotti A, Maina G. Bipolar disorders: is there an influence of seasonality or photoperiod? Braz J Psychiatry. 2018 Jan-Mar;40(1):6-11