‘Nice to know there is help if needed’; Review of Quality of Life for Lymphoma patients on supported self-management pathway

Alia Nizam and Hilary Plant

Macmillan Support and Information Service at University College London Hospital NHS Trust

Introduction

The importance of supported self - management for people living with and beyond cancer has been recognised as one of the key priorities in the Cancer Strategy for England (2015-2020). Supported self-management is a collaborative partnership between patients and healthcare professionals that empowers the individual to self-manage their condition and well-being without the need to participate in routine follow-up appointments. There is evidence to suggest that routine follow-up appointments do not always meet individual needs and disease recurrences are more likely to be picked up in between appointments. Supported self-management pathways aim to give individuals the confidence to be able to self-manage, live actively and well, and recognise the signs and symptoms of when to contact the clinical team for advice if needed.

At University College London Hospital NHS Trust supported self-management pathway was introduced in curative lymphoma patients with stable disease. Following treatment, eligible patients are seen in follow-up appointments by the haematologists every three months for a year. Within that year the lymphoma nurse specialists meet the patients to perform their holistic needs assessments, educate them on long term symptom management and signs of recurrence, and offer health and wellbeing advice. The patients are given contact details to enable rapid re-access to the specialist team as required. After the year, patients are no longer followed up in clinic but are on an open access pathway for 4 years where the specialist team relies on the patients to contact them should any problems arise.

Whilst on the open access pathway, the EQ-5D-5L quality of life (QoL) questionnaire and a feedback questionnaire is sent to the patients annually for 4 years. This questionnaire aims to collect data on the quality of life of lymphoma patients as a means of assessing how well patients are reintegrating into their normal lives following treatment. This is important data as measuring and understanding long term quality of life of cancer patients will give healthcare professionals instrumental information of where to focus care, support and resources in the survivorship phase. The feedback questionnaire which is separate to the QoL questionnaire is a locally agreed questionnaire which aims to assess the patients’ confidence in self-management.

Methods and results

The questionnaires are sent via post annually.

The EQ-5D-5L questionnaire covers problems with walking about, washing, dressing, doing usual activities, pain or discomfort and anxiety or depression

Between January 2016 and November 2018, 56 questionnaires have been reviewed and summarised in graph 1 below

Graph 1

Graph 2

Between January 2016 and November 2018, 46 feedback questionnaires have been reviewed for confidence in self-management

Graph 3

Discussion

Currently there is no measure to assess how well patients are supported after treatment for cancer. This does not only include their physical needs, but also other support such as help at home or financial advice. In England there is work in progress in developing a ‘quality of life metric’ with the use of two validated questionnaires. One of the questionnaires is the EQ-5D-5L one. This is to help to understand the quality of life of people who have been treated for cancer and how effective support is.

Lymphoma team has been forward thinking in assessing the quality of life of patients on the supported self-management pathway. The numbers may be low but they provide a unique insight to how patients feel about their care and wellbeing. From the observation of the scale between slight and severe in graph 1, anxiety or depression appears to be the prominent issue, followed by pain or discomfort and mobility. It can be debated that mobility issues could be due to other co-morbidities of the patients and pain/ discomfort due to late effects of treatment. Anxiety and depression is also a common theme that mirrors with results of the holistic needs assessment which is done in the UK as part of the recovery package interventions. To have an in depth understanding about anxiety and depression, it would be useful to review the patients individually to unpick more specific issues. This would enable healthcare professionals to understand and provide more support and sign posting prior to treatment relating to anxiety and depression.

Half the patients appear to be confident in self-management which is a positive reflection on the consultation with the lymphoma nurses.

Conclusion

More data needs to be captured to provide a more robust overview of how patients feel after treatment. On review of the entire pathway, there has been no delayed diagnosis and in cases where relapse has occurred patients have contacted the specialist team for advice. Cancer survival rates are at a record high, it is vital that cancer patient experience is prioritised and care tailored to their needs. Assessing and measuring patients’ quality of life using a standardised metric is a unique way to assess where things are going well and where improvements can be made.