Accelerometers as a novel tool for automatic analysis of infants' spontaneous motility

Fontana C¹, Ottaviani V², Sforza SE¹, Veneroni C², Picciolini O¹, Passera S¹, Fumagalli M¹, Mosca F¹, Dellacà R²

1. NICU, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, University of Milan, Department of Clinical Sciences and Community Health, Italy
2. Dipartimento di Elettronica, Informazione e Bioingegneria, Politecnico di Milano, Italy

BACKGROUND AND AIM: General movements (GM) assessment evaluates infants' spontaneous motility to early detect neuromotor impairments; it depends, however, on examiners' experience and training.

Aim of the study is to develop a novel tool to assess GMs based on automatic measurements and data processing.

METHODS: Observational study. Neonates admitted to NICU without congenital malformations or genetic syndromes, entered the study. At Term equivalent age (TEA), both clinical video-based General Movements (GMs) assessment and automatic recording of GMs were performed.

Clinical GMs was classified: normal (N), cramped synchronized (CS) or poor repertoire (PR). Automatic recording was carried out with four accelerometers placed on the infants' ankles and wrists (Figure1); accelerations were recorded continuously at 150 Hz for at least 10 minutes. The module of acceleration and its variation during time (jerk) have been computed for each limb. Moreover, the Kurtosis index and the Cross-Correlation index have been evaluated: the former evaluates movements variability (the higher the value the lower the variability of movements); the latter measures movements synchronicity (higher values represent greater synchronicity in movements).

RESULTS: 42 infants were enrolled. Infants' characteristics are presented in Table 1. A progressive trend of increasing **kurtosis** values was observed moving from N to CS groups in all limbs (Figure 2), showing a higher variability in movements in the N group. **Cross-correlation** index of the lower limbs showed a statistically significant difference between N and CS and between PR and CS, showing more synchronicity in CS movements. No differences were observed in the upper limbs (Figure 3)

Figure 1 – Experimental setting

Demographic feature	N (n=26)	PR (n=11)	CS (n=5)	p.value
Gestational Age at Birth, mean±SD	31 ± 2.6	30.2 ± 4.4	32.6 ± 4.4	0.407*
Birth Weight, mean±SD	1416 ± 376	1405 ± 828	1654 ± 697	0.669*
Twin Pregnancy, n (%)	15 (58)	4 (36)	2 (40)	0.537*
Small for Gestational Age, n (%)	5 (19)	2 (18)	1 (20)	>0.999*
Severe BPD, n (%)	0(0)	3 (27)	0(0)	0.033*
No Brain Lesions, n (%)	23 (88)	2 (18)	1 (20)	<0.001*
IVH, n (%)				
Low-grade IVH (IVH 1-2)	3 (12)	4 (36)	1 (20)	0.036*
High-grade IVH (IVH 3-4) including PHVD	0(0)	0(0)	2 (40)	
Punctate White Matter Lesions, n (%)	0(0)	4 (36)	0(0)	0.006*
Extensive White Matter Damage, n (%)	0(0)	0(0)	2 (40)	>0.999*
White Matter and Cortical Damage, n (%)	0(0)	1 (9)	0(0)	0.381*
Cerebellar Hemorrhage, n (%)	0(0)	1 (9)	0(0)	0.381*
Intraparenchymal hemorrhage, n (%)	0(0)	1 (9)	0(0)	0.381*
Gestational Age at Discharge, mean±SD	38.1 ± 1.8	41.2 ± 4.9	46.7 ± 5.4	<0.001*
Gestational Age at Time of Analysis, mean±SD	42.6 ± 2.8	42 ± 2.8	42 ± 0.8	0.773*

* One-Way ANOVA, ^Kruskal-Wallis test, *Fisher Exact Test

Table 1 – Baseline characteristics

CONCLUSIONS: Even if preliminary, our study suggests a possible new method to automatically assess GMs in high risk infants in early post-natal life. The tool we developed was feasible and allowed the distinction of normal from cramped synchronized movements. Further studies are required to confirm these preliminary results and to explore other parameters.

Figure 2 – Kurtosis index for each limb
(* p<0.05 compared to N GMs).

The higher the value the lower the variability of movements

Figure 3 – Cross Correlation Index for: left hand-right hand and left foot-right foot

(* p<0.05 compared to N GMs and O p=0.033 compared to CS GMs)
Higher values represent greater synchronicity in movements