Qualitative Analysis of Medical Records of Preanaesthetic Evaluations

J. Pedreira¹, C. Lanzaro¹, L. Ferreira¹, A. Bernardino¹, M. L. Bela¹

1. Centro Hospitalar e Universitário de Coimbra, Portugal

Background

Performance and quality indicators are essential tools in management and quality of medical units. Preanaesthetic evaluation is a central component of clinical anaesthesia. Consultation records should always exist, be easily accessible and meet minimum information requirements.

Goal: evaluate the technical-scientific care in anaesthesiology consultation, from the quality of the consultation records, based on standard 029/2013 (updated 24/04/2015) of *Direção Geral de Saúde* (DGS)¹. This study also aims to evaluate and compare two methods of computerized consultation record.

Materials and Methods

Type of Study	Retrospective descriptive analytical
Object of Study	Preanaesthetic consultation records of adult patients proposed for elective surgery Evaluation period: November 2016 to November 2017
Evaluated quality criteria (based on standard 029/2013 of DGS ¹)	<ol style="list-style-type: none"> Existence of registration (on specific anaesthetic evaluation form or in consultation diary, free text) Complete clinical evaluation (pathology, medication, allergies, habits) Complete objective examination (cardiopulmonary auscultation, weight/height) Complete airway assessment (Mallampati, dentition, neck, thyromentonian distance) Calculated clinical risks (physical status - ASA, Apfel scale, cardiac risk associated with the type of surgery) Defined anaesthetic plan and conclusion
Statistical analysis	SPSS 24.0® Descriptive: frequencies and percentages Analytical: chi-squared test

Results

Sample n = 653 consultations
607 (93%) – 1st consultation

647 (99,1%) – with consultation record
6 (0,9%) – without consultation record

Chart 1. Type of preanaesthetic evaluation record

Chart 2. Compliance rate of quality criteria

Chart 3. Quality criteria with least compliance (failure rates)

Table 1. Comparison of compliance rates between the two registration methods: specific anaesthetic evaluation form and consultation diary (chi-squared test)

Quality criteria	Specific anaesthetic evaluation form vs Consultation diary
Clinical evaluation	p < 0,001*
Objective examination	p < 0,001*
Airway assessment	p = 0,819
Calculated clinical risks	p < 0,001*
Anaesthetic plan/conclusion	p < 0,001*

Discussion and Conclusions

The **calculated clinical risk** was the criteria with the **highest failure rate**. Only 17,6% of consultations contained registry of **calculated cardiac risk**.

When comparing **records made** in an **specific anaesthetic evaluation form** with those on a consultation diary, the specific form records achieved **better compliance rates** in most of the quality criteria evaluated in this study (p < 0,001), with better quality of registration.

1. Saúde, Direção Geral da. Avaliação Pré-Anestésica para Procedimentos Eletivos. s.l.: 09/2013 (atualizada a 24/04/2015), 2013. Norma