SALIVARY A-AMYLASE AS A MARKER FOR SALIVARY GLAND TOXICITY IN HEAD AND NECK CANCER PATIENTS RECEIVING RADIOTHERAPY/ CHEMORADIOTHERAPY

Maria Kouri¹, Ourania Nicolatou-Galitis¹, Athina Vadalouca², Vasilios Kouloulis³, Georgios Maropoulos⁴, Panagiotis Kanellopoulos⁴, Efthymios Kyrodimos⁵, Erofilia Papadopoulou¹, Emmanouil Vardas¹, Evangelos Galitis⁶, Miltiadis Trichas⁷, Anna Zigogianni⁸, Zoi Liakouli⁸, Amanda Psyrris⁹, Ioannis Kouvaris⁸

INTRODUCTION:

Painful oral mucositis (OM), salivary gland dysfunction and xerostomia are significant radiation-toxicity related complications in head and neck cancer (HNC) patients. Salivary A-amylase can be a good salivary gland function indicator.

PATIENTS & METHODS:

PARAMETERS		N=34	%
Gender	Male	27	79.4
	Female	7	20.6
Age (Mean)		65.12	
Radiotherapy	Postoperative	21	61.76
	Radical	13	20.31
Total Dose (Mean, Gy)		63.18	
Concomitant Chemotherapy		25	73.53
Tumor Site	Oral/ Oropharyngeal	16	47.06
	Laryngeal	7	20.59
	Parotid	4	11.76
	Other	7	20.59

A-AMYLASE EVALUATION:

2-Chloro-NPG3-Substrate-Photometric method.
Whole saliva a-amylase was assessed:
before radiotherapy , 5th-7th weeks.

ORAL MUCOSITIS GRADING (EORTC / RTOG criteria):

Grades: 1-4

ORAL PAIN AND XEROSTOMIA EVALUATION:

Verbal Analogue Scale (VAS). Scores: 0 – 10;
1 - 4: mild, 5 – 7: moderate, 8 – 10: severe.

RESULTS:

	Before radiotherapy	5 th -7 th week of radiotherapy	N=30 (%)			N(%)	p
AMYLASE (U/ml) Reduced	165.09	49.48	25 (83.3)	Mucositis	Mild	7(23.3)	<0.005
					Moderate/ Severe	23(76.7)	
				Pain	Mild	14(46.7)	>0.005
					Moderate/ Severe	16(53.3)	
				Xerostomia	Mild	11(36.7)	<0.005
					Moderate/ Severe	19(63.3)	
AMYLASE (U/ml) Increased	241.17	389.91	5 (16.7)	Mucositis	Mild / Moderate	5(100)	>0.005
					Severe	0	
				Pain	Mild	5(100)	
					Moderate/ Severe	0	
				Xerostomia	Mild	5(100)	
					Moderate/ Severe	0	

	Before radiotherapy	2 nd -4 th week of radiotherapy	N=8 (%)			N(%)	p
AMYLASE (U/ml) Reduced	154.08	29.09	5 (62.5)	Mucositis	Mild	1 (20)	Not Available
					Moderate/ Severe	4(80)	
				Pain	Mild	1(20)	
					Moderate/ Severe	4(80)	
				Xerostomia	Mild	0	
					Moderate/ Severe	5(100)	
AMYLASE (U/ml) Increased	12.62	267.45	3 (37.5)	Mucositis	Mild	2(66.7)	Not Available
					Moderate/ Severe	1(33.3)	
				Pain	Mild	2(66.7)	
					Moderate/ Severe	1 (33.3)	
				Xerostomia	Mild	2(66.7)	
					Moderate/ Severe	1 (33.3)	

CONCLUSIONS:

The reduction of α-amylase in most assessments (30/38) was considered related to salivary gland dysfunction and radiation-toxicity. A-amylase may serve as a marker for the radiation-induced salivary gland toxicity.

2018
28-30 JUNE
VIENNA, AUSTRIA

**SUPPORTIVE CARE
MAKES EXCELLENT
CANCER CARE POSSIBLE**

**HELLENIC REPUBLIC
UNIVERSITY OF ATHENS**

**SCHOOL OF DENTISTRY
CLINIC OF HOSPITAL DENTISTRY
Head Professor :
Prof. O. NICOLATOU-GALITIS**