

Chapter

Behavioral Effects of Vitamin D₃ at Estrogen Deficiency in Females of Different Age

Julia Fedotova

Abstract

The present study was performed to determine the behavioral effects of cholecalciferol treatment at different doses as an adjunctive therapy alone or in a combination with low dose of 17 β -estradiol on anxiety-like behavior of the middle-aged (12–14 months) and old (16–18 months) female rats at 12 weeks after ovariectomy. Vitamin D₃ supplementation individually (as cholecalciferol at doses of 1.0, 2.5 or 5.0 mg/kg/day, s.c.) or in co-administration with 17 β -estradiol (17 β -E₂, 0.5 μ g/ rat, s.c.) were given to the ovariectomized (OVX) rats of different age at 12 weeks after ovariectomy. Anxiety-related state was tested in the elevated plus maze (EPM) and light-dark test (LDT), as well behavioral reactivity was registered in the open field test (OFT). The results of the study indicated that vitamin D₃ supplementation at doses of 1.0 or 5.0 mg/kg/day decreased manifestations of anxiety-like profile in the middle-aged or old OVX rats, respectively. Vitamin D₃ (1.0 mg/kg/day) plus 17 β -E₂ resulted in more profound anxiolytic-like effects in the old OVX rats than effects of both drugs administered alone. Cholecalciferol (1.0 mg/kg/day, s.c.) in the old OVX rats produced elevated estradiol and 25-OH-VD₃ levels for these rats as compared to the old OVX females treated with solvent.

Keywords: vitamin D₃, anxiety, menopause, estradiol, aging, female rats

1. Introduction

Anxiety disorders are twice common in women than in men and the risk increased more during the menopausal transition [1, 2]. Menopause indicates the termination of a woman's reproductive life. It is defined as the permanent cessation of menstruation induced by the loss of ovarian follicular activity [2, 3]. Menopause represents an important stage in female lives, which is often associated with a plethora of complaints and sufferings. Mood disturbances, especially anxiety and depression, are commonly associated with menopause [3, 4]. For women, data suggest that estrogens are strongly implicated in the regulation of mood and behavior, as well as in the pathophysiology of mood disorders [5–7]. Nowadays, there is a strong tendency to increase the duration of living in the whole world, however, the median age of programmed termination of a woman's reproductive life failed to alter [5–7]. Thus, majority of women in the aging population lives significant period of their lifetime in a postmenopausal state which is clearly associated with very low

estrogen levels that could be one of the marked trigger factors for development of affective-related disorders [5, 7].

A strategy to alleviate the mood disorders associated with menopause is hormonal replacement therapy (HRT) [8]. However, controversial results related to the effectiveness of such treatment have been frequently reported [9]. These discrepancies could be associated to various factors, one of them being the time when estrogen restitution is initiated after the beginning of menopause [10, 11].

There is a growing interest about the potential of diet and nutrients to improve the mental health of the women population and for the treatment of psychiatric disorders [10, 11]. In the case of mood disorders, the limitations of psychotropic drugs to achieve adequate rates of clinical remission and functional recovery have promoted the search for complementary approaches [12–15]. Menopausal women are now choosing to take alternate and complementary therapies marketed as “natural” treatments that offer the positive health effects of estrogens without the unwanted side effects [12, 13]. Among other nutraceuticals, one of such “natural” substances for treatment of affective-related diseases could be vitamin D (VD) [14, 15].

VD is a neuroactive secosteroid with well-known skeletal physiological role and diverse “non-skeletal” functions in the human body [16, 17]. “Non-skeletal” functions of VD are connected with its different range of outcomes in the central nervous system, such as neuroplasticity, apoptosis, cell proliferation and differentiation [18, 19]. All functions of VD in the body (classical functions, i.e., effect upon calcium-phosphate management and the non-classical ones) are imposed by the nuclear VD receptor (VDR), regulating directly the gene expression [20, 21]. Nuclear VDR are member of receptors family for transcription factors which are activated by numerous ligands [20, 21]. VDR are present in most tissues and cells in the body, and within the brain it shows some specificity to the prefrontal cortex, hippocampus, cingulate gyrus, thalamus, hypothalamus and substantia nigra [22]. This is of relevance as many of those brain regions have been implicated in the physiology of affective-related disorders.

Estrogen deficiency effects on affective-related behavior are restricted to certain periods of age after ovary removal [23, 24]. Preclinical data suggest that onset age of menopause can be important to obtain behavioral positive or negative results. Thus, it is of great interest to evaluate the effects of repeated cholecalciferol administration on anxiety-related behavior in the middle-aged and old female rats with long-term estrogen deficiency.

The aim of the present study was to determine if repeated systemic treatment with cholecalciferol affects anxiety-like behavior in the middle-aged and old female rats after long-term ovariectomy.

2. Materials and methods

2.1 Animals

Female albino Wistar rats (12–14 months, middle-aged rats or 16–18 months, old rats, weighing 230–240 or 260–270 g, respectively) from the special biocollection of Koltushi vivarium (St. Petersburg, Russia) were used in the present study. All rats were allocated in groups and were allowed to accommodate for 1 week in the animal house at I.P. Pavlov Institute of Physiology, of the Russian Academy of Sciences, before subjecting them to behavioral testing and pharmacological treatments. They were provided with a standard pellet diet and were given water *ad libitum*. The

animals were kept at a temperature of $23 \pm 2^\circ\text{C}$ and a 12 h light/dark cycle as well as a constant relative humidity ($50 \pm 10\%$) during all experimental sessions. Female rats of different age were randomly separated into experimental groups accordingly to their age, including the control groups.

Vitamin D₃ and 17 β -E₂ treatments, as well as anxiety-related tests were carried out by double-blind method by using rules of the Health guide for the care and use of Laboratory animals (1978) formulated by the National Institute of Health. Females of different age were placed in the special room for behavioral trials at least 1 h prior to the beginning of the experimental sessions which were performed from 09:00 am to 12:00 am. The experimental protocols of this study were approved by the Institutional Animal Ethics Committee of I.P. Pavlov Institute of Physiology, Russia (protocol 1095/1 from June 25, 2012).

2.2 Surgery

Long-term ovariectomy surgery was performed as previously described [25]. Briefly, middle-aged and old female rats were anesthetized with ketamine (70 mg/kg b.w.) mixed with xylazine (10 mg/kg b.w.). To avoid inflammation, the rats were administered with meloxicam (1 mg/kg b.w.). The fallopian tube was crushed and the ovary was removed by cutting. The effectiveness of long-term ovariectomy or 17 β -estradiol (17 β -E₂) application was assessed by vaginal smears. The ovariectomized (OVX) females of different age were housed in groups of five in cages separated by groups. To assure the long-term absence of estrogens, all rats after surgery were remained to the housing facilities for 12 weeks.

2.3 Drug treatments

17 β -estradiol, 17 β -E₂ (Sigma, USA) at low dose of 5.0 $\mu\text{g}/\text{rat}$ [7, 26] and vitamin D₃ as cholecalciferol (Sigma, USA) at several doses (1.0, 2.5 or 5.0 mg/kg) [27] were subcutaneously (s.c.) administered once daily starting 14 days prior to the cognitive experiments. 17 β -E₂ was dissolved in sterile sesame oil, VD₃ was dissolved in 95% ethanol solvent, aliquoted and stored at -80°C . The stock of VD₃ was dissolved in sterile water, resulting in a solution of cholecalciferol with 2% ethanol. All drug solutions were freshly prepared before each behavioral testing. 17 β -E₂ and cholecalciferol were injected in a volume of 0.1 ml. The middle-aged OVX females were 15.5–17.5 months middle-aged at the onset for drug treatments.

The estrogen, 17 β -E₂ (E-8875, Sigma Chemical Co., St. Louis, MO, USA) was dissolved in sterile sesame oil. Cholecalciferol (C-9756, Sigma Chemical Co., St. Louis, MO, USA) was dissolved in 95% ethanol, aliquoted and stored at -80°C . The stock of cholecalciferol was diluted in sterile water, resulting in a solution of cholecalciferol with 2% ethanol. 17 β -E₂ was injected subcutaneously (s.c. at a dose of 0.5 $\mu\text{g}/\text{rat}$). The low dose of 17 β -E₂ (5.0 $\mu\text{g}/\text{rat}$ subcutaneously, s.c.) was chosen from the studies performed by Estrada-Camarena and co-workers [26, 28]. Three doses of cholecalciferol (1.0, 2.5 or 5.0 mg/kg, s.c.) were chosen from the behavioral study performed by Idrus and co-workers [27]. All solutions were freshly prepared before each experimental series. All preparations were administered in a volume of 0.1 ml. Following 12 weeks after ovariectomy, cholecalciferol, 17 β -E₂ and oil solvent were injected once daily for 14 days.

2.4 Animal groups

Female rats (middle-aged or old, intact and OVX) were randomly divided into 24 groups, accordingly to their age, with 8 rats in each group.

The following experimental groups for the middle-aged and old female rats were created in the present study:

- 1 and 2—middle-aged or old intact female rats + solvent.
- 3 and 4—middle-aged or old intact female rats + cholecalciferol 1.0 mg/kg (middle-aged or old intact – vitamin D₃ 1.0).
- 5 and 6—middle-aged or old intact female rats + cholecalciferol 2.5 mg/kg (middle-aged or old intact – vitamin D₃ 2.5).
- 7 and 8—middle-aged or old intact female rats + cholecalciferol 5.0 mg/kg (middle-aged or old intact – vitamin D₃ 5.0).
- 9 and 10—middle-aged or old OVX + solvent (middle-aged or old OVX – Sol).
- 11 and 12—middle-aged or old OVX rats + 17β-E₂ (middle-aged or old OVX – 17β-E₂).
- 13 and 14—middle-aged or old OVX rats + cholecalciferol 1.0 mg/kg (middle-aged or old OVX – vitamin D₃ 1.0).
- 15 and 16—middle-aged or old OVX rats + cholecalciferol 2.5 mg/kg (middle-aged or old OVX – vitamin D₃ 2.5).
- 17 and 18—middle-aged or old OVX rats + vitamin D₃ 5.0 mg/kg (middle-aged or old OVX – vitamin D₃ 5.0).
- 19 and 20—middle-aged or old OVX rats + cholecalciferol 1.0 mg/kg + 17β-E₂ (middle-aged or old OVX – vitamin D₃ 1.0 – 17β-E₂).
- 21 and 22—middle-aged or old OVX rats + cholecalciferol 2.5 mg/kg + 17β-E₂ (middle-aged or old OVX – vitamin D₃ 2.5 – 17β-E₂).
- 23 and 24—middle-aged or old OVX rats + cholecalciferol 5.0 mg/kg + 17β-E₂ (middle-aged or old OVX – vitamin D₃ 5.0 – 17β-E₂).

The treatment period for animals was 14 days, and at the end of the treatment period (1 h after the last dose of solvent, vitamin D₃ or 17β-E₂), all animals were subjected to the elevated plus maze (EPM), light-dark test (LDT) and the open field test. During testing sessions in all behavioral tests, the control and experimental groups of rats were also given with solvent, vitamin D₃ or 17β-E₂.

2.5 Behavioral tests

2.5.1 Elevated plus maze test

EPM is commonly accepted as standard test of anxiety-like behavior and was used to assess anxiety-like behavioral responses [29, 30]. This test is sensitive to putative anxiogenic-like and anxiolytic-like drugs [31]. EPM consist of two open arms (50 × 10 cm²) and two closed arms (40 × 10 cm²) with a central platform (10 × 10 cm²) and elevated 50 cm above the floor level. All female rats from control and experimental groups were randomly placed at the center of the EPM and allowed them to freely move in the apparatus for 5 min. The number of entries and total time spent in open arms were accepted as parameters of anxiolytic-like effects of treatments. The apparatus was cleaned with damp cloth after each trial to avoid place preference and the influence of olfactory stimuli.

2.5.2 Light/dark test

The apparatus consist of two identical boxes (30 × 40 × 40 cm), one of which with white walls and floor and illuminated by a 60 Watt light from above, while the other of the box was painted black and had a lid so it was not illuminated [32, 33]. The number of entrance and the total time in the light box were registered for 5 min [7]. The increase in the number of entrances and the total time in the light box were postulated as manifestation of anxiolytic-like effects of treatments. The apparatus was cleaned with damp cloth after each trial to avoid place preference and the influence of olfactory stimuli.

2.5.3 Open field test

The effect of cholecalciferol on locomotor, rearing and grooming activities was evaluated automatically using an open-field computer-aided controlling system as described previously [34]. The apparatus consists of a square platform (80.0 cm × 80.0 cm; wall height 36.0 cm). The floor of the platform was divided into 16 equal squares of 19.5 cm × 19.5 cm. A video camera fixed at the top, and the apparatus was illuminated by a light source of 120 Lux on the ceiling. Each rat was placed at the center of the apparatus and allowed to explore freely for 5 min. Total number of central and peripheral square crossings were recorded for each animal. The apparatus was cleaned with damp cloth after each trial to avoid place preference and the influence of olfactory stimuli.

2.6 Determination of estradiol, 25-OH-VD₃ and calcium levels in the blood serum

Blood samples were collected in tubes and centrifuged. After centrifugation, serum was separated, frozen and stored at -20°C until biochemical assessment. Estradiol levels were assessed using commercial available ELISA kit (DRG Diagnostics, Marburg, Germany). The sensitivity of the estradiol ELISA kit was 1.0 pg/ml. Measurement of 25-hydroxyvitamin D₃ (25-OH-VD₃) levels was performed by ELISA kit (CSB-E08098r, Cusabio Biotech Co., Ltd., Wuhan, P.R. China). Technical variability for 25-OH-VD₃ ELISA kit was low with coefficients of variation of <10% intra-assay and < 15% inter-assay. Detection range of 25-OH-VD₃ levels was 20–100 µg/L. The sensitivity of the 25-OH-VD₃ using ELISA kit was 5.0 µg/L. Calcium concentrations were detected by spectrophotometric method using calcium assay colorimetric kit (ab102505, Abcam, France). The sensitivity of the calcium kit was 0.1 m/M. All the procedures of estradiol, 25-OH-VD₃ and calcium kits were conducted following the manufacturer's instruction manual.

2.7 Statistical analysis

Data were expressed as means ± standard error (SEM). Differences among means were postulated as significant at $p \leq 0.05$. Behavioral and biochemical data were analyzed using a two-way ANOVA and subsequent *post-hoc* analysis was conducted with Dunnett's multiple comparison test. Statistical calculation was carried out using SPSS software 19 version (SPSS Inc., Chicago, IL., USA).

3. Results

3.1 Vitamin D₃ in different doses decreases anxiety-like profile of the middle-aged and old OVX and OVX rats given with 17β-estradiol after long-term absence of estrogen as measured in the EPM test

For vitamin D₃ supplementation, two-way ANOVA analysis revealed a significant interaction between hormone condition and treatments ($[F(5,44) = 12.83, p < 0.05]$ and $[F(5,44) = 9.47, p < 0.01]$, respectively), with significant effects of hormone conditions ($[F(5,44) = 9.47, p < 0.01]$ and $[F(5,44) = 7.88, p < 0.01]$, respectively) and treatment ($[F(5,44) = 15.24, p < 0.05]$ and $[F(5,44) = 11.02, p < 0.05]$, respectively) in the time spent into the open arms or the number of entries into the open arms of the middle-aged OVX rats. The *post-hoc* test demonstrated significant differences for these groups ($p < 0.05$).

Vitamin D₃ application at all tested doses did not modify the time spent into the open arms and the number of entries into the open arms in the intact middle-aged rats as compared to the control rats (**Figure 1a, b**, $p > 0.05$). Following 12 weeks of post-surgery period, the middle-aged female rats showed a profound decrease of the time spent in the open arms and the number of entries into the open arms as compared to the control females (**Figure 1a, b**, $p < 0.05$). The 17 β -E₂ injection (0.5 μ g/kg, s.c.) resulted in an increase of the time spent in the open arms and the number of entries into the open arms in the middle-aged OVX rats as compared to the middle-aged OVX rats administered with solvent (**Figure 1a, b**, $p < 0.05$), but did not reach the values of control rats (**Figure 1a, b**, $p < 0.05$).

Vitamin D₃ treatment at dose of 5.0 mg/kg significantly increased the time spent in the open arms and the number of entries into the open arms of the middle-aged OVX rats as compared to the middle-aged OVX rats given with solvent (**Figure 1a, b**, $p > 0.05$). Co-administration of vitamin D₃ at dose of 5.0 mg/kg and 17 β -E₂ to the middle-aged OVX rats produced a more greater increase of the time spent in the open arms and the number of entries into the open arms than that of the middle-aged OVX rats given 17 β -E₂ or solvent (**Figure 1a, b**, $p > 0.05$). However, vitamin D₃ treatment at doses of 1.0 and 2.5 mg/kg individually or plus 17 β -E₂ failed to alter the time spent in the open arms and the number of entries




Figure 1. Effects of cholecalciferol administration on anxiety-like behavior of the middle-aged ovariectomized (OVX) rats following long-term estrogen deficiency in the elevated plus maze. (a) Time spent into the open arms, sec; (b) The number of entries into the open arms. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol (17 β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17 β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.

into the open arms of the middle-aged OVX as compared to the middle-aged OVX rats treated with 17 β -E₂ **Figure 1a, b**, $p > 0.05$). The time spent in the open arms and the number of entries into the open arms in the middle-aged OVX rats administered with vitamin D₃ at these doses in combination with 17 β -E₂ were lower than that for middle-aged control rats and were higher than for middle-aged OVX rats (**Figure 1a, b**, $p > 0.05$).

Two-way ANOVA test demonstrated a significant hormone condition \times treatment interaction ($[F(5,44) = 11.44, p < 0.05]$ and $[F(5,44) = 11.12, p < 0.01]$, respectively), significant effect of hormone conditions ($[F(5,44) = 9.22, p < 0.01]$ and $[F(5,44) = 16.88, p < 0.01]$, respectively) and significant effect for treatment ($[F(5,44) = 11.56, p < 0.05]$, and $[F(5,44) = 12.56, p < 0.05]$, respectively) for the time spent into the open arms or the number of entries into the open arms of the old OVX rats. *Post-hoc* analyses revealed differences among the groups for the anxiety-like state during experimental sessions ($p < 0.05$).

The old intact rats treated with vitamin D₃ at doses of 1.0, 2.5 or 5.0 mg/kg failed to alter the time spent into the open arms and the number of entries into the open arms as compared to the old control rats (**Figure 2a, b**, $p > 0.05$). The old OVX rats given with solvent displayed a significant decrease of the time spent into the open arms and the number of entries into the open arms as compared to the old control rats (**Figure 2a, b**, $p > 0.05$).




Figure 2.

Effects of cholecalciferol administration on anxiety-like behavior of the old ovariectomized (OVX) rats following long-term estrogen deficiency in the elevated plus maze. (a) Time spent into the open arms, sec; (b) The number of entries into the open arms. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol (17 β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17 β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.

Vitamin D₃ supplementation administered in all doses to the old OVX rats resulted in increase of the time spent into the open arms and the number of entries into the open arms as compared to the old OVX rats treated with solvent (**Figure 2a, b**, $p > 0.05$). Combined administration of cholecalciferol at dose of 1.0 mg/kg with 17 β -E₂ to the old OVX rats more significantly increased the time spent into the open arms and the number of entries into the open arms as compared to the old OVX rats treated with solvent or 17 β -E₂ (**Figure 2a, b**, $p > 0.05$). The old OVX rats administered with cholecalciferol at doses of 2.5 mg/kg and 5.0 mg/kg plus 17 β -E₂ showed similar values of the time spent into the open arms and the number of entries into the open arms like as old OVX rats treated with 17 β -E₂ (**Figure 2a, b**, $p > 0.05$).

3.2 Vitamin D₃ at different doses reverses anxiety-like profile of the middle-aged and old OVX and OVX rats given with 17 β -estradiol after long-term absence of estrogen as measured in light: dark box

Two-way ANOVA statistical test revealed a significant interaction between hormone condition and treatments ([F(5,44) = 11.52, $p < 0.01$) and [F(5,44) = 16.75, $p < 0.05$], respectively), with significant effects of hormone conditions ([F(5,44) = 19.22, $p < 0.05$ and [F(5,44) = 9.56, $p < 0.001$], respectively) and treatment ([F(5,44) = 9.88, $p < 0.01$) and [F(5,32) = 7.26, $p < 0.05$], respectively) in the time spent and number of entries in the light compartment of the middle-aged OVX rats. The *post-hoc* test demonstrated significant differences for these groups ($p < 0.05$).

Vitamin D₃ supplementation at all doses did not alter the time spent and number of entries in the light compartment in the intact middle-aged rats as compared to the control rats (**Figure 3a, b**, $p > 0.05$). Following 12 weeks of post-ovariectomy period, the middle-aged female rats showed a profound decrease of the time spent and number of entries in the light compartment as compared to the control females (**Figure 3a, b**, $p < 0.05$). The 17 β -E₂ injection (0.5 μ g/kg, s.c.) resulted in an increase of the time spent and number of entries in the light compartment in the middle-aged OVX rats as compared to the middle-aged OVX rats administered with solvent (**Figure 3a, b**, $p < 0.05$), but did not reach the values of control rats (**Figure 3a, b**, $p < 0.05$).

Vitamin D₃ treatment at dose of 5.0 mg/kg markedly elevated time spent and number of entries in the light compartment of the middle-aged OVX rats as compared to the middle-aged OVX rats given with solvent (**Figure 3a, b**, $p > 0.05$). Co-administration of vitamin D₃ at dose of 5.0 mg/kg and 17 β -E₂ to the middle-aged OVX rats produced greater increase of time spent and number of entries in the light compartment than that of the middle-aged OVX rats given 17 β -E₂ or solvent (**Figure 3a, b**, $p > 0.05$). Vitamin D₃ treatment at doses of 1.0 mg/kg and 2.5 mg/kg individually or plus 17 β -E₂ did not change time spent and number of entries in the light compartment of the middle-aged OVX as compared to the middle-aged OVX rats treated with 17 β -E₂ (**Figure 3a, b**, $p > 0.05$). The time spent and number of entries in the light compartment in the middle-aged OVX rats administered vitamin D₃ at these doses in combination with 17 β -E₂ were lower than that for middle-aged control rats and were higher than for middle-aged OVX rats (**Figure 3a, b**, $p > 0.05$).

Two-way analysis of variance showed a significant hormone condition \times treatment interaction ([F(5,44) = 11.52, $p < 0.01$) and [F(5,44) = 12.74, $p < 0.05$], respectively), significant effect of hormone conditions ([F(5,44) = 11.22, $p < 0.05$) and [F(5,44) = 11.56, $p < 0.001$], respectively) and significant effect for treatment ([F(5,44) = 10.08, $p < 0.01$) and [F(5,32) = 12.26, $p < 0.05$], respectively) for time spent and number of entries in the light compartment of the old OVX rats. *Post-hoc* analyses revealed differences among the groups for the anxiety-like state during experimental sessions ($p < 0.05$).




Figure 3. Effects of cholecalciferol administration on anxiety-like behavior of the middle-aged OVX rats following long-term estrogen deficiency in the light/dark test. (a) Time spent in the light box, sec; (b) The number of entrances in the light box. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of sham-operated rats, #— $p < 0.05$ as compared to the old OVX rats treated with solvent, ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol. Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day s.c., once daily, for 14 days. The administered dose of 17 β -estradiol (17 β -E₂) was 0.5 μ g/rat s.c., once daily, for 14 days.

The old intact rats treated with vitamin D₃ at doses of 1.0, 2.5 or 5.0 mg/kg failed to modify time spent and number of entries in the light compartment as compared to the old control rats (Figure 4a, b, $p > 0.05$). The old OVX rats given with solvent displayed a significant decrease of time spent and number of entries in the light compartment as compared to the old control rats (Figure 4a, b, $p > 0.05$).

Vitamin D₃ supplementation administered in all doses to the old OVX rats induced increase of the time spent and number of entries in the light compartment as compared to the old OVX rats treated with solvent (Figure 4a, b, $p > 0.05$). Combined administration of cholecalciferol at dose of 1.0 mg/kg with 17 β -E₂ to the old OVX rats more significantly increased time spent and number of entries in the light compartment as compared to the old OVX rats treated with solvent or 17 β -E₂ (Figure 4a, b, $p > 0.05$). The old OVX rats administered with vitamin D₃ at doses of 2.5 mg/kg and 5.0 mg/kg plus 17 β -E₂ showed identical parameters of time spent and number of entries in the light compartment like as old OVX rats treated with 17 β -E₂ (Figure 4a, b, $p > 0.05$).

3.3 Vitamin D₃ administration changes behavioral reactivity of the middle-aged and old OVX and OVX rats treated with 17 β -estradiol

Accordingly to the two-way ANOVA test, there were significant differences for the grooming behavior between hormone conditions ($[F(5,44) = 8.12, p < 0.05]$ and $[F(5,44) = 9.22, p < 0.05]$, respectively) between drug treatment ($[F(5,44) = 12.51,$




Figure 4.

Effects of cholecalciferol administration on anxiety-like behavior of the old OVX rats following long-term estrogen deficiency in the light/dark test. (a) Time spent in the light box, sec; (b) The number of entrances in the light box. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of sham-operated rats, #— $p < 0.05$ as compared to the old OVX rats treated with solvent, ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol. Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day s.c., once daily, for 14 days. The administered dose of 17 β -estradiol (17 β -E₂) was 0.5 μ g/rat s.c., once daily, for 14 days.

$p < 0.01$] and [F(5,44) = 12.56, $p < 0.01$], respectively) and an interaction between hormone condition and treatments ([F(5,44) = 7.16, $p < 0.01$] and [F(5,44) = 9.26, $p < 0.01$], respectively) in the middle-aged and old OVX rats. Further *post-hoc* test revealed differences for grooming between experimental groups of the OVX rats with different age ($p < 0.05$).

Vitamin D₃ injected at several doses failed to demonstrate any changes of behavioral reactivity of the middle-aged and old intact females in the OFT as compared to the middle-aged control rats (Tables 1 and 2, $p > 0.05$).

A significant decrease of grooming behavior was registered in the middle-aged and old OVX rats given with solvent as compared to the control (Tables 1 and 2, $p < 0.05$). 17 β -E₂ significantly reduced grooming reactions in the middle-aged and old OVX rats as compared to the middle-aged OVX rats (Tables 1 and 2, $p < 0.05$). The middle-aged and old OVX rats treated with vitamin D₃ in all tested doses alone or in a combination with 17 β -E₂ did not demonstrate any modifications of motor and rearing activities as compared to the middle-aged OVX rats given with solvent (Tables 1 and 2, $p < 0.05$).

However, the middle-aged and old OVX rats treated with vitamin D₃ at doses of 1.0, 2.5 and 5.0 mg/kg demonstrated an increase of grooming behavior as compared to the middle-aged OVX rats. A co-administration of vitamin D₃ at these doses with 17 β -E₂ decreased grooming behavior as compared to both middle-aged, as well as old intact and OVX rats received with solvent or 17 β -E₂ (Tables 1 and 2, $p < 0.05$).

3.4 Modifications of 25-hydroxyvitamin D₃, estradiol and calcium levels in the blood serum following vitamin D₃ administration in the middle-aged and old OVX and OVX females treated with 17 β -estradiol

The middle-aged intact rats treated with cholecalciferol at doses of 1.0, 2.5 and 5.0 mg/kg increased 25-OH-VD₃ levels (**Figure 5**, $p < 0.05$) and failed to alter estradiol levels in the serum blood as compared to the control rats (**Figure 6**, $p > 0.05$).

Long-term ovariectomy in the middle-aged female rats resulted in a significant decrease of estradiol and 25-OH-VD₃ levels in the blood as compared to the middle-aged control females (**Figures 5 and 6**, $p < 0.05$). The 17 β -E₂ supplementation (0.5 μ g/kg, SC) failed to modify 25-OH-VD₃ levels in the blood of the middle-aged OVX rats as compared to the middle-aged OVX rats administered with solvent (**Figure 5**, $p > 0.05$), and the value of this parameter in the middle-aged OVX/17 β -E₂ females were lower than that of the value of middle-aged control rats. However, 17 β -E₂ supplementation significantly increased estradiol levels in the blood of the middle-aged OVX rats as compared to the middle-aged OVX rats given with solvent (**Figure 6**, $p < 0.05$).

The middle-aged OVX rats treated with cholecalciferol at all tested doses significantly increased estradiol levels in the serum blood as compared to the middle-aged OVX rats treated with solvent (**Figure 6**, $p < 0.05$). However, the value of estradiol levels in the middle-aged OVX rats treated with cholecalciferol at these doses were lower than that of the value of middle-aged control rats. The middle-aged OVX rats treated with cholecalciferol at doses of 2.5 and 5.0 mg/kg significantly increased 25-OH-VD₃ levels in the serum blood as compared to the middle-aged OVX rats

Groups	Crossing	Rearing	Grooming
Middle-aged control rats + solvent	73.3 \pm 4.2	12.1 \pm 0.8	3.0 \pm 0.2
Middle-aged intact rats + cholecalciferol 1.0 mg/kg	68.0 \pm 2.4	12.0 \pm .05	3.2 \pm 0.2
Middle-aged intact rats + cholecalciferol 2.5 mg/kg	59.3 \pm 5.6	10.5 \pm 0.8	3.2 \pm 0.2
Middle-aged intact rats + cholecalciferol 5.0 mg/kg	69.5 \pm 4.2	11.7 \pm 0.8	3.5 \pm 0.2
Middle-aged OVX rats + solvent (OVX/solvent rats)	62.4 \pm 2.3	12.3 \pm 0.6	1.2 \pm 0.5*
Middle-aged OVX rats +17 β -E ₂ (OVX/17 β -E ₂ rats)	60.3 \pm 2.6	13.2 \pm 0.3	3.3 \pm 0.4#
Middle-aged OVX rats + cholecalciferol 1.0 mg/kg	65.2 \pm 2.5	12.1 \pm 0.6	4.0 \pm 0.2#
Middle-aged OVX rats + cholecalciferol 2.5 mg/kg	76.9 \pm 4.2	13.2 \pm 0.8	3.8 \pm 0.4#
Middle-aged OVX rats + cholecalciferol 5.0 mg/kg	72.3 \pm 4.4	11.0 \pm 0.5	4.1 \pm 0.6#
Middle-aged OVX rats + cholecalciferol 1.0 mg/kg + 17 β -E ₂	62.1 \pm 2.8	12.6 \pm 0.6	0.6 \pm 0.2*,#,#
Middle-aged OVX rats + cholecalciferol 2.5 mg/kg + 17 β -E ₂	73.2 \pm 2.4	10.2 \pm 0.4	0.5 \pm 0.2*,#,#
Middle-aged OVX rats + cholecalciferol 5.0 mg/kg + 17 β -E ₂	65.4 \pm 5.6	12.5 \pm 0.8	0.8 \pm 0.2*,#,#

* $p < 0.05$ as compared to the control group of the old sham-operated rats.

$p < 0.05$ as compared to the old OVX rats treated with solvent.

$p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol.

The obtained results show the mean \pm S.E.M. Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day, s.c., once daily, for 14 days. 17 β -Estradiol (17 β -E₂) was given at 0.5 μ g/rat, s.c., once daily, during 14 days.

Table 1.

Cholecalciferol influences on behavioral parameters of the middle-aged OVX rats following long-term estrogen deficiency in the open field test for 5 min.

Groups	Crossing	Rearing	Grooming
Old control rats + solvent	69.7 ± 5.2	11.5 ± 0.3	2.9 ± 0.2
Old intact rats + cholecalciferol 1.0 mg/kg	71.8 ± 2.9	10.7 ± 0.3	3.2 ± 0.2
Old intact rats + cholecalciferol 2.5 mg/kg	66.9 ± 3.6	10.4 ± 0.2	3.1 ± 0.2
Old intact rats + cholecalciferol 5.0 mg/kg	69.0 ± 4.2	12.2 ± 0.8	3.0 ± 0.2
Old OVX rats + solvent (OVX/solvent rats)	72.1 ± 2.3	12.1 ± 0.6	1.0 ± 0.2 [*]
Old OVX rats + 17β-E ₂ (OVX/17β-E ₂ rats)	64.3 ± 4.6	11.7 ± 0.8	3.1 ± 0.3 [#]
Old OVX rats + cholecalciferol 1.0 mg/kg	63.2 ± 3.5	12.6 ± 0.9	4.2 ± 0.2 [#]
Old OVX rats + cholecalciferol 2.5 mg/kg	67.2 ± 5.2	10.2 ± 0.8	3.9 ± 0.2 [#]
Old OVX rats + cholecalciferol 5.0 mg/kg	70.3 ± 4.4	11.5 ± 0.5	4.3 ± 0.2 [#]
Old OVX rats + cholecalciferol 1.0 mg/kg + 17β-E ₂	72.1 ± 6.8	12.2 ± 0.6	0.7 ± 0.2 ^{*,##,##}
Old OVX rats + cholecalciferol 2.5 mg/kg + 17β-E ₂	78.5 ± 8.4	11.8 ± 0.4	0.6 ± 0.2 ^{*,##,##}
Old OVX rats + cholecalciferol 5.0 mg/kg + 17β-E ₂	69.4 ± 6.6	10.9 ± 0.8	0.9 ± 0.2 ^{*,##,##}

^{*} *p* < 0.05 as compared to the control group of the old sham-operated rats.

[#] *p* < 0.05 as compared to the old OVX rats treated with solvent.

^{##} *p* < 0.05 as compared to the old OVX rats treated with 17β-estradiol.

The obtained results show the mean ± S.E.M. Each group comprised a minimum of 8 rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day, s.c., once daily, for 14 days. 17β-Estradiol (17β-E₂) was given at 0.5 µg/rat, s.c., once daily, during 14 days.

Table 2.

Effects of cholecalciferol administration on behavioral impairments of the old OVX rats following long-term estrogen deficiency in the open field test for 5 min.




Figure 5.

Effects of cholecalciferol administration on 25-OH-VD₃ level of the middle-aged ovariectomized (OVX) rats following long-term estrogen deficiency in the serum blood. The obtained results show the mean ± standard error of the mean (SEM). *—*p* < 0.05 as compared to the control group of the old sham-operated rats; #—*p* < 0.05 as compared to the old OVX rats treated with solvent; ##—*p* < 0.05 as compared to the old OVX rats treated with 17β-estradiol (17β-E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17β-estradiol was 0.5 µg/rat s.c., once daily, for 14 days.

treated with solvent (**Figure 5**, *p* < 0.05). The value of 25-OH-VD₃ content in the middle-aged OVX rats treated with cholecalciferol at doses of 2.5 and 5.0 mg/kg were lower than that of the value of middle-aged control rats. Moreover, cholecalciferol administered at a dose of 1.0 mg/kg into the middle-aged OVX rats failed to change 25-OH-VD₃ levels in the serum blood as compared to the middle-aged OVX rats treated with solvent (**Figure 5**, *p* > 0.05). Cholecalciferol treatment at doses of




Figure 6. Effects of cholecalciferol administration on estradiol level of the middle-aged ovariectomized (OVX) rats following long-term estrogen deficiency in the serum blood. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol (17 β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17 β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.

2.5 and 5.0 mg/kg in combination with 17 β -E₂ more significantly elevated 25-OH-VD₃ levels for the middle-aged OVX rats as compared to the OVX females treated with oil solvent or 17 β -E₂ (Figure 3, $p < 0.05$). Combined administration of cholecalciferol at a dose of 1.0 mg/kg and 17 β -E₂ in the middle-aged OVX rats failed to change 25-OH-VD₃ levels as compared to the OVX rats administered with 17 β -E₂ or solvent (Figure 5, $p > 0.05$). Cholecalciferol at all doses in combination with 17 β -E₂ significantly increased estradiol levels when middle-aged OVX rats/cholecalciferol in tested doses plus 17 β -E₂ rats were compared with the middle-aged OVX/solvent and OVX/17 β -E₂ rat groups (Figure 6, $p > 0.05$).

The old intact rats treated with cholecalciferol at doses of 1.0, 2.5 and 5.0 mg/kg increased 25-OH-VD₃ levels (Figure 7, $p < 0.05$) and failed to alter estradiol levels in the serum blood as compared to the control rats (Figure 8, $p > 0.05$). Long-term ovariectomy in the old female rats resulted in a significant decrease of estradiol and 25-OH-VD₃ levels in the blood as compared to the old control females (Figures 7 and 8, $p < 0.05$). The 17 β -E₂ supplementation (0.5 μ g/kg, s.c.) failed to modify




Figure 7. Effects of cholecalciferol administration on 25-OH-VD₃ level of the old ovariectomized (OVX) rats following long-term estrogen deficiency in the serum blood. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol (17 β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17 β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.




Figure 8. Effects of cholecalciferol administration on estradiol level of the old ovariectomized (OVX) rats following long-term estrogen deficiency in the serum blood. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ###— $p < 0.05$ as compared to the old OVX rats treated with 17β -estradiol (17β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.

25-OH-VD₃ levels in the blood of the old OVX rats as compared to the old OVX rats administered with solvent (Figure 7, $p > 0.05$), and the value of this parameter in the old OVX/ 17β -E₂ females were lower than that of the value of old control rats. However, 17β -E₂ supplementation significantly increased estradiol levels in the blood of the old OVX rats as compared to the old OVX rats given with solvent (Figure 8, $p < 0.05$).

The old OVX rats treated with cholecalciferol at all doses significantly increased 25-OH-VD₃ and estradiol levels in the serum blood as compared to the old OVX rats treated with solvent (Figure 8, $p < 0.05$). However, the values of 25-OH-VD₃ and estradiol levels in the old OVX rats treated with cholecalciferol at all doses were lower than that of the values of old control rats.

Co-administration of vitamin D₃ (1.0 mg/kg) and 17β -E₂ markedly enhanced estradiol levels in the old OVX rats as compared to the groups of old OVX rats received with solvent or 17β -E₂ (Figure 7, $p < 0.05$). Vitamin D₃ supplementation (2.5 and 5.0 mg/kg) plus 17β -E₂ did not modify estradiol concentrations in the serum blood of the old OVX rats as compared to the OVX rats given with 17β -E₂




Figure 9. Effects of cholecalciferol administration on calcium level of the middle-aged ovariectomized (OVX) rats following long-term estrogen deficiency in the serum blood. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ###— $p < 0.05$ as compared to the old OVX rats treated with 17β -estradiol (17β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.




Figure 10. Effects of cholecalciferol administration on calcium level of the old ovariectomized (OVX) rats following long-term estrogen deficiency in the serum blood. The obtained results show the mean \pm standard error of the mean (SEM). *— $p < 0.05$ as compared to the control group of the old sham-operated rats; #— $p < 0.05$ as compared to the old OVX rats treated with solvent; ##— $p < 0.05$ as compared to the old OVX rats treated with 17 β -estradiol (17 β -E₂). Each group comprised a minimum of eight rats. Cholecalciferol was given at 1.0, 2.5 or 5.0 mg/kg/day subcutaneously (s.c.), once daily, for 14 days. The administered dose of 17 β -estradiol was 0.5 μ g/rat s.c., once daily, for 14 days.

(Figure 7, $p > 0.05$). Cholecalciferol at all doses in combination with 17 β -E₂ significantly increased 25-OH-VD₃ levels when these old OVX rats were compared with the old OVX/solvent and OVX/17 β -E₂ groups (Figure 8, $p > 0.05$).

The two-way ANOVA failed to show any significant differences in the calcium levels in the blood serum between hormone conditions, drug treatments, and an interaction between hormone condition and treatments in the middle-aged and old OVX rats with long-term estrogen deficiency (Figures 9 and 10, $p > 0.05$). The *post-hoc* test did not find any differences among the experimental groups for the calcium levels ($p > 0.05$).

4. Discussion

In the present study, the effects of chronic cholecalciferol treatment at different doses (1.0, 2.5 and 5.0 mg/kg, s.c.) for 14 days on anxiety-like behavior in the middle-aged and old female rats with long-term estrogen deficiency and 17 β -E₂ supplementation in a low dose were examined. Endogenous estrogens were removed by ovariectomy and only after 12 weeks post-ovariectomy period, these rats were used in all experiments. The results of behavioral testing for the anxiety-related effects of cholecalciferol were compared in both old OVX rats and OVX female rats treated with 17 β -E₂. Simultaneously, the effects of cholecalciferol at similar doses on anxiety-like behavior were tested in middle-aged and old intact female rats. For this purpose, the elevated plus maze (EPM) and light-dark test (LDT) were made use in the present study. It was also investigated whether the effects of cholecalciferol at different doses were specific in the EPM and LDT, measuring its effects on the behavioral activity in the OFT of the middle-aged and old intact and OVX rats after long-term absence of estrogen.

Cholecalciferol at all investigated doses did not produce any changes of anxiety-like behavior of the middle-aged and old intact female rats in the EPM and LDT. Analyzing the results from biochemical assay, an increase of 25-OH-VD₃ concentrations and absence of any modifications of estradiol levels in the serum blood of the middle-aged and old intact rats given with different doses of cholecalciferol were found.

These results suggest that cholecalciferol induced the increasing of 25-OH-VD₃ levels in the blood serum of the middle-aged and old intact-ovary rats are not associated with absence of anxiety-like profile alterations in the behavioral tests.

Furthermore, the ovary-intact female rats of different age are also needed to evaluate the behavioral effects of cholecalciferol administered at several doses in the EPM and LDT paradigms.

The results showed that in the middle-aged and old OVX rats following 12 weeks of post-ovariectomy period, there were marked anxiety-like behavior as assessed by EPM and LDT. Although 17β -E₂ supplementation resulted in significant anxiolytic-like effect of the middle-aged and old OVX rats with long-term absence of estrogen, the 17β -E₂ administration was not able to completely diminish anxiety-like behavior to the level of the middle-aged and old control intact animals. According to these results, we conclude that middle-aged and old OVX rats following 12 weeks of postovariectomy period display significant anxiety-related behavior, while 17β -E₂ administration to the middle-aged and old OVX rats attenuates the estrogen deficiency-induced anxiety-like behavior to some extent. In fact, these experiments showed that the effects of 17β -E₂ supplementation on anxiety-like behavior did not associated with absence of its effects on 25-OH-VD₃ levels in the old OVX rats. The long-term effect of ovariectomy on anxiety-like behavior in female rats of different age that were submitted in a standard behavioral tests [24, 35, 36].

Cholecalciferol at dose of 5.0 mg/kg/day per se had a significant anxiolytic-like effect in the middle-aged OVX rats following long-term ovariectomy. On the contrary, cholecalciferol at doses of 1.0 and 2.5 mg/kg/day failed to induce any modifications of anxiety-like behavior in the middle-aged OVX rats with long-term absence of estrogen. Combined application of vitamin D₃ (1.0 mg/kg/day) with 17β -E₂-induced synergic anxiolytic-like action of both preparations in the anxiety-related tasks of the old OVX rats. Cholecalciferol at all doses per se had a significant anxiolytic-like effect in the old OVX rats following long-term ovariectomy. Unexpectedly, that the old OVX rats with 12 weeks post-ovariectomy period administered with cholecalciferol at doses 2.5 and 5.0 mg/kg/day in combination with 17β -E₂ showed similar anxiety-like profile like the OVX rats given with 17β -E₂. Thus, we did not observe any synergic anxiolytic-like effects of cholecalciferol at doses of 2.5 or 5.0 mg/kg in the old OVX rats given with 17β -E₂. It might suppose that there are some concurrent relation between 17β -E₂ and cholecalciferol at doses of 2.5 and 5.0 mg/kg/day. In fact, application of 17β -E₂ interfere with anxiolytic-like action of cholecalciferol at doses of 2.5 or 5.0 mg/kg in the old OVX rats after long-term ovariectomy. Simultaneously, cholecalciferol treatment in all tested doses similarly increased grooming, did not change locomotor activity and rearing of the middle-aged and old OVX rats after long-term ovariectomy. Thus, the present results suggest that anxiolytic-like effects of vitamin D₃ are specific in the OVX rats of different age given with solvent or 17β -E₂, since data of the OFT were not able to demonstrate any of alterations in motor or rearing activities in these rats. Combined application of vitamin D₃ (5.0 mg/kg/day) with low dose of 17β -E₂ synergistically decreased anxiety-like profile of the middle-aged OVX rats. However, vitamin D₃ at all doses significantly decreased grooming behavior in the middle-aged OVX rats treated with 17β -E₂.

Unexpectedly, the middle-aged OVX rats treated with vitamin D₃ at doses 1.0 and 2.5 mg/kg/day in combination with 17β -E₂ have practically identical parameters of anxiety-like state as the OVX rats received only 17β -E₂. Thus, this study failed to show any anxiolytic-like effects of vitamin D₃ at doses of 1.0 or 2.5 mg/kg in the middle-aged OVX treated with 17β -E₂. The present results might speculate that there exists some concurrent relation between 17β -E₂ and vitamin D₃ at doses of 2.5 and 5.0 mg/kg/day for the brain structures implicated in the mechanisms of anxiety. In fact, additional application of 17β -E₂ to the middle-aged OVX rats interfered with anxiolytic-like action of vitamin D₃ when it administered at doses 1.0 or

2.5 mg/kg. Thus, the effects of vitamin D₃ administration in a combination with low dose of 17β-E₂ on anxiety-like state of the middle-aged OVX rats are determined by dose of treatment.

ELISA assays demonstrated that administration of cholecalciferol only at doses of 2.5 and 5.0 mg/kg resulted in elevated 25-OH-VD₃ levels in the blood serum of the middle-aged OVX rats with long-term absence of estrogen. Moreover, application of cholecalciferol at these doses with low dose of 17β-E₂ induced more profound increase of 25-OH-VD₃ levels in the blood serum of the middle-aged OVX rats. These data suggest that the different effects of cholecalciferol application in the middle-aged OVX rats with long-term absence of estrogen on anxiety-like behavior in the EPM and LDT did not associated with its effects on 25-OH-VD₃ levels.

Biochemical analysis showed that administration of cholecalciferol at all doses alone or in a combination with 17β-E₂ resulted in elevated 25-OH-VD₃ levels in the blood serum of the old OVX rats with long-term absence of estrogen. Cholecalciferol administered alone at all doses similarly increased estradiol levels in the blood serum of the old OVX rats after long-term ovariectomy. On the other hand, only application of cholecalciferol at a dose of 1.0 mg/kg with low dose of 17β-E₂ induced more profound increase of estradiol levels in the blood serum of the old OVX rats. Moreover, cholecalciferol in several doses failed to induce any changes in calcium concentrations in the blood serum of the old OVX rats given with solvent or 17β-E₂.

These data suggest that the different effects of cholecalciferol application per se in the middle-aged and old OVX rats with long-term absence of estrogen on anxiety-like behavior in the EPM and LDT did not associated with its effects on estradiol, 25-OH-VD₃ and calcium levels.

However, we can speculate that behavioral effects of cholecalciferol treatment with low dose of 17β-E₂ in the EPM and LDT tests might connected with fluctuations of estradiol levels in the blood serum of the middle-aged and old OVX rats. It is possible that specific sites of action involved in the anxiolytic-like effects of cholecalciferol that also modulated by estrogens are affected by the endocrine milieu that prevails at different period for the middle-aged and old female rats. Moreover, after a long-time absence of ovarian fluctuations an adaptive process may contribute to a better response for cholecalciferol administration at a dose of 1.0 mg/kg in the old female rats and for cholecalciferol at a dose of 5.0 mg/kg in the middle-aged female rats.

The role of ovarian hormones in anxiety and stress sensitivity is of great interest for women transitioning through menopause [37, 38]. Mood disorders during menopause could be partly due to loss of estrogen with menopause because estrogen is known to have neuroprotective effects on brain [39, 40]. Menopausal hormonal therapy (MHT) may improve the symptoms of affective-related state in people or decrease the risk of developing mood disturbances in older women, but this is unclear because in some studies MHT does not stop the development of anxiety-like symptoms in elderly postmenopausal women [41]. The exact role of estrogen still needs to be defined.

Menopause are also at higher risk of developing VD deficiency due to decreased dietary intake, less sun exposure, restricted outdoor activity and a decreased capacity to produce enough calcitriol as a result of an age related decline in hydroxylation by kidneys [42–44].

Vitamin D₃ is a hormone precursor which is transformed into 1,25-dihydroxyvitamin D (1,25-(OH)₂D₃) in the liver and kidney [45]. Through decades, this active form of VD has been involved in the brain development and functions of the central nervous system (CNS) [46, 47]. Hormonal form 1,25(OH)₂D₃ enters the brain via the blood-brain barrier to act directly on cells containing its nuclear receptor, the VDR [48, 49].

The presence of VD receptors (VDR) outside the skeletal system, enterocytes and renal tubular cells was confirmed in many cell types including immune cells, neurons, pancreatic cells, myocytes, cardiomyocytes, endothelium cells, which stress pleiotropic activity of VD [50, 51]. The active form of vitamin D is transferred to astrocytes where it can bind to VDR and initiate gene transcription or be inactivated when in excess [48, 49]. Alternatively, $1,25(\text{OH})_2\text{D}_3$ can induce autocrine or paracrine rapid non-genomic actions since all brain cell types express the other membrane receptor of VD [48]. It is possible that the behavioral effects of cholecalciferol are mediated by multiple target regions, including brain centers that are involved in the mechanisms of anxiety-like behavior. Regardless, it cannot presently exclude possible indirect effects of vitamin D_3 on different neurotransmitter circuits. Further research is required to investigate the underlying mechanisms in its anxiolytic-like effects.

It is well-established both systemic effects of VD on calcium metabolism and neuroprotective effects of VD on the brain [52]. Low VD levels has been implicated in the pathophysiological mechanisms of cardiovascular diseases, depression, anxiety, cognitive disorders, obesity, metabolic syndrome, type 2 diabetes, various types of cancer and immune disorders [10, 52]. According to Gaugris and co-workers [53], the prevalence of low VD levels appears to be high in postmenopausal women. Additionally, the decline of estrogens after menopause decreases the activity of $1\alpha\text{-OHase}$, which results in lower synthesis of the active VD form [53, 54]. Application of vitamin D_3 in specific periods of women's life, seems to be of great importance, because it may reduce the risk of affective-related diseases during menopausal period [55–57]. However, the current data for vitamin D_3 application studies are very incomplete and need of more intensive investigations. The main point of question is to examine how the interaction between vitamin D_3 and estradiol might alter at specific periods of women's life, and the impacts of such alterations elsewhere in the postmenopausal woman. VDR have been identified throughout the female reproductive tract [58, 59]. Some studies have demonstrated a direct modulation by VD of estradiol, estrone and progesterone production in human ovarian cells [60–62]. It could be supposed that estrogens and VD share similar targets of the brain to induce their anxiolytic-like effects. However, the behavioral manifestations of VD at various doses are completely different in the middle-aged and old OVX females. It is likely that VD acts through a various mechanisms that are sensitive in female rats of different age with long-term absence of ovarian hormones. Moreover, it is completely needed to understand the precise mechanisms of how VD treatment alone or in a combination with $17\beta\text{-estradiol}$ supplementation may affect women's anxiety-related state.

These points illustrate how the current state of VD treatment research is incomplete and in need of more intensive research. Working toward uncovering how the interaction between VD and estradiol changes after menopause, and the implications of these changes elsewhere in the postmenopausal woman, is necessary for providing the most complete understanding of how VD treatment alone or in a combination with $17\beta\text{-estradiol}$ supplementation may affect women's affective-related state.

VD as changes in VDR impact on various brain neurotransmitters, and thus suggest a potential role of vitamin D in causing and redressing mood disorders [63]. It could be supposed, even though estrogens and cholecalciferol share similar targets on monoaminergic or another neurotransmitter system to induce their anxiolytic-like effects, the behavioral manifestations of cholecalciferol are completely different in the old OVX females. It is likely that cholecalciferol acts through a different mechanisms of action that is sensitive to the age of female rats with long-term absence of ovarian hormones.

In conclusion, the results of this study can be summarized as follows: all tested doses of cholecalciferol given alone are produced anxiolytic-like effects in the

EPM and LDT in the old OVX female rats; the one specific dose of cholecalciferol (1.0 mg/kg/day) in the old OVX rats and another one dose of cholecalciferol (5.0 mg/kg/day) in the middle-aged OVX rats are able to induce synergic anxiolytic-like effect in the EPM and LDT; effects of cholecalciferol on anxiety-like behavior in the middle-aged and old OVX rats after long-term absence of estrogen are dependent from absence or presence of additional hormonal treatment as 17β -E₂. Further investigations is to be addressed in relation to such issues: whether different effects of cholecalciferol on anxiety-like behavior are dependent from different age of rats, or whether different doses of cholecalciferol on anxiety-like behavior in OVX rats with different age rats might lead to negative versus positive effects. Further research, with properly designed experimental studies, is needed to test this hypothesis. In addition, further research is needed to elucidate the biochemical mechanism/(s) of cholecalciferol effects on the anxiety-like behavior and their physiological relevance for development of mood impairment at estrogen deficiency at aging. Furthermore, the mechanism by which cholecalciferol produces anxiolytic-like effect in the middle-aged and old OVX rats and the implications of this in brain function need to be investigated in future research. Moreover, further studies are needed to evaluate the association of VD with estrogen-related pathways and to conduct detail experiments together with biochemical studies of these subjects to verify the significance of this study.

5. Conclusions

The present data of the preclinical study indicates that chronic cholecalciferol at a dose of 5.0 or 1.0 mg/kg, s.c. treatment decreased anxiety-related behavior after impairment induced by long-term ovariectomy in the middle-aged and old female rats with long-term absence of estrogen, respectively. The data also indicate that the combination of cholecalciferol at a dose of 5.0 mg/kg and 17β -E₂ is more effective than 17β -E₂ alone in the middle-aged OVX rats inducing more synergic anxiolytic-like effects in the EPM and LDT. Moreover, a combination of cholecalciferol at a dose of 1.0 mg/kg s.c. and 17β -E₂ is more effective than 17β -E₂ alone in the old OVX rats inducing a more synergic anxiolytic-like effects in the EPM and LDT. Furthermore, this is the first study to show a beneficial effect of chronic cholecalciferol at dose of 1.0 mg/kg s.c. administration on anxiety-related states induced by long-term ovariectomy in the old female rats. Importantly, these results suggest that 17β -E₂ administration interfered with anxiolytic-like action of cholecalciferol administered alone at doses of 2.5 or 5.0 mg/kg to the old OVX rats with long-term absence of estrogens. This work promotes more effective creating of the novel therapeutic targets and strategies for anxiety-like treatment in the middle-aged and old subjects with long-term estrogen deficiency.

Acknowledgements

The reported study was funded by Russian Science Foundation (RSF) accordingly to the research project No. 16-15-10053.

Conflict of interest

The author declares no conflict of interest.

Author contributions

Author Julia Fedotova designed the study, performed the experiments, analyzed the data and wrote the protocol and manuscript.

Author details

Julia Fedotova^{1,2}

1 Laboratory of Neuroendocrinology, I.P. Pavlov Institute of Physiology of the Russian Academy of Sciences, St. Petersburg, Russia

2 Faculty of Food Biotechnologies and Engineering, ITMO University, St. Petersburg, Russia

*Address all correspondence to: julia.fedotova@mail.ru

IntechOpen

© 2018 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Terauchi M, Hiramitsu S, Akiyoshi M, Owa Y, Kato K, Obayashi S, et al. Associations between anxiety, depression and insomnia in peri- and postmenopausal women. *Maturitas*. 2012;**72**:61-65. DOI: 10.1016/j.maturitas.2012.01.014
- [2] Castanho TC, Moreira PS, Portugal-Nunes C, Novais A, Costa PS, Palha JA, et al. The role of sex and sex-related hormones in cognition, mood and well-being in older men and women. *Biological Psychology*. 2014;**103**:158-166. DOI: 10.1016/j.biopsycho.2014.08.015
- [3] Soares CN, Maki PM. Menopausal transition, mood, and cognition: An integrated view to close the gaps. *Menopause*. 2010;**17**:812-814. DOI: 10.1097/gme.0b013e3181de0943
- [4] Maki PM, Freeman EW, Greendale GA, Henderson VW, Newhouse PA, Schmidt PJ, et al. Summary of the National Institute on aging-sponsored conference on depressive symptoms and cognitive complaints in the menopausal transition. *Menopause*. 2010;**17**:815-822. DOI: 10.1097/gme.0b013e3181d763d2
- [5] Arevalo M-A, Azcoitia I, Garcia-Segura LM. The neuroprotective actions of oestradiol and oestrogen receptors. *Nature Reviews Neuroscience*. 2015;**16**:17-29. DOI: 10.1038/nrn3856
- [6] Soares CN. Mood disorders in midlife women: Understanding the critical window and its clinical implications. *Menopause*. 2014;**21**:198-206. DOI: 10.1097/GME.0000000000000193
- [7] Walf AA, Frye CA. ER(beta)-selective estrogen receptor modulators produce antianxiety behavior when administered systemically to ovariectomized rats. *Neuropsychopharmacology*. 2005;**30**:1598-1609
- [8] Soares CN. Depression in peri- and postmenopausal women: Prevalence, pathophysiology and pharmacological management. *Drugs & Aging*. 2013;**30**:677-685. DOI: 10.1007/s40266-013-0100-1
- [9] Soares CN, Almeida OP, Joffe H, Cohen LS. Efficacy of estradiol for the treatment of depressive disorders in perimenopausal women: A double-blind, randomized, placebo-controlled trial. *Archives of General Psychiatry*. 2001;**58**:529-534
- [10] Pae CU, Mandelli L, Kim TS, Han C, Masand PS, Marks DM, et al. Effectiveness of antidepressant treatments in pre-menopausal versus post-menopausal women: A pilot study on differential effects of sex hormones on antidepressant effects. *Biomedicine & Pharmacotherapy*. 2009;**63**:228-235. DOI: 10.1016/j.biopha.2008.03.010
- [11] Vedder LC, Bredemann TM, McMahon LL. Estradiol replacement extends the window of opportunity for hippocampal function. *Neurobiology of Aging*. 2014;**35**:2183-2192. DOI: 10.1016/j.neurobiolaging.2014.04.004
- [12] Genaro PS, Pereira GAP, Pinheiro MM, Szejnfeld VL, Martini LA. Relationship between nutrient intake and vitamin D status in osteoporotic women. *International Journal for Vitamin and Nutrition Research*. 2007;**77**:376-381. DOI: 10.1024/0300-9831.77.6.376
- [13] Scheid V, Ward T, Ch WS, Watanabe K, Liao X. The treatment of menopausal symptoms by traditional east Asian medicines: Review and perspectives. *Maturitas*. 2010;**66**:111-130. DOI: 10.1016/j.maturitas.2009.11.020
- [14] Studd J, Nappi RE. Reproductive depression. *Gynecological*

Endocrinology. 2012;**28**(Suppl. S1):42-45. DOI: 10.3109/09513590.2012.651932

[15] Peng W, Sibbritt DW, Hickman L, Adams J. Association between use of self-prescribed complementary and alternative medicine and menopause-related symptoms: A cross-sectional study. *Complementary Therapies in Medicine*. 2016;**23**:666-673

[16] Holick MF, Frommer JE, McNeill SC, Richtand NM, Henley JW, Potts JT Jr. Photometabolism of 7-dehydrocholesterol to previtamin D3 in skin. *Biochemical and Biophysical Research Communications*. 1977;**76**:107-114

[17] Groves NJ, McGrath JJ, Burne TH. Vitamin D as a neurosteroid affecting the developing and adult brain. *Annual Review of Nutrition*. 2014;**34**:117-141. DOI: 10.1146/annurev-nutr-071813-105557

[18] Christakos S, Raval-Pandya M, Wernyj RP, Yang W. Genomic mechanisms involved in the pleiotropic actions of 1,25-dihydroxyvitamin D3. *The Biochemical Journal*. 1996;**316**:361-371

[19] Mizwicki MT, Norman AW. The vitamin D sterol-vitamin D receptor ensemble model offers unique insights into both genomic and rapid-response signaling. *Science Signaling*. 2009;**2**:re4. DOI: 10.1126/scisignal.275re4

[20] Cui X, McGrath JJ, Burne TH, Mackay-Sim A, Eyles DW. Maternal vitamin depletion alters neurogenesis in the developing rat brain. *International Journal of Developmental Neuroscience*. 2007;**25**:227-232

[21] De Luca HF. History of the discovery of vitamin D and its active metabolites. *Bone Key*

Reports. 2014;**3**:479. DOI: 10.1016/j.ijdevneu.2007.03.006

[22] Eyles DW, Smith S, Kinobe R, Hewison M, McGrath JJ. Distribution of the vitamin D receptor and 1 α -hydroxylase in human brain. *Journal of Chemical Neuroanatomy*. 2005;**29**:21-30

[23] De Chaves G, Moretti M, Castro AA, Dagostin W, da Silva GG, Boeck CR, et al. Minireview: Vitamin D: Is there a role in extraskeletal health? *Endocrinology*. 2011;**152**:2930-2936

[24] Estrada-Camarena E, Lopez-Rubalcava C, Azucena Hernandez-Aragon A, Silvia Mejia-Mauries S, Picazo O. Long-term ovariectomy modulates the antidepressant-like action of estrogens, but not of antidepressants. *Journal of Psychopharmacology*. 2017;**25**:1365-1377

[25] Bose R, Di Paolo T. Dopamine and GABA receptor imbalance after ovariectomy in rats: Model of menopause. *Journal of Psychiatry & Neuroscience*. 1995;**20**:364-371

[26] Estrada-Camarena E, Fernandez-Guasti A, Lopez-Rubalcava C. Interaction between estrogens and antidepressants in the forced swimming test in rats. *Psychopharmacology*. 2004;**173**:139-145. DOI: 10.1007/s00213-003-1707-4

[27] Idrus NM, Happer JP, Thomas JD. Cholecalciferol attenuates perseverative behavior associated with developmental alcohol exposure in rats in a dose-dependent manner. *The Journal of Steroid Biochemistry and Molecular Biology*. 2013;**136**:146-149

[28] Estrada-Camarena E, Fernandez-Guasti A, Lopez-Rubalcava C. Antidepressant-like effect of different estrogenic compounds in the forced swimming test.

Neuropsychopharmacology.
2003;28:830-838. DOI: 10.1038/
sj.npp.1300097

[29] Pellow S, File SE. Anxiolytic and anxiogenic drug effects on exploratory activity in an elevated plus-maze: A novel test of anxiety in the rat. *Pharmacology, Biochemistry, and Behavior*. 1986;24(3):525-529. DOI: 10.1016/0091-3057(86)90552-6

[30] Pellow S, Chopin P, File SE, Briley M. Validation of open: Closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. *Journal of Neuroscience Methods*. 1985;14:149-167

[31] Menzaghi F, Howard RL, Heinrichs SC, Vale W, Rivier J, Koob GF. Characterization of a novel and potent corticotropin-releasing factor antagonist in rats. *The Journal of Pharmacology and Experimental Therapeutics*. 1994;269:564-572

[32] Edinger KL, Frye CA. Sexual experience of male rats influences anxiety-like behavior and androgen levels. *Physiology & Behavior*. 2007;92:443-453

[33] Pan H-Z, Chen H-H. Hyperalgesia, low-anxiety, and impairment of avoidance learning in neonatal caffeine-treated rats. *Psychopharmacology*. 2006;191:119-125

[34] Fedotova J, Soultanov V, Nikitina T, Roschin V, Ordayn N. Ropren[®] is a polyphenol preparation from coniferous plants that ameliorates cognitive deficiency in a rat model of beta-amyloid peptide-(25-35)-induced amnesia. *Phytomedicine*. 2012;19:451-456. DOI: 10.1016/j.phymed.2011.09.073

[35] Nelly M, Rivera V, Tenorio AG, Fernández-Guasti A, Estrada-Camarena E. The post-ovariectomy interval

affects the antidepressant-like action of citalopram combined with ethynyl-estradiol in the forced swim test in middle aged rats. *Pharmaceuticals*. 2016;9:21

[36] Okada M, Hayashi N, Kometani M, Nakao K, Inukai T. Influences of ovariectomy and continuous replacement of 17 β -estradiol on the tail skin temperature and behavior in the forced swimming test in rats. *Japanese Journal of Pharmacology*. 1997:93-96

[37] Burger H. The menopausal transition-endocrinology. *The Journal of Sexual Medicine*. 2008;5:2266-2273. DOI: 10.1111/j.1743-6109.2008.00921.x

[38] MacLennan AH, Taylor AW, Wilson DH. Hormone therapy use after the women's health initiative. *Climacteric*. 2004;7:138-142

[39] Przybelski R, Binkley N. Is vitamin D important for preserving cognition? A positive correlation of serum 25-hydroxyvitamin D concentration with cognitive function. *Archives of Biochemistry and Biophysics*. 2007;460:202-220

[40] Wilkins C, Sheline Y, Roe C, Birge S, Morris J. Vitamin D deficiency is associated with low mood and worse cognitive performance in older adults. *The American Journal of Geriatric Psychiatry*. 2006;14:1032-1040

[41] Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: Principal results from the women's health initiative randomized controlled trial. *Journal of the American Medical Association*. 2002;288:321-333

[42] Cheema C, Grant BF, Marcus R. Effects of estrogen on circulating "free" and total 1,25-dihydroxyvitamin

D and on the parathyroid-vitamin D axis in postmenopausal women. *The Journal of Clinical Investigation*. 1989;**83**:537-542. DOI: 10.1172/JCI113915

[43] Schnatz PF, Marakovits KA, O'Sullivan DM, Ethun K, Clarkson TB, Appt SE. Response to an adequate dietary intake of vitamin D3 modulates the effect of estrogen therapy on bone density. *Journal of Women's Health* (2002). 2012;**21**:858-864. DOI: 10.1089/jwh.2011.3244

[44] Robbins JA, Aragaki A, Crandall CJ, Manson JE, Carbone L, Jackson R, et al. Women's health initiative clinical trials: Interaction of calcium and vitamin D with hormone therapy. *Menopause*. 2014;**21**:116-123. DOI: 10.1097/GME.0b013e3182963901

[45] Stewart A, Wong K, Cachat J, Elegante M, Gilder T, Mohnot S, et al. Neurosteroid vitamin D system as a nontraditional drug target in neuropsychopharmacology. *Behavioural Pharmacology*. 2010;**21**:420-426. DOI: 10.1097/FBP.0b013e32833c850f

[46] Holick MF. High prevalence of vitamin D inadequacy and implications for health. *Mayo Clinic Proceedings*. 2006;**81**:353-373

[47] Penckofer S, Kouba J, Byrn M, Estwing Ferrans C. Vitamin D and depression: Where is all the sunshine? *Issues in Mental Health Nursing*. 2010;**31**:385-393. DOI: 10.3109/01612840903437657

[48] Eyles DW, Burne TH, McGrath JJ. Vitamin D, effects on brain development, adult brain function and the links between low levels of vitamin D and neuropsychiatric disease. *Frontiers in Neuroendocrinology*. 2013;**34**:47-64. DOI: 10.1016/j.yfrne.2012.07.001

[49] Eyles DW, Liu PY, Josh P, Cui X. Intracellular distribution of the vitamin D receptor in the brain:

Comparison with classic target tissues and redistribution with development. *Neuroscience*. 2014;**268**:1-9. DOI: 10.1016/j.neuroscience.2014.02.042

[50] Holick MF. Vitamin D deficiency. *New England Journal of Medicine*. 2007;**357**:266-281

[51] Drevets WC, Price JL, Furey ML. Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Structure and Function*. 2008;**213**:93-118. DOI: 10.1007/s00429-008-0189x

[52] Fernandes de Abreu DA, Eyles D, Féron F. Vitamin D, a neuro-immunomodulator: Implications for neurodegenerative and autoimmune diseases. *Psychoneuroendocrinology*. 2009;**34**:S265-S277. DOI: 10.1016/j.psyneuen.2009.05.023

[53] Gaugris S, Heaney RP, Boonen S, Kurth H, Bentkover JD, Sen SS. Vitamin D inadequacy among post-menopausal women: A systemic review. *QJM: An International Journal of Medicine*. 2005;**98**:667-676

[54] Bikle DD. Vitamin D metabolism, mechanism of action and clinical applications. *Chemistry and Biology*. 2014;**21**:319-329. DOI: 10.1016/j.chembiol.2013.12.016

[55] Garcion E, Wion-Barbot N, Montero-Menei C, Berget F, Wion D. New clues about vitamin D functions in the nervous system. *Trends in Endocrinology and Metabolism*. 2005;**13**:100-105

[56] Kalueff AV, Eremin K, Tuohimaa P. Mechanisms of neuroprotective action of vitamin D3. *The Biochemist*. 2004;**69**:738-741

[57] Eyles D, Brown J, Mackay-Sim A, McGrath J, Feron F. Vitamin D3 and brain development. *Neuroscience*. 2003;**118**:641-653

[58] Halloran BP, De Luca HF. Effect of vitamin D deficiency on fertility and reproductive capacity in the female rat. *The Journal of Nutrition*. 1980;**110**:1573-1580

[59] Kwiecinski GG, Petrie GI, De Luca HF. 1,25-Dihydroxyvitamin D₃ restores fertility of vitamin D-deficient female rats. *The American Journal of Physiology*. 1989;**256**:E483-E487. DOI: 10.1152/ajpendo.1989.256.4.E483

[60] Kinuta K, Tanaka H, Moriwake T, Aya K, Sato S, Seino Y. Vitamin D is an important factor in estrogen biosynthesis of both female and male gonads. *Endocrinology*. 2004;**141**:1317-1324. DOI: 10.1210/endo.141.4.7403

[61] Luk J, Torrealday S, Neal Perry G, Pal L. Relevance of vitamin D in reproduction. *Human Reproduction* (Oxford, England). 2012;**27**:3015-3027. DOI: 10.1093/humrep/des248

[62] Ozkan S, Jindal S, Greenseid K, Shu J, Zeitlian G, Hickmon C, et al. Replete vitamin D stores predict reproductive success following in vitro fertilization. *Fertility and Sterility*. 2010;**94**:1314-1319. DOI: 10.1016/j.fertnstert.2009.05.019

[63] Kiraly SJ, Kiraly MA, Hawe RD, Makhani V. Vitamin D as neuroactive substance: Review. *The Scientific World Journal*. 2006;**6**:125-139. DOI: 10.1100/tsw.2006.25