Personality patterns of people with medically refractory epilepsy – does the epileptogenic zone matter?

F. Novais¹, G. Andrade¹, J. Crawford¹, L. Câmara Pestana¹, M. Andrea¹, S. Loureiro¹, M.L. Figueira¹

¹Centro de Referência de Epilepsia Refratória, Hospital Santa Maria – Centro Hospitalar Universitário Lisboa Norte

Epilepsy painting, Adrian Sibley. Available at saatchiart.

Introduction: The high prevalence of dysfunctional personality traits among people with epilepsy was noted many decades ago and it has been discussed over the years. Kraepelin (1923) described certain distinctive personality characteristics in these patients, such as meticulousness, slowness, circumstantiality, lability, irritability, explosiveness and a particular proneness to religiosity (1). Between 1973 and 1986, Norman Geschwind wrote substantially about what was later called the “Geschwind syndrome”, in people with temporal lobe epilepsy (2).

Recent research has shown that people with epilepsy have certain personality characteristics or traits, such as lower self-consideration and self-esteem (3). Some personality traits may be associated with epilepsy-related factors, namely, earlier age of onset, longer duration of epileptic history and higher seizure frequency (4).

Aims:

- to determine the rate of dysfunctional personality patterns before and after epilepsy surgery in people with refractory epilepsy.
- to determine their types before surgery;
- to determine if the epileptogenic zone is a predictor of any dysfunctional personality pattern.

Methods: We conducted an ambispective observational study, including refractory epilepsy surgery candidates.

Demographic, psychiatric and neurological data were recorded. Evaluation of personality was made using the Millon Clinical Multiaxial Inventory-II (MCMI-II).

Pre-surgical predictors of personality patterns were determined using a linear regression model.

The proportion of patients with dysfunctional personality patterns, before and after surgery, was compared using the McNemar’s test. Then a generalized estimating equation model was performed to include predictors of changes in this rate.

Results: 192 participants were included.

70% had a dysfunctional personality pattern before surgery. After surgery, this percentage dropped to 58%. The difference was statistically significant after adjusting for potential confounders ($p=0.013$).

Figure 1 – Dysfunctional personality rates before and one-year after surgery.

Socio-demographical and clinical characteristics of the participants.

Age, years	38.8 ± 11.6
Males, n (%)	85 (42.7)
Education, years	10.1 ± 4.5
Active workers, n (%)	93 (64.6)
Unemployed, n (%)	39 (21.7)
Retired, n (%)	25 (13.9)
Married, n (%)	74 (50.0)
Age at onset, years	15.9 ± 11.3
Duration of epilepsy, years	22.7 ± 13.0
Temporal epileptogenic zone, n (%)	168 (86.6)
• Mesial, n (%)	106 (65)
• Neocortical, n (%)	57 (35)
Extratemporal epileptogenic zone, n (%)	26 (13.4)
• Frontal epileptogenic zone, n (%)	13 (6.7)
• Multilobar epileptogenic zone, n (%)	10 (5.2)
• Occipital epileptogenic zone, n (%)	2 (1.0)
• Parietal epileptogenic zone, n (%)	1 (0.5)
Laterality of the epileptogenic focus	
• Left	97 (50.8)
• Right	84 (44.0)
• Bilateral	10 (5.2)
Number of antiepileptic drugs	2.3 (0.6)

Table 1 – Socio-demographical and clinical characteristics of the participants.

The most common types were Cluster C personality patterns. Temporal epileptogenic zone was a significant predictor of higher scores of the Avoidant (Coef. 11.8; CI -0.59 23.7; $p=0.051$) and Compulsive (Coef. 9.55; CI 2.48 16.6; $p=0.008$) personality patterns and lower scores of Histrionic (Coef. -11.4; CI -21.2 -1.55; $p=0.024$) and Antisocial (Coef. -8.4; CI -15.6 -1.25; $p=0.022$) personality patterns, compared to extratemporal epileptogenic zone.

Conclusion: People with refractory epilepsy have high rates of dysfunctional personality patterns. These patterns differ according to the epileptogenic zone.

REFERENCES:

1. Mendez MF, Doss RC, Taylor JL, Argiello R. Relationship of seizure variables to personality disorders in epilepsy. J Neuropsychiatry Clin Neurosci. 1993 Aug 1;5(3):283–6.
2. Devinsky J, Schachter S. Norman Geschwind’s contribution to the understanding of behavioral changes in temporal lobe epilepsy: The February 1974 lecture. Epilepsy Behav. 2009 Aug 1;15(4):417–24.
3. Balottin L, Mannarini S, Candeloro D, Mita A, Chiappedi M, Balottin U. Rorschach Evaluation of Personality and Emotional Characteristics in Adolescents With Migraine Versus Epilepsy and Controls. Front Neurol. 2018;9:160.
4. Wang X, Lv Y, Zhang W, Meng H. Cognitive Impairment and Personality Traits in Epilepsy: Characterization and Risk Factor Analysis. J Nerv Ment Dis. 2018 Oct;206(10):794–9.