Estimated blood loss: A new hemoglobin mass-based model

S. Jaramillo-Selman¹, A. Blasi-Ibañez¹, M. Montane-Muntane¹, D. Capitan¹, F. Aguilar², R. Navarro-Ripoll¹

¹Department of Anesthesiology, Hospital Clinic de Barcelona, Universitat de Barcelona - Barcelona (Spain), ²European Foundation for the Study of Chronic Liver Failure - Barcelona (Spain)

BACKGROUND AND GOAL OF STUDY

- Estimation of blood loss (EBL) has relevant clinical and research applications.
- Volume-based EBL formulas have been found to be significantly inaccurate.
- Main goal: Testing a new hemoglobin (Hb) mass-based model to estimate surgical blood loss and to predict hemoglobin drop 48h after surgery.
- Secondary goal: Comparison between the new proposed hemoglobin mass-based model and main volume-based formulas.

MATERIALS AND METHODS

- 100 consecutive patients undergoing laparoscopic urologic surgery were studied.
- Hb was measured from the suction canister.
- Blood samples were drawn from the patient before surgery and 24, 48 and 72h after surgery.
- A model was designed to estimate blood loss in grams of hemoglobin mass and to predict hemoglobin drop 48h after surgery.
- Concordance between estimates and directly measured values was assessed using the Intraclass Correlation Coefficient (ICC) and Bland-Altman analysis.
- Main volume-based formulas were compared with the proposed model using the ICC and Bland-Altman analysis.

DISCUSSION

- Excellent consistency of ICC was found between estimated and measured hemoglobin mass loss (0.980, $p < 0.001$), and between predicted and measured hemoglobin level drop (0.974, $p < 0.001$).
- Narrow limits of agreement for estimated blood loss (± 21.7 g) and predicted hemoglobin drop (± 0.54 g/dl) were observed.
- The proposed model was more reliable than volume-based formulas.

CONCLUSIONS

- This is the first model aimed at estimating blood loss based on the quantification of lost hemoglobin mass, which also accurately predicts hemoglobin drop 48h after surgery.
- Further studies are needed to validate the new model in other scenarios.

RESULTS

Fig A: Bland Altman plot, comparison of estimated Hb mass loss with measured Hb mass loss.
Fig B: Bland Altman plot, comparison of predicted perioperative Hb drop with observed perioperative Hb drop.

EBL method	ICC Consistency	ICC Absolute Concordance	Bias + limits of agreement (α 0,05)
New hemoglobin mass-based model	0,980 ($p < 0,001$)	0,907 ($p < 0,001$)	22,4 $\pm$ 21,7 g
Bourke et al. (1974)	0,665 ($p < 0,001$)	0,489 ($p < 0,001$)	370 $\pm$ 1230 ml
Ward et al. (1980)	0,585 ($p < 0,001$)	0,440 ($p < 0,001$)	361 $\pm$ 1247 ml
Gross et al. (1983)	0,577 ($p < 0,001$)	0,465 ($p < 0,001$)	355 $\pm$ 1238 ml
OSTHEO study (2001)	0,578 ($p < 0,001$)	0,416 ($p < 0,001$)	573 $\pm$ 1227 ml
Meunier et al. (2008)	0,879 ($p < 0,001$)	0,664 ($p < 0,001$)	-413 $\pm$ 893 ml
López-Picado et al. (2017)	0,282 ($p < 0,001$)	0,218 ($p < 0,001$)	-345 $\pm$ 1244 ml

Table: Summary of results of the intraclass correlation coefficient (ICC) and Bland-Altman analysis of the new haemoglobin mass-based model and volume-based formulae studied.

REFERENCES

Goodnough LT, Panigrahi AK: Estimating Blood Loss. AnesthAnalg 2017; 125 : 13-4.