Floods in Madeira Island: A Classic Portuguese Case of Post-Traumatic Stress Disorder in a child

Hugo Canas Simião^{*a}; Rui Ferreira Carvalho^b; Nuno de Moura^a; Leonor Santana^a; Ana Corina^c

No Financial Interest
* hugo.simiao@gmail.com

^a Department of Psychiatry and Mental Health, Centro Hospitalar de Lisboa Ocidental, Rua da Junqueira 126, 1340-019, Lisbon

^b Child and Adolescence Psychiatry Service, Centro Hospitalar de Lisboa Norte, Av. Prof. Egas Moniz, 1649-035, Lisbon

^c Child and Adolescence Psychiatry Service, Hospital Dr. Nélio Mendonça, Av. Luís de Camões 6180, 9000-177 Funchal, Ilha da Madeira

Introduction and Objectives

The most frequently diagnosed mental illness in population affected by floods is post-traumatic stress disorder (PTSD). According to reviews and meta-analyses, the prevalence of PTSD is 30–40% among direct disaster survivors. The benefits of early targeted interventions on acute psychological distress after traumatizing events have been thoroughly studied. Our aim is to discuss a clinical case of PTSD in the aftermath of the floods and mudslides of 20/02/2010 in Madeira Island, Portugal, which caused the death of 42 people and left over a hundred injured, and to review and present the most relevant literature regarding PTSD in childhood and adolescence, transposing its applicability to the clinical practice and reflecting on the transitional challenges between Child and Adolescent Psychiatry and Adult Psychiatry.

Methods

A structured and integrated psychiatric interview was applied on an 18-year-old patient with a retrospective focus on the consequences of exposure to a traumatic event when he was 10 years old, after obtaining his informed consent. A non-systematic review of the literature was performed using a combination of terms: “PTSD”, “childhood/child and adolescence/adolescent”, “trauma” and “psychiatry”.

Clinical Case

- R., 10-years-old, father with anxiety, no other relevant medical history.
- When R. was 10 years old, living with his parents, the family got a call for help from an aunt who lived nearby because of flooding and rock sliding. On the way, they found a road blocked by a boulder and tried to return, but got surrounded by mud and rocks. In great anguish, they were able to leave the car. In the aftermath, they found a group of people, having R.’s mother told R. to go with them while the parents were going to help the aunt. A wall fell and the group of people R. was supposed to join were dragged by the mud and rockslide, not surviving. R. was then submersed to his chest with mud, being saved by a lady which took the family in her home. The next day, they went to the ER because of R.’s complaints of anxiety and complete insomnia. He then initiated regular visits for psychology and child and adolescence psychiatry.

- In the first visit, R.’s drew spontaneously a car at a muddy floor and with sad sun (Picture 1). R. avoided being in a car, would get very anguished with rain, including in videogames, would refuse to get out of the house in bad weather, refused to go through the street where the event had taken place in.
- For about 3 years, he had intrusive flashbacks of the day and had frequent monothemed nightmares related to floodings, mud and rocks. He also mentioned panic attacks, palpitations, increased breath rate, separation anxiety, insomnia, recurring nightmares and depersonalization experiences, when he would relieve the moment “as if I was seeing from the outside” (sic) when peers and teachers would try to know more about the traumatic event. A regular follow-up and a multidisciplinary team were always present.
- After 3 years, R. obtained remission of symptoms. Picture 2 shows a second drawing by R, with a clean floor under the car and a happy sun.

Picture 1 – R.’s first drawing.

Picture 2 – R.’s second drawing.

Discussion and Conclusions

- Several studies have shown that a majority of people will likely experience at least one traumatic event in their lives. However, many will not develop PTSD. Adults with PTSD can have other problems as well, including depression, drug and alcohol abuse, or other anxiety disorders.
- Discussing PTSD, promoting its **screening after potentially traumatic events** and pursuing an **integrated approach** is necessary and, as illustrated by this case, is proven to be beneficial.

REFERENCES

Asim M, Mekkodathil M, Sathian B, Elayedath R, Kumar RN, Simkhada P, van Teijlingen E. (2019) PostTraumatic Stress Disorder among the Flood Affected Population in Indian Subcontinent. Nepal J Epidemiol. 755-758.
James LE, Welton-Mitchell C, Noel JR, James AS (2019). Integrating mental health and disaster preparedness in intervention: a randomized controlled trial with earthquake and flood-affected communities in Haiti. Psychological Medicine 1–11
Gagne, C., Dayan, P., & Bishop, S. J. (2018). When planning to survive goes wrong: predicting the future and replaying the past in anxiety and PTSD. Current Opinion in Behavioral Sciences, 24, 89-95.
Maia, Â., & Fernandes, E. M. (2003). Epidemiologia da perturbação pos stress traumático (PTSD) e avaliação de resposta ao trauma.
Cloitre, M., Stolbach, B. C., Herman, J. L., Kolk, B. V. D., Pynoos, R., Wang, J., & Petkova, E. (2009). A developmental approach to complex PTSD: Childhood and adult cumulative trauma as predictors of symptom complexity. Journal of traumatic stress, 22(5), 399-408.