WHITMAN COLLEGE

Cyanation Methods for Boron Cluster Anions

Marcus A. Juhasz

ACS National Meeting, Spring 2020

A c k n o w l e d g m e n t s

Students

Woody Walters
Grant Domecus

Austin Kamin

Ethan Raffman

Natalie Hodis

Will Mendelsohn

Greg Hof

Julian Reed

Austin Shaff

Walker Bruhn

Clint Vorauer

Greg Dwulet

Rosie Midget

Graham Matheson

Paul Chang

Aaron Rosenbaum

Donna Phan

Nathan Neff-Mallon

Laura Coulson

Kieran Burkholder

Matt Badura

Michael van Heyst

John Heurich

Michelle Starr

Elizabeth Deeran

Whitman College

Colleagues & Collaborators

Doug Juers (X-ray, Whitman)

Funding:

Murdock Charitable Trust

NSF (CHE-1764344)

Whitman College

The CB_{11}^- carborane cluster

CB_{11}^- clusters:

- Delocalized bonding
- Stable skeleton: “ σ -aromatic”
- Weakly-coordinating anions
- Potential applications
  - Superacidity
  - Counteranion: [Catalyst⁺]
  - Ionic liquids
  - Boron source for BNCT
  - Optical materials

The “parent” anion:

Synthesis of the parent cluster: *closo*- $\text{CB}_{11}\text{H}_{12}^-$

Typical Results:

10 g batch

50 – 60% yield

2 weeks

Methods for derivatizing CB_{11} borons (*not many!*)

$CB_{11}H_{12}^-$ is *inorganic* and σ -aromatic, but often reacts like a π -aromatic organic compound.

Most B-substituted derivatives result from two approaches

Electrophilic substitution

Cross-coupling

MWI accelerates CB_{11}^- reactions

Observation: MWI shortens Kumada coupling reactions

Michael van Heyst, Allegheny 2008
Nathan Neff-Mallon, Whitman 2012

Reactions that require
reflux for days complete in
minutes under MWI

Published examples:

Electrophilic methylation

Valášek, M.; Štursa, J.; Pohl, R.; Michl, J. *Inorg. Chem.* **2010**, 49 (22), 10247–10254.

Kumada cross-coupling reactions

Himmelsbach, A.; Reiss, G. J.; Finze, M. *Inorg. Chem.* **2012**, 51 (4), 2679–2688.

CN: Cyana ted CB₁₁- derivative s

5114

J. Am. Chem. Soc. 2000, 122, 5114–5124

Gas-Phase Acidities of Some Neutral Brønsted Superacids: A DFT and ab Initio Study

Ilmar A. Koppel,^{*†} Peeter Burk,[†] Ivar Koppel,[†] Ivo Leito,[†] Takaaki Sonoda,[‡] and Masaaki Mishima[§]

Cont
Jakob
Kas
6-10-
Recei

12972

J. Phys. Chem. A 2009, 113, 12972–12978

Gas-Phase Brønsted Superacidity of Some Derivatives of Monocarpa-*closo*-Borates: a Computational Study

Lauri Lipping,^{*} Ivo Leito, Ivar Koppel, and Ilmar A. Koppel^{*}

University of Tartu, Institute of Chemistry, 2 Jakobi St. 51014 Tartu, Estonia

Received: June 10, 2009; Revised Manuscript Received: September 14, 2009

The structures and gas-phase acidities (GA) of several CB₁₁H₁₂H-based carborane acid derivatives (HA) have been calculated with DFT B3LYP method using 6-311+G**, 6-311++G** basis sets. In order to verify the obtained GA values, several systems were also studied at G3(MP2) level of theory. Inserted substituents (CF₃, F, Cl, Br, I, CN, CH₃, etc.) followed the “belts” of the monocarborane cage starting from the boron antipodal to the carbon. In general, the predicted intrinsic gas-phase acidities of the systems varied according to the substituents in the following order of decreasing strength: CF₃ > F > Cl > Br > I > CN > CH₃.

Methods for derivatizing CB_{11} borons (*not many!*)

$\text{CB}_{11}\text{H}_{12}^-$ is *inorganic* and σ -aromatic, but often reacts like a π -aromatic organic compound.

Most B-substituted derivatives result from two approaches

Electrophilic substitution

Cross-coupling

Synthesis of $12\text{-CN-CB}_{11}\text{H}_{11}^-$

Donna Phan

Electrophilic cyanation: CNBr

MS, calc'd: 168

Methods for derivatizing CB_{11} borons (*not many!*)

$\text{CB}_{11}\text{H}_{12}^-$ is *inorganic* and σ -aromatic, but often reacts like a π -aromatic organic compound.

Most B-substituted derivatives result from two approaches

Electrophilic substitution

Cross-coupling

Alternative: Cross-coupling

Pd-catalyzed cyanation¹

Kumada Coupling

5 min
up to 85% yield

(1) e.g. Sundermeier, M.; Zapf, A.; Beller, M. *Eur. J. Inorg. Chem.* **2003**, 2003, 3513–3526.

Cu-promoted cyanation of CB_{11}^-

Aaron Rosenbaum

Observations: **CuCN required**, Pd accelerates reaction

Entry	CN^- Source	Catalyst (10 mol% Pd)	Time (min)	Temp ($^{\circ}\text{C}$)	Conversion (^{11}B NMR)
1	NaCN or KCN	$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$	30	240	None
2	NaCN or KCN	$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2 + \text{CuI}$ (18 mol%)	30	240	None
3	NaCN or KCN	$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2 + \text{CuCN}$ (18 mol%)	90	240	None
4	CuCN	$\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$	15	180	Complete
5	CuCN	$\text{Pd}(\text{OAc})_2$	30	200	Complete

Cu-promoted cyanation of CB_{11}^-

Aaron Rosenbaum

^{11}B NMR

MS negative ion, m/z (% abund. > 50%): 167(82), 168(100), 169(79)
Calc'd for $[\text{CB}_{11}\text{H}_{11}\text{CN}]^-$: 167(74), 168(100), 169(82)

IR, KBr disc, cm^{-1} : 3066 (C-H), 2572-2531 (B-H), **2205 ($\text{C}\equiv\text{N}$)**
Calc'd (B3LYP/6-311++G(d,p): 3089 (C-H), 2569-2536 (B-H), **2222 ($\text{C}\equiv\text{N}$)**

Rosenbaum, A. J.; Juers, D. H.; Juhasz, M. A. *Inorg. Chem.* **2013**, 52, 10717-10719.

X-ray structure : $[\text{Et}_4\text{N}^+][12\text{-CN-CB}_{11}\text{H}_{11}^-]$

Aaron Rosenbaum

- Monoclinic, $P2_1$
- $\text{C}\equiv\text{N}$ 1.128 Å
- B-CN 1.557 Å
- $\angle\text{B-C-N}$: 178.56°
- Anion nearly C_{5v}

Rosenbaum, A. J.; Juers, D. H.; Juhasz, M. A. *Inorg. Chem.* **2013**, 52, 10717-10719.

Dic y a n a t i o n o f CB_{11}^-

Aaron Rosenbaum

MS negative ion, m/z (% abund. > 50%): 192(68), 193(100), 194(70)

Calc'd for $[\text{CB}_{11}\text{H}_{10}(\text{CN})_2]^-$: 192(74), 193(100), 194(82)

IR, KBr disc, cm^{-1} : 3060 (C-H), 2593-2533 (B-H), **2214 (C $\equiv$ N)**

Calc'd (B3LYP/6-311++G(d,p): 3089 (C-H), 2586-2551(B-H), **2233 (C $\equiv$ N)**, **2234 (C $\equiv$ N)**

Rosenbaum, A. J.; Juers, D. H.; Juhasz, M. A. *Inorg. Chem.* **2013**. 52, 10717-10719.

X-ray structure : $[\text{Et}_4\text{N}^+][7,12\text{-(CN)}_2\text{-CB}_{11}\text{H}_{10}^-]$

Greg Dwulet, A. Rosenbaum

- Monoclinic, $P2_{1/c}$
- $\text{C}\equiv\text{N}$ 1.143 Å, 1.120 Å
- B-CN 1.556 Å, 1.568 Å
- $\angle\text{B-C-N}$: 179.45° 178.60°
- C_s symmetric anion

Juhasz, M. A.; Juers, D. H.; Dwulet, G. E.; Rosenbaum, A. J. *Acta Cryst. E* **2014**, 70, 411–412.

Heavalicyanated clusters: Preliminary results

- **LC-MS:** mix of $(\text{CN})_n\text{I}_{(6-n)}$ products
- >65% of the mixture is $\text{CB}_{11}\text{H}_6(\text{CN})_6^-$

Cyanated $B_{12}H_{12}^{2-}$ derivatives

Calculated electronic stability of $B_{12}(CN)_{12}^{2-}$ suggests promise for use as a battery electrolyte.

Zhao, H.; Zhou, J.; Jena, P. *Angew. Chem. Int. Ed.* **2016**, 55 (11), 3704–3708.

<https://doi.org/10.1002/anie.201600275>.

Communications

Cluster Compounds Very Important Paper

International Edition: DOI: 10.1002/anie.201600275
German Edition: DOI: 10.1002/ange.201600275

Stability of $B_{12}(CN)_{12}^{2-}$: Implications for Lithium and Magnesium Ion Batteries

Hongmin Zhao⁺, Jian Zhou^{+,*} and Puru Jena^{*}

Abstract: Multiply charged negative ions are seldom stable in the gas phase. Electrostatic repulsion leads either to autodetachment of electrons or fragmentation of the parent ion. With

rules. For example, a multidecker $Cr[BC_5(CN)_6]_2^{2-}$ organometallic complex, which simultaneously satisfies the octet rule (of CN moieties), aromatic Hückel's rule (of organic BC

“We hope that these results will stimulate the synthesis of $B_{12}(CN)_{12}^{2-}$ ”

Cyanation of closo - $B_{12}H_{12}^{2-}$

Cyanation by UV-irradiation: [Trofimenko, 1966](#)

Cyanation by UV-irradiation: [Jenne, 2019](#)

MWI Cyanation of closo- $B_{12}H_{12}^{2-}$

Cyanation by MWI cross coupling: [Kamin, 2020](#)

Kamin, A. A.; Juhasz, M. A. *Inorg. Chem.* **2020**, 59 (1), 189–192.

X-ray structure of
 $(CH_3CN)_3Cu[\mu-B_{12}(CN)_{12}]Cu(CH_3CN)_3$

Conclusions

- MWI has made possible the cyanation of CB_{11}^- and B_{12}^{2-} boron cluster anions
- Monocyanated and dicyanated CB_{11}^- derivatives have been prepared:
 $12\text{-CN-CB}_{11}\text{H}_{11}^-$ and $7,12\text{-(CN)}_2\text{-CB}_{11}\text{H}_{10}^-$
- Percyanation of the dodecaborate anion has been achieved: **$\text{B}_{12}(\text{CN})_{12}^{2-}$**

A c k n o w l e d g m e n t s

Students

Woody Walters
Grant Domecus

Austin Kamin

Ethan Raffman
Natalie Hodis
Will Mendelsohn
Greg Hof
Julian Reed
Austin Shaff

Walker Bruhn
Clint Vorauer
Greg Dwulet
Rosie Midget
Graham Matheson
Paul Chang
Aaron Rosenbaum
Donna Phan
Nathan Neff-Mallon

Laura Coulson
Kieran Burkholder
Matt Badura
Michael van Heyst
John Heurich
Michelle Starr
Elizabeth Deeran

Whitman College

Colleagues & Collaborators

Doug Juers (X-ray, Whitman)

Funding:

Murdock Charitable Trust
NSF (CHE-1764344)
Whitman College