Assesing empiric antibiotherapy in complicated community-acquired intra-abdominal infection.

C. Garcia Bernedo, A. Benitez Cano, D. Bande Julian, L. Bosch Duran, M. Rueda Gamboa, L. Castellort Masco .
Hospital del Mar . Barcelona (Spain)

BACKGROUND AND GOAL OF THE STUDY

Appropriate and early empiric antibiotic therapy for complicated intra-abdominal infection is essential for patient prognosis. Community-acquired intra-abdominal infection (CA-IAI) usually involves sensitive pathogens but incidence of multidrug resistant bacteria is increasing. The goal of the study is to assess suitability of empiric antibiotherapy for CA-IAI in our institution.

MATERIAL AND METHODS

A retrospective clinical review was conducted in our institution from Jan 2016 to Sept 2017. We included patients with CA-IAI and positive intraoperative cultures. We review bacteriology and susceptibility to the main agents used for CA-IAI. Common regimes as empiric antibiotherapy for CA-IAI in our institution includes beta-lactam (3rd generation cephalosporin) + metronidazole or beta-lactamase inhibitor combination (amoxicillin-clavulanic acid or piperacillin-tazobactam). We, carefully, consider extended-spectrum beta-lactamase-producing Enterobacteriaceae, ampicillin-resistant Enterococcus spp, Ps. aeruginosa, and Candida spp. as the main pathogens resistant to empiric coverage in our institution.

RESULTS AND DISCUSSION

137 patients with CA-IAI and positive intraoperative culture were identified. Gram negative bacteria (GNB) remain the major pathogens in CA-IAI (80%), gram positive bacteria are involved in 48% CA-IAI cases, anaerobes 18% and fungi 7%. Extended-spectrum beta-lactamase-producing Enterobacteriaceae was involved in 15% of CA-IAI, ampicillin-resistant Enterococcus spp 4%, Ps. aeruginosa 9% and Candida spp 7%.

CONCLUSIONS

Extended-spectrum beta-lactamase Enterobacteriaceae increasing incidence in community-acquired intra-abdominal infections represents a major problem for the future. In severe complicated intra-abdominal infection with non optimum surgical source control carbapenem should be considered for empiric antibiotherapy despite community-acquired origin.

REFERENCES

1. Guirao X. Rev Esp Quimioter 2009.
2. Jones RN. J Clin Microbiol 2001.

* Antibiotic sensitivity

	AMK	A-C	AZT	CTZ	CTX	CFZ	CIPRO	CEFE	CEFU	PTZ	GEN	IMI
E. coli	95	74	81	81	81	72	71	83	78	76	84	100
Klebsiella spp	95	90	95	95	95	90	90	95	95	90	95	100
Citrobacter spp	100		71	71	71		100	100		71	100	100
Enterobacter spp	100		62	69	54		100	92		62	100	100
Ps. aeruginosa	100		8				100	92		92	100	100

Gram positive bacteria

* (AMK, amikacine; AMP, ampiciline; A-C, amoxicillin-clavulanic; AZT, aztreonam; CTZ, ceftazidime; CTX, cefotaxime; CFZ, cefazolin; CIPRO, ciprofloxacin; CEFE, cefepime; CEFU, cefuroxime; GEN, gentamicin; IMI, imipenem; LEV, levofloxacin; PTZ, piperacillin-tazobactam; TET, tetracycline; TEI, teicoplanin.)

* Antibiotic sensitivity

	AMP	CTX	GEN	LEV	LIN	TET	TEI	VAN
E. Faecalis	100		91	91	100	55	100	100
E. Faecium	63		88	81	100	81	100	100
Enterococcus spp	100		100	100	100	100	100	33
Streptococcus spp	89	100	100	100	100		100	100