PATIENTS WITH NEWLY DIAGNOSED DIABETES HAVE WORSE LONG-TERM OUTCOMES THAN KNOWN DIABETICS AFTER ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION

Mahmoud Abdelnabi ¹, Abdallah Almaghraby ², Ola Abdelkarim ², Yehia Saleh ³, Haitham Badran ⁴

¹Cardiology and Angiology Unit, Department of Clinical and Experimental Internal Medicine, Medical Research Institute, University of Alexandria, Egypt.

²Department of Cardiology, University of Alexandria, Egypt. ³Michigan State University, East Lansing, Michigan, United States of America.

⁴Department of Cardiology, Aim Shams University, Cairo, Egypt

INTRODUCTION

A significant portion of patients presenting with ST elevation myocardial infarction (STEMI) are diagnosed incidentally with diabetes at the time of presentation. The effect of undiagnosed diabetes is thought to be comparable to known diabetics when it comes to long term outcomes.

AIM OF THE WORK

In this study we aimed to determine frequency of undiagnosed diabetic patients and their cardiovascular outcomes in comparison to those with known diabetes.

PATIENTS & METHODS

1109 Patients presented with STEMI at 4 tertiary care centers in the period of January 2011 to December 2017. All patients eligible for the study were either diagnosed with diabetes mellitus (DM) previously or had a glycated hemoglobin (HbA1c) value of 6.5 g/dl or more on presentation with the exclusion of non diabetic patients. We compared their in-hospital and 1-year outcomes and mortality (MACCE).

RESULTS

Of 1109 patients who were eligible, 919 (82.87%) were known diabetics, 190 (17.13%) were newly diagnosed diabetics. There was no significant difference in baseline data of both groups. However, newly diagnosed diabetes had a higher HbA1c in comparison to known diabetics (7.3 vs 6.8 g/dl). In-hospital outcomes were similar in both groups regarding death, myocardial infarction and MACCE, however, the known diabetics group had a higher stroke rate (0.04% vs 0%). For the 1 year follow up, the newly diagnosed diabetics group had a significantly higher MACCE rate.

Table (1): Baseline characteristics

	Total (n=1109)	Known DM (n=919)	P value	New DM (n=190)
Age		51 (±6.9)	0.9	50 (±9.1)
Male Sex		550 (59.8%)	0.71	124 (65.3%)
Hypertension		535 (58.2%)	0.75	119 (62.6%)
Smoking		492 (53.5%)	0.69	111 (58.4%)
Dyslipidemia		101 (11%)	0.91	23 (12.1%)
HbA1C		6.8 (±1)	<0.001	7.3 (±0.8)

All data are represented in mean (± SD) or Number (Percentage)
P value is significant if < 0.05

Table (2): In-Hospital and 1 year MACCE

	Total (n=1109)	Known DM (n=919)	P value	New DM (n=190)
In-hospital MACCE	Death	41 (4.5%)	0.81	8 (4.2%)
	MI	12 (1.3%)	0.73	2 (1.1%)
	TVR	4 (0.4%)	0.9	1 (0.5%)
	CVA/TIA	4 (0.4%)	<0.001	0 (0%)
	MACCE	61 (6.6%)	0.42	11 (5.8 %)
1 Year MACCE	Death	63 (6.9%)	0.91	13 (6.8%)
	MI	19 (2.1%)	0.74	7 (3.7%)
	TVR	9 (1%)	0.005	6 (3.2%)
	CVA/TIA	15 (1.6%)	0.02	4 (2.1%)
	MACCE	106 (11.5%)	0.01	30 (15.8%)

All data are represented in mean (± SD) or Number (Percentage)
P value is significant if < 0.05

CVA: Cerebrovascular accident, MI: Myocardial Infarction, TIA: Transient Ischemic Attack, TVR: Target Vessel Revascularization

CONCLUSION

In the setting of STEMI, newly diagnosed diabetics have comparable in-hospital outcomes to patients with known diabetes. However, they have a higher MACCE in the one year follow up.

EAS

with thanks to the EAS for support
in the form of a Young Investigator Fellowship