The effectiveness of muscle relaxation: consideration of the operating surgeon during laparoscopic procedures

E. Nicolayenko¹, D. Kurenkov¹, E. Evdokimov², S. Chizevskaya¹. ¹ Research and Clinical Center of Russian Railways. Department of Anesthesiology and Intensive Care. ²Russian Medical Academy of Postgraduate Education. Moscow, Russian Federation.

Summary. Abstract presents the results of the study, which evaluated the effectiveness of maintaining a deep muscle relaxation in laparoscopic surgery under general anesthesia from the perspective of convenience of surgical procedures, based on a subjective assessment by the surgeon and evaluated the safety of this method.

Background and Goal of Study. The issue of the optimum level of neuromuscular blockade (NMB) in relation to different types of surgery is debatable. Some studies show the benefits of deep NMB, but this approach is not widely used because of high risk of residual relaxation. The aim of this study was to evaluate the efficacy of maintaining a deep NMB during laparoscopic procedures from the perspective of convenience of surgical procedures, and evaluate the safety of this method.

Materials and methods. We examined 38 patients undergoing laparoscopic procedures under general anesthesia with NMB: ASA I-II, mean age 52.3 (26-69 years), male – 14 (37%). Exclusion criteria were: suspected difficult tracheal intubation; disorder affecting neuromuscular transmission (NMT); known or suspected significant renal dysfunction; known or suspected severe hepatic dysfunction; history of malignant hyperthermia; allergy to opioids, neuromuscular blocking drugs or other medications used during general anesthesia; pregnancy and breastfeeding.

Patients were randomized to two groups: in group I (n=18) we maintained a deep NMB at PTC 2 until the end of the operation; in group II (n=20) we maintained an average depth NMB at T2 level. NMB reversion has been performed after surgery with sugammadex: 4 mg/kg in group I; 2 mg/kg in group II. Tracheal extubation has been implemented only after the recovery NMT to the TOF level ≥ 0.9 . After the end of intervention the operating surgeon was invited to give a personal assessment of the adequacy of NMB (unsatisfactory, satisfactory or optimal) on a scale at the start, at the main stage and at the final part of surgery. To analyze the results we used nonparametric two-sample Kolmogorov-Smirnov test. Stat. significance was assumed at $p < 0.005$.

Figure 1. Comparison of good assessment of relaxation quality is provided at the figure: satisfactory and optimal (left) and optimal only (right).

Results and discussion. Recovery of NMT to TOF 0.9 level occurred within 4 min after the administration of sugammadex and tracheal extubation within 11 min after the end of surgery. There were no incidences of residual relaxation. Comparison of the results of surgeon's survey shows that the percentage of positive rating of NMB (satisfactory and optimal) in the group I was higher than in the group II of an average of 11%. When we compared only the optimal ratings of the NMB - percentage in the group I was higher than in the group II of an average of 44%. Decrease in surgeon's satisfaction by quality of a relaxation to the end of operation has been revealed (Fig.1).

Conclusion. Maintaining a deep NMB during laparoscopic surgery provides better conditions for surgical procedures from the perspective of the operating surgeon. The ending of surgery is the most "problem" period from the point of view of the inadequacy of NMB. Maintain a deep NMB to the last stitch, followed by a complete recovery suggests an objective NMT monitoring and reversion.