The role of scavenger receptors stabilin 1 and 2 on atherosclerosis development in zebrafish

R. Verwilligen, A. van Dijke, M. Hoekstra, J. Bussmann, M. van Eck
Research division of BioTherapeutics, LACDR, Leiden University, the Netherlands

LACDR

Background and relevance

Scavenger receptors are a superfamily of membrane bound receptors that are known to bind and internalize different ligands including oxidized LDL (oxLDL). Two scavenger receptors that are known to mediate the uptake of oxLDL are stabilin 1 (*STAB1*) and stabilin 2 (*STAB2*). *STAB1* & *STAB2* are highly expressed on liver endothelial cells. Several studies, including a microarray study of our research group, have already identified *STAB1* and *STAB2* as risk factors for atherosclerosis.

Hypothesis

AIM

We aim to show in an atherosclerosis zebrafish model whether endothelial and macrophage scavenger receptors *STAB1* and *STAB2* are involved in 1) foam cell formation and 2) early atherogenesis.

Conclusion and perspectives

The uptake of oxLDL by endothelial cells is diminished in *STAB2* zebrafish mutants and is absent in *STAB1* and *STAB2* double mutants, suggesting that zebrafish deficient for *STAB1* and *STAB2* are susceptible to foam cell formation as macrophages will compensate for the loss of function of endothelial *STAB1* and *STAB2*.

A follow-up study will be performed to examine whether *STAB1* and *STAB2* double mutants develop more atherosclerosis.

Preliminary results

A) Experimental set-up; labeled oxLDL and fluorescent hyaluronic acid (FluoHA) - a known ligand for *STAB2* - were injected in the blood circulation of 3 day old zebrafish (days post fertilization; dpf) B) Uptake of FluoHA and oxLDL by endothelial cells and macrophages in WT, *STAB2*^{-/-} and *STAB1*^{-/-} *STAB2*^{-/-}. C) Uptake of labeled LDL by macrophages in WT and *STAB1*^{-/-} *STAB2*^{-/-}. D) Follow-up atherosclerosis study. HCD = High cholesterol diet. Confocal Imaging: magnification 10x overview - 40x caudal vein