Effects of local anesthesia of the skin on pain during lumbar puncture and patient satisfaction

C. Massoth¹, S. Weidlich², D. Poepping¹, S. Gurlit², M. Moellmann², M. Wenk¹

¹Department of Anesthesiology, Intensive Care and Pain Medicine, University Hospital of Muenster - Muenster (Germany), ²Department of Anesthesiology and Intensive Care, St. Franziskus Hospital Muenster - Muenster (Germany)

Background

Although spinal anaesthesia has a more favourable risk-profile, patients usually prefer surgery under general anaesthesia. Besides fear of severe but rare neurological complications, typical reasons for rejection are fear of backache or anticipated pain during puncture. Providing local anaesthesia before lumbar puncture is not a standard procedure.

Objective

To evaluate the efficacy in pain reduction and the effect on patient satisfaction when using different local anaesthetic treatments before i.t. puncture

Methods

Results

Variable	Group	Mean	SD	F	P
Age	1	61,28	15,21	2,188	0,119*
	2	67,56	13,68		
	3	68,19	12,02		
BMI	1	28,13	6,67	0,096	0,908*
	2	27,51	4,05		
	3	27,91	4,77		
ASA	1	2,00	0,60	0,635	0,532*
	2	2,11	0,58		
	3	2,19	0,68		

Tab.1: Demographic data: 1=Control, 2=EMLA, 3=Prilocaine
*Levene's test for homogeneity of variance p-value >0.05: population variances are equal

Fig.1: Patient's rating of pain as less (grey) or greater (blue) than expected, Chi-squared test=24,56 (*p<0.001)

Fig.1: Mean VAS Scores ± SD for pain during lumbar puncture, *p<0.01

The **total procedure time** was significantly longer in Prilocaine-Group in comparison to EMLA-Group (mean duration 5.63min ±1.94 vs. 4.0min ± 2.51)

- no group differences for
- the **number of attempts** to successful spinal puncture
 - Patients' **future decision for spinal anesthesia** (95.2%)

Conclusion

- irrespective of the type, the use of local anaesthesia of the skin was found to significantly reduce pain during puncture as compared to no local anaesthesia
- there was a trend towards better pain reduction in the EMLA-group
- overall pain during puncture was not relevant in terms of decision for or against future spinal anaesthesia in all groups
- our data supports the use of local anaesthesia prior to spinal puncture in daily routine