PRESCRIPTION OF ANTIMICROBIALS TO HOSPITALISED NEONATES AND CHILDREN IN JAPAN

Kotaro Araki¹, Takemi Murai¹, Kahoru Fukuoka¹, Yuta Aizawa¹, Junichi Suwa², Yuho Horikoshi¹

1 Division of Infectious Diseases, Department of Pediatrics, Tokyo Metropolitan Children’s Medical Center, Tokyo, Japan

2 Department of Pharmacy, Tokyo Metropolitan Children’s Medical Center, Tokyo, Japan

Introduction

The inappropriate use of antimicrobials is one of the main factors for the emergence of antimicrobial resistance. Antimicrobial stewardship programs have been introduced in many hospitals to optimize the use of antimicrobials. In order to effectively enforce, appropriate assessment of antibacterial usage is needed. This study aims to describe antimicrobial prescribing patterns at a tertiary care children’s hospital in Japan.

Methods

- **Study design** : one-point prevalence survey (PPS)
1st:May-June 2016, 2nd:September-October 2016, 3rd:December 2016-February 2017
A part of the Global Antimicrobial Resistance, Prescribing and Efficacy in Neonates and Children study
- **Setting** : Tokyo Metropolitan Children’s Medical Center in Japan
- **Inclusion** : children and neonates receiving antimicrobials on the day of PPS
- **Data collection** : antimicrobial agents, dose, frequency, route of administration, reasons for treatment

Results

Table1. Patient characteristics

	N=275	n (%)
Male	188	(68.4)
Children age(years) :Median(IQR)	264	(96.0) 6 (3-17.1)
Neonate	11	(4.0)

Table2. Details of prescription

	N=414	n (%)
Reasons for prescription		
Treatment	184	(44.4)
Prophylaxis	230	(55.6)
Route of administration		
Parental agents	191	(46.1)
Oral agents	223	(53.8)

Figure1. Prescription for treatment (n=184)

Figure2. Prescription for prophylaxis (n=230)

Figure3. Prescribed parental antimicrobial agents (n=191)

Conclusion

- Penicillins and 3rd generation cephalosporins were widely prescribed for treatment at a tertiary care children’s hospital in Japan.
- More than 50% were prophylactic administration.