Authors: Victoria Armenteros-Yeguas. Arantza Tomás-López. Estíbaliz Cristóbal-Domínguez. Inmaculada Moraza-Dulanto. Lucía Gárate-Echenique . Lara Meléndez-Fernández. Eva Merino-Romero. Erika Miranda-Serrano.

Integrated Health Organization (IHO) - Araba University Hospital. Basque Health Service. Vitoria-Gasteiz (Spain)

INTRODUCTION

The aim of this study is to evaluate the effectiveness of an intravenous therapy team (ITT) in relation to vascular care safety and satisfaction in multimorbid patients.

- SETTING:** IHO-Araba University Hospital and IHO-Alto Deba hospital. (Spain)
- STUDY PERIOD:** December 2013 - May 2015
- VARIABLES:** Demographics.
Patient satisfaction (visual analog scale)
The main variable: “Safety of vascular care”
- DATA ANALYSIS:** Absolute frequencies, percentages, means, standard deviation and relative rates per 1.000 days of treatment. Chi-square, Mann-Whitney U tests and incidence rates were used to compare groups.

- THE MAIN VARIABLE: “Safety of vascular care”**

 - Presence of difficult venous access (DIVA)
 - Number / kind of vascular access devices (VAD) needed
 - Locations of VAD
 - Number of attempts
 - Average duration of VAD
 - Number of adverse events

PATIENTS	CONTROL GROUP N=26	EXPERIMENTAL GROUP N=26	p
Men	19 (73.1%)	12 (46.2%)	0.048
Women	7 (26.9%)	14 (53.8%)	
Age mean	79.1 (SD:8.2)	80.4 (SD:10.4)	0.638
Presence of DIVA*	19 (73.1%)	23 (88.5%)	0.159

*DIVA: no visible or palpable veins and/or previous history of more than two attempts to insert an intravenous device

VARIABLES		CONTROL GROUP	EXPERIMENTAL GROUP				p*
			TOTAL	During delay time	ITT intervention	p**	
Number of VAD		51 (41.5%)	72 (58.5%)	43 (34.9%)	29 (23.6%)	<0.001	<0.001
Type of VAD	Peripheral	47 (92.2%)	45 (62.5%)	43 (59.7%)	2 (2.8%)	<0.001	<0.001
	Midline	3 (5.9%)	27 (35.5%) all inserted by ITT				
	PICC	1 (2%)	0				
Number of attempts	1	44 (93.6%)	36 (85.7%)				0.297
	2 or >	3 (6.4%)	6 (14.3%)	5 (11.9 %)	1 (2.4%)	0.011	
Average duration		5.4 (SD:3.9)	9.37 (SD 12.33)	2.8 (SD:1.39)	19.03 (SD:14.9)	<0.001	0.902

*inter-group (control and experimental group)
** intra-group (experimental group).

Delay time of ITT: 2.6 days (SD:1.8)

- Innappropriate sites:** Hand, wrist, leg, and cubital fossa.
- Innappropriate VAD for prescribed therapy :** Peripheral venous catheter for the administration of irritants or hyperosmolar substances.

ADVERSE EVENTS	CONTROL GROUP	EXPERIMENTAL GROUP				p*
		TOTAL	During delay time	ITT intervention	p**	
n (%)	21 (41.2%)	22 (30.6%)	19 (26.4%)	3 (4.2%)	0.002	0.224
n /1.000 catheter days	90.52	35.43				0.003

*inter-group (control and experimental group)
** intra-group (experimental group).

SATISFACTION	CONTROL GROUP	EXPERIMENTAL GROUP	p
Visual Analog Scale (0-10)	7.78 (SD: 0.89)	8.76 (SD: 1.38)	0.015

DISCUSSION & CONCLUSION

ITT provided more safety in relation to adequacy of VAD to the prescribed therapy and incidence of adverse events. Delay time influenced the intervention group negatively in relation to number of VAD, the average duration and the number of attempts to insert them.

REFERENCES
1.Brunelle, D. (2003). Impact of a dedicated infusion therapy team on the reduction of catheter-related nosocomial infections. *Journal of Infusion Nursing*, 26(6), 362-366.
2.Robinson, M. K., Mogensen, K. M., Grudinskas, G. F., Kohler, S., & Jacobs, D. O. (2005). Improved care and reduced costs for patients requiring peripherally inserted central catheters: the role of bedside ultrasound and a dedicated team. *Journal of Parenteral and Enteral Nutrition*, 29(5), 374-379.