THE PREVALENCE OF PSYCHOACTIVE SUBSTANCE USE AMONG ADOLESCENTS BEFORE THE FIRST PSYCHIATRIC CONSULTATION.

M. Pilecki¹, K. Krzykawska², A. Polańska², J. Ryniak², K. Wierba-Rak², K. Urbanek².

¹Jagiellonian University Medical College- Chair of Psychiatry, Department of Child and Adolescent Psychiatry, Krakow, Poland.

²Krakow University Hospital, The Department of Adult- Child and Adolescent Psychiatry, Krakow, Poland.

Objectives:

The aim of the study was to evaluate the frequency of psychoactive substance use among adolescents referred for psychiatric consultation and how it correlates with selected socio-environmental factors.

Background:

Studies provide clinical data of increasing psychoactive substance use and abuse in the population of Polish teenagers and frequent coexistence of substance use and psychiatric disorders.

Materials and Methods:

The study was conducted within 3 months of 2016. It included 60 patients applying for the first consultation to the Children and Adolescent Psychiatry Clinic of the University Hospital in Krakow. The study consisted of anonymous completion of the ESPAD (European School Survey Project on Alcohol and Drugs) standardized questionnaire. It was used to assess the prevalence of substance use and to analyze the associated factors. Afterwards the results were compared with the nationwide Polish data concerning adolescents above 16 years of age.

Results and Conclusions:

In the studied group, 52% were boys and 48% girls, 2% did not specify their gender. 68% of participants have 15 yr. or more. 7% did not specify their age. The use of alcohol among patients above the age of 16 in our study group was smaller, while the use of other psychoactive substances was significantly higher compared to the general population. In addition, there is a positive correlation between substance use and, among others, older adolescent's age, modern media use, reduction of interests (e.g. reading), time spent outdoors with friends and less parental knowledge about it. For the purpose of the analysis we use Spearman's rank correlation coefficient. For some part of the results the correlation was significant at the level 0.01 and for the other part at the level 0.05. There is a significantly greater prevalence of psychoactive substance use among patients seeking for psychiatric attention. For that reason this issue should be taken into account both during the medical interview and treatment programs.

The limitation of the study is that some of the teenagers could seek help due to problems with substances.

Comparison between the results of studies conducted at The Department of Adult, Child and Adolescent Psychiatry of Krakow University Hospital and the nationwide study in Poland in the older age group - over 16 years old.