THE COLOSTRUM-KIT

A quality improvement project implementing the evidence-based strategy to provide colostrum to the sick or premature newborn infant

Emilija Wilson^{1,2}; Ann Dsilna Lindh¹; Ia Roos¹; Maria Appelqvist¹; Jeanette Ekeberg¹; Eleni Ioannidou¹; Boubou Hallberg^{1,2}; Birgitta Berglund¹; Siri Lilliesköld¹; Dirk Wackernagel^{1,2}; Mireille Vanpee^{1,3}; Kajsa Bohlin^{1,2}; Anna Gustafsson^{1,2}

1.Department of Neonatology, Karolinska University Hospital, Stockholm, Sweden 2.Department of Clinical Science, Intervention and Technology, Karolinska Institutet, Stockholm, Sweden, 3.Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden

Background

In Sweden approximately 10% of all newborn infants need admission to neonatal care. Establishing and maintaining a milk production can be a major challenge for a mother whose child is admitted to neonatal care. Mother's own milk (MOM) provides several benefits for all newborn infants and colostrum may be applied to the oral mucosa as immunological protection.

Hypothesis and aim

We hypothesize that if parents are provided with information to be able to make an informed choice about colostrum, the proportion of infants receiving colostrum within the first hours of life will increase, leading to early stimulation of breast milk production, resulting in a higher rate of infants receiving mother's own milk (MOM) during hospitalization and at discharge from the neonatal intensive care unit (NICU).

Methods

Local lactation and nutrition teams at Karolinska University Hospital, Department of Neonatology, Stockholm developed an information brochure about early administration of colostrum and early initiation of breastmilk expression.

A "colostrum-kit", with all tools needed for colostrum expression and administration was composed and handed out together with the information.

Breastfeeding at discharge from NICU was investigated prior to implementation of the colostrum-kit and investigated four month after implementation.

Breastfeeding was defined as exclusive, partial and no breastfeeding regardless of route of administration

Exclusive breastfeeding	Only MOM
Partial breastfeeding	Formula in addition to MOM
No breastfeeding	No MOM

Results

Nutritional discharge data was available for 95,5% (171/179) of infants discharged home during the pre implementation period January to April 2018. During the post implementation period May to August 2018, 98,2% (167/170) had nutritional data available at discharge. After implementation the group of exclusively breastfed infants increased and less infants were partially breastfed. The group of no breastfeeding did not increase.

Breastfeeding at discharge before and after implementation of the kolostrum-kit

Conclusion

By implementation of the colostrum-kit, we support and inform parents based on best practice and existing evidence. We speculate that this quality improvement project may increase exclusive MOM rates at discharge as it supports early stimulation of breast milk production of those mother's who wish to breastfeed. Further analysis are required to be able to assess associations.

Next steps

Data on "time of initiation of breast milk expression" and "time of first colostrum" will be collected and the association with breastfeeding at discharge from NICU will be investigated as well as associated neonatal outcomes.

References: 1.Parker LA, et al. Effect of early breast milk expression on milk volume and timing of lactogenesis stage II among mothers of very low birth weight infants; a pilot study. Journal of perinatology (2012) 32, 205-209 2. Victora CG et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. Lancet 2016; 387: 475-90. 3. Hanson LA, Silfverdal SA. The mother's immune system is a balanced threat to the foetus, turning to protection of the neonate. Acta Paediatr. 2009 Feb;98(2):221-8. 4. <http://www.ihl.org/resources/Pages/Tools/PlanDoStudyActWorksheet.aspx>

Department of Neonatology, Karolinska University Hospital
in Huddinge, Solna and Danderyd

Corresponding author:

Emilija Wilson, RNRM, PhD
E-mail: emilija.wilson@sl.se

Karolinska
Institutet

KAROLINSKA
UNIVERSITY HOSPITAL