A rare case of a 34 years old parturient with elephantiasis

Maria Tzima, Paschalitsa Serchan, Argyro Fileli, Panagiota Bilali, Konstantinos Katsanoulas, Eleni Katsika

Anesthesiology department, Hippokrateion General Hospital of Thessaloniki, Greece

Elephantiasis nostras verucosa is a rare form of chronic lymphedema, which affects mainly gravity-dependent parts of the body. It causes progressive cutaneous hypertrophy and severe dysformation of the affected part. . The pathogenesis of elephantiasis is still unclear, although several conditions are found to be accused, such as trauma, filarial infections, radiation, neoplasms, chronic venous stasis, congestive heart failure, hypothyroidism. In our case, a 34 years old primigravita parturient is scheduled for elective cesarian section at 38th week of gestation. Elephantiasis appeared at the age of 7. After an ankle injury, an inflammation fired the appearance of the oedema. Initially, the oedema concerned only the right leg but the last year it appeared to the left leg, as well. Her condition began to deteriorate when she became pregnant. She gained 50 kilograms (her weight was 135kgr)most of them were due to the extensive leg oedema, which rendered her incapable even for walking. She presented diabetes and hypothyreoidism during her pregnancy. She was under thyroxine and she was euthyroidic. Her airway seemed manageable (Mallampati I-II). The cesarean section was performed under general anaesthesia, keeping low the airway pressure (up to 20mmHg) and PEEP 5mmHg.

Aspiration prophylaxis was given; ranitide 40 mg and metoclopramide 10mg IV. Rapid sequence induction was performed, using propofol 200mg and succinylcholine 100mg. Anaesthesia was maintained with O2/N2O (50%/50%) and sevoflurane 1%.

After the delivery of the baby, she received morphine 5 mg for analgesia, atracurium 50 mgr, paracetamol 1gr and ondasetron 4 mgr. The neonate's APGAR score was 7, 7, 10 at 1st, 5th, and 10th min. respectively, after delivery. She recovered well, without complications.

The targets in this case were to avoid worsening the oedema, and not to impair the lymphatic contractile activity. Performing regional anaesthesia, we were aware of vasodilation and fluid overloading for treating possible hypotension. On the other hand, local anesthetics have shown a decrease in lymphatic function, which could worsen the oedema. General anaesthesia was chosen, with low PEEP and low intrathoracic pressure, so as not to affect oedema and lymph flow.