Induced periimplantitis in a novel mini-pig model-
description and defect characterization

Umberto Demoner Ramos* (umbertodramos@gmail.com)
Barbara Furtado Masalskas
Arthur Belém Novaes Jr.

Ribeirão Preto School of Dentistry Ribeirão Preto-SP, Brazil
São Paulo Research Foundation

Abstract

This study has the original intention of compare treatments for induced Periimplantitis. The collected data until this time point will be presented in order to characterize the disease induction in this new model.

Methods: Eight Mini-pigs from the same breeder (Minipig br1) received eight implants each. After 8 weeks, healing abutments were installed and are 2 more weeks, silk ligatures were placed and left to accumulate plaque in association with a was wetted to a slurry food during 8 more weeks for plaque accumulation disease induction; after 8 weeks radiographic data when collected to evaluate bone loss and the ligature were maintained for additional 8 weeks, when a bone loss of at least 30% of the implant could be detected, the ligatures were removed ; after 3 more weeks, the clinical signs of disease were evaluated and periimplant bone loss was measured by x-rays. The treatment of the induced lesions were performed and the defects characterization was recorded trans-surgically. This study will show the characteristics of the defects and microtomographic data from two animals that died before the end of the study.

Results: The ligatures successfully induced progressive bone loss and ligature removal reduced the signs of acute inflammation. The progression rate was slower than the obtained in a previous study of our group in dog model. There were circumferential defects as well as horizontal bone loss, buccal and lingual bone dehiscence were not frequent and occurred mainly in the presence of horizontal bone loss. Vertical defects appeared mainly at interproximal areas ($2.84 \pm 2.35\text{mm}$). The defects configuration was similar to humans but, as this model does not control the pattern of bone loss, different defects can be found in the same experimental unit.

Conclusions: The use of mini-pig as a model to study periimplantitis produced defects comparable to human periimplantitis lesions with acute and chronic characteristics and might be a feasible alternative to the already described dog model.

Methods and Materials

Eight Mini-pigs from the same breeder (Minipig br1) had their mandibular P2, P3, P4 and M1 teeth extracted (fig A). After 12 weeks, each animal received eight implants, 4 each side of the jaw, with an interimplant distance of 7 mm minimum (figure B).

After eight weeks, healing abutments were installed and after 4 weeks, silk ligatures were placed and associated with soft diet in order to provoke plaque accumulation and inflammation (Figure C).

The ligatures remained in place for 6 weeks, when x-rays were taken to evaluate bone loss. At this time, bone loss was not sufficient for the second part of the study (not shown in this poster), thus, the ligatures were maintained for 6 additional weeks, when a bone loss of approximately 30% of the implant was achieved.

After disease induction, a flap was raised for the second part of the study (data not shown) and the defects were measured in order to characterize the disease induction achieved (figure D).

Results

Disease induction was successfully carried out with silk ligatures producing defects similars to the ones found in humans, with infra-crestal em supra crestal componentes (Table 1). It is advisable to use a minimum of 10 weeks of ligatures place to first evaluat boné loss and avoid excessive sedations of the animals. The Lesions resulted on bone loss and tissue inflammation with bleeding on probing similar to clinical findings in humans.

Table 1. Defects mesured transurgically

	Mesial	Distal	Buccal	Lingual
Total defect	3,25 ± 2,27	3,59±2,97	3,44±2,54	2,33±1,65
Infraosseous	2,18±2,74	2,85±2,35	2,30±2,22	1,74±1,20
Supraosseous	1,87±0,94	1,83±0,95	1,71±0,85	1,61±0,74

At the end of the experiment, 5 implants were lost., 4 of them in the same animal. The implants exhibited positive bleeding on probing and probing pocket dept of $4,55 \pm 0,93\text{mm}$ na Clinical Attachment level of $4,59 \pm 0,88\text{mm}$ (figure E).

Background and Aim

Periimplantitis is a common biologic complication in implant dentistry. Its prevalence ranges from 1 to 47% depending on the bone loss treshold(1). To date, there is no standard treatment protocolo well stabilished to clean and treat periimplantitis lesions and fully recover osseointegration, In addition, there is still a great gap in literature regarding the biologic process of bone loss after osseointegration and successful prosthetic rehabilitation.

The most common model to study periimplant biology and disease progression and treatment lies on a dog model of induced disease with ligatures(2). Despite its effectivity, this model is having its use reduced specially due to a social pressure of not using dogs for animal experiments and law regulations that made the experiments economically difficult. Thus, a new model with posibility of creating lesions similar the one found in humans is needed in order to study bone biology and disease progression and treatment.

The use of minipigs for studies is not new and has been used (3) previously to asses treatment response to induced periimplantitis. Still, there is no data on the characterization of the defects and disease progression.

Conclusion

Periimplantitis lesions induced with silk ligatures active for at least 12 weeks successfully induced periimplant lesions similar to the ones found in humans and may be a feaseble alternative to the dog model to study boné biology and treatment response

References

- Derks, J. & Tomasi, C. (2015) Peri-implant health and disease. A systematic review of current epidemiology. *J Clin Periodontol* 42 Suppl 16, S158-171. doi:10.1111/jcpe.12334.
- Lindhe, J., Berglundh, T., Ericsson, I., Liljenberg, B. & Marinello, C. (1992) Experimental breakdown of peri-implant and periodontal tissues. A study in the beagle dog. *Clin Oral Implants Res* 3, 9-16
- Singh G, Oneal RB, Brennan WA, Strong SL, Horner JA, Vandyke TE. Surgical Treatment of induced peri-implantitis-clinical and histological analisys. *Journal of periodontology* 1993;64.