EARLY BIOMARKER PATTERNS SPECIFIC TO INFANTS DEVELOPING SEVERE RETINOPATHY OF PREMATURITY

Chatarina Löfqvist^{1,2}, Eva M Andersson³, Bernhard Bührlen², Gunnel Hellgren^{2,4}, Karin Sävman⁵,
Lois EH Smith⁶, Ann Hellström²

¹Institute of Health and Care Sciences and ²Department of Ophthalmology, Institute of Neuroscience and Physiology and ³Department of Public Health and Community Medicine, Institute of Medicine and ⁴Institute of Biomedicine and ⁵Department of Pediatrics, Institute of Clinical Sciences, Sahlgrenska Academy at University of Gothenburg, Gothenburg, SWEDEN, ⁶Department of Ophthalmology, Harvard Medical School, Boston, US

Aim

The goal of this project was to identify early patterns of biomarkers, which predicted severe retinopathy of prematurity (ROP) needing treatment.

Background

Biomarkers have become essential to clinical studies of retinopathy of prematurity, yet there are no biomarkers which reliably or uniquely predict the subsequent clinical course.

Results

The best predictive biomarkers could be divided into four categories:
1) growth (Fig 1), 2) enteral nutrition (Fig 2), 3) oxygen use (Fig 3) and
4) biomarkers in blood (IGF and adiponectin) (Fig 4).

GROWTH

Figure 1: Schematic illustration of growth variables associated with development of severe ROP needing treatment.

Larger size of circle indicate stronger association.

NUTRITION

Figure 2: Schematic illustration of nutritional variables associated with development of severe ROP needing treatment.

Larger size of circle indicate stronger association, arrow indicate dependence between variables.

Study subjects and methods

A study in 90 infants with gestational age (GA) < 28 weeks with a mean (SD) gestational age at birth of 25.2 (1.4) weeks and mean (SD) birth weight of 882 (220) g. ROP was determined according to the International ROP classification. ROP stages were assessed in 78 infants and grouped as ROP stage 0 (n=17), ROP stage 1 (n=8), ROP stage 2 (n=22) and ROP stage 3 and/or treated (n=31).

Longitudinal (up to 10 measurements per infant) data including clinical as well as blood biomarkers were analysed. It was assessed if these predictors were associated with severe ROP that needed treatment.

Funding Source: This study is supported by grants provided by the Swedish Research Council (DNR# 2011-2432) and Gothenburg County Council (ALFGBG-426531) long-term support by De Blindas Vänner, Kronprinsessan Margaretas Arbetsnämnd för synskadade,, European Commission FP7 project 305485 PREVENT-ROP.

Conclusions

We have identified biomarkers, clinical as well as in blood, that will serve as a basis for development of an algorithm that can be used to safely and timely to identify infants at risk for severe ROP needing treatment on an individual level.

OXYGEN USE

Figure 3: Schematic illustration of oxygenation variables associated with development of severe ROP needing treatment.

Larger size of circle indicate stronger association.

BIOMARKERS IN BLOOD

Figure 4: Schematic illustration of blood-based variables associated with development of severe ROP needing treatment.

Retinopathy of prematurity disease progression – rational for longitudinal pattern analysis

The eye starts to develop at about 16 weeks of pregnancy, when the blood vessels of the retina begin to form at the optic nerve in the back of the eye. The blood vessels grow gradually toward the edges of the developing retina, supplying oxygen and nutrients. During the last 12 weeks of a pregnancy, the eye develops rapidly. When a baby is born full-term, the retinal blood vessel growth is mostly complete. If a baby is born prematurely, before these blood vessels have reached the edges of the retina, normal vessel growth may stop. The periphery of the retina might then send out signals to other areas of the retina for nourishment. As a result, new abnormal vessels might begin to grow. These new blood vessels are fragile and can leak, leading to retinal detachment and blindness if not identified and treated timely.

www.rop.gu.se