Pain and Depression in Rheumatologic Patients

N.P. Chernus¹, Gorenkov R. B¹, S.K. Sivkova¹, A.S. Sivkov¹, A.M. Zolotovickaja¹, T. S. Savina¹.

1 - I.M. Sechenov First Moscow State Medical University: Moscow, Russia

Background: Chronic recurrent pain is a clinical symptom of rheumatologic diseases with primary joint damage. The mechanisms of chronic pain syndrome development are often associated with depression.

Methods: The study enrolled 67 patients with osteoarthritis (OA), rheumatoid arthritis (RA) and lower back pain. The average subject age was $59, 5 \pm 5, 3$; 24 were males and 43 were females. In order to achieve the study objective, Luscher test, CES-D; TAS; Spielberg-Hanin's scale; MMPI; VAS scale and McGill Pain Questionnaire were used.

Results: Following the testing results, 32 (47,8%) of 67 patients were diagnosed with chronic pain syndrome associated with depression respectively $p=0,002$. The average pain syndrome (PS) level by VAS scale in patients with depression was almost 1,7 times higher than in patients without psychological disorders, $8,6 \pm 0,9$ vs. $5,2 \pm 1,09$; $p < 0,001$. Under adaptive tension conditions, when the MMPI profile exceeded 70 T-scores, PS tended to exacerbate ($p < 0,001$), while manifestation of depression symptoms correlated with PS ($r = 0,627$) and the presence of personality traits accentuation ($r = 0,531$), along with increased anxiety ($r = 0,425$), asthenic condition ($r = 0,392$) and general TAS alexithymia score ($r = 0,473$). After the course of therapy involving antidepressant drugs, PS decreased by 2,9 times as compared to the baseline level, $p < 0,001$.

Conclusion: Indications for antidepressant use in patients with PS include the chronic pain syndrome associated with depression at the background of basic anti-inflammatory therapy.