Late Diagnosis of HIV in Northern Ireland

SEJ Todd, E Walker, M Hunter, CM Donnelly, WW Dinsmore, CR Emerson, SP Quah, EJ McCarty

Department of Genito-Urinary Medicine, Belfast HSC Trust, Royal Victoria Hospital, Belfast

Background: Highly active antiretroviral therapy (HAART) has transformed the outcome for HIV, however, there continues to be significant morbidity and mortality especially with late diagnosis (CD4 count <350 cells/mm³). A national audit by British HIV Association (BHIVA) in 2006 showed that of deaths occurring amongst HIV positive adults in the UK, 24% were directly attributable to the diagnosis of HIV being made too late for treatment¹. Furthermore, many of the late diagnoses had been seen in recent past by healthcare professionals².

Late diagnosis of HIV infection has been associated with increased mortality and morbidity³, impaired response to HAART⁴ and increased cost to healthcare services⁵. Late diagnoses across UK declined from 57% (2004) to 39% (2015), however, a review within our region in 2013-14 revealed much higher proportion of late diagnoses than UK average, with multiple missed opportunities for testing in majority cases. Data was presented at educational meetings and feedback given to clinicians when delay in diagnosis occurred. We sought to assess the impact of late diagnoses and mortality within our population.

Methods: A retrospective chart analysis of all new HIV diagnoses from March 2015 - February 2016 (Period 2) to determine proportion of late diagnosis, missed opportunities for testing and mortality. Comparison was made with results of a previous review during July 2013 - June 2014 (Period 1).

Results: There were 76 new HIV diagnoses during period 1 and 71 in period 2. Late diagnosis (CD4 count <350 cells/mm³) decreased from 59% (45/76) to 49% (35/71). Those that had a CD4 count <200 cells/mm³ decreased from 41% (31/76) to 31% (22/71) (Figure 1A).

The majority of late diagnoses were male; 76% in period 1 and 86% in period 2. The percentage of late diagnoses, born in the UK decreased; 70% in period 1 and 66% in period 2 (Figure 1B).

Figure 1 - A. Total number of new HIV diagnoses and number of patients with CD4 count <350 and <200 cells/mm³ during period 1 (July 2013-June 2014) and period 2 (March 2015-February 2016); B. Late diagnoses of HIV: Sex and % patients born in UK during period 1 and 2; C. Late diagnoses of HIV: Mode of acquisition during period 1 and 2; D. Number of patients with AIDS defining condition at diagnosis.

The proportion diagnosed through Genitourinary Medicine (GUM) increased from 24% (period 1) to 34% (period 2). The remainder were diagnosed in a wide range of other specialities, most commonly general medicine and general practice settings (Figure 2). Mode of transmission in period 1 and 2 respectively; Men who have sex with men (MSM) 49% vs 71%, heterosexual 47% vs 23%, intravenous drug use (IVDU) 7% vs 11% (Figure 1C).

References.

1. British HIV Association (BHIVA). 2005–6 mortality audit. <http://www.bhiva.org/files/file1001379.ppt>
2. Sullivan AK, Curtis H, Sabin CA et al. Newly diagnosed HIV infections: review in UK and Ireland. BMJ, 2005, 330, 1301 –2. <http://www.bmj.com/cgi/content/full/330/7503/1301>
3. Health Protection Agency (HPA), Centre for Infections. The UK Collaborative Group for HIV and STI Surveillance. Testing Times. HIV and other sexually transmitted infections in the United Kingdom: 2007. http://www.hpa.org.uk/web/HPAweb&HPAwebStandard/HPAweb_C/12030843559418.
4. Stöhr W, Dunn DT, Porter K et al. on behalf of the UK CHIC Study. CD4 cell count and initiation of antiretroviral therapy: trends in seven UK centres, 1997–2003. HIV Medicine, 2007, 8, 135–41.
5. Krentz HB, Auld MC, Gill MJ. The high cost of medical care for patients who present late (CD4<200 cells/μL) with HIV infection. HIV Medicine, 2004, 5, 93 –8.

Figure 2. Late diagnoses of HIV: Diagnosis by each specialty

Prior to diagnosis, number of patients presenting to other healthcare settings with clinical indicator diseases significantly decreased from 84% (38/45) in period 1 to 49% (17/35) period 2. Most common indicators included oral candidiasis, unexplained weight loss and blood dyscrasias. 57% (25/45) in period 1 and 57% (20/35) in period 2 attended care for investigation prior to diagnosis (Figure 3). Most common investigations requested by healthcare physicians included auto-immune and coeliac screen. The number of patients with acquired immunodeficiency syndrome (AIDS) defining conditions at diagnosis significantly decreased from 44% (20/45) in period 1 to 34% (12/35) period 2 (Figure 1D).

Figure 3. Late diagnoses of HIV: Type of investigation requested prior to diagnosis

The proportion of patients diagnosed in hospital decreased from, 44% (20/45) in period 1 to 37% (13/35) in period 2. Average inpatient stay was 19 days in period 1 and 31 days in period 2. Overall mortality more than halved from 7% (4/45) in period 1 to 3% (1/35) period 2. 91% (41/45) in period 1 and 97% (34/35) in period 2 commenced HAART (Figure 4).

Figure 4. Late diagnoses of HIV: Number of patients diagnosed as hospital inpatient, average hospital stay, number requiring intensive care treatment, mortality and number that commenced antiretroviral treatment (ARVT)

Conclusion & Recommendations: Whilst there has been a decrease in number of late diagnoses and mortality rate, the proportion being diagnosed late remains higher in Northern Ireland compared to other UK regions. 66% of late diagnoses presented to a wide range of healthcare settings with over half being investigated prior to diagnosis. Unfortunately, opportunities for early testing are still being missed and ongoing education is required. Continued education regarding clinical indicator conditions and HIV testing at educational meetings to hospital physicians and general practitioners is needed. Individual feedback should be given to clinicians when delay in diagnosis occurred.