Utilization and effectiveness of multimodal pain management strategies in spine surgery.

Jashvant Poeran MD PhD¹, Ottokar Stundner MD, MBA², Crispiana Cozowicz MD³, Nicole Zubizarreta MPH¹, Mary Hargett BSciEd³, Eric Schwenk MD¹, Madhu Mazumdar PhD¹, Stavros G. Memtsoudis MD PhD FCCP^{1,3}

1) Institute for Healthcare Delivery Science, Department of Population Health and Science, Icahn School of Medicine at Mount Sinai, New York, NY, USA

2) Department of Anesthesiology, Paracelsus Medical University, Salzburg, Austria

3) Department of Anesthesiology, Hospital for Special Surgery, New York, NY, USA

BACKGROUND

Given evidence supporting multimodal analgesia in spine surgery, and the lack of population-based data, we investigated 1) utilization patterns in spine surgery, and 2) associations between number of analgesic modes utilized and opioid utilization dose (assuming 10% reduction as clinically significant), complications and resource utilization.

METHODS

Retrospective cohort study of posterior lumbar fusion patients (Premier national claims database; 2006-2016). Opioid only analgesia versus multimodal analgesia (systemic opioids + acetaminophen, steroids, gabapentin/pregabalin, ketamine, NSAIDs, COX-2 inhibitors, intrathecal or epidural analgesia), categorized into 1, 2, or 2+ additional analgesic modes. Multilevel multivariable models measured associations between multimodal analgesia and opioid prescription dose, cost/length of hospitalization, and opioid-related adverse events (respiratory, gastrointestinal, genitourinary, central nervous system complications).

RESULTS

N = 265,538 cases, while 61.1% (n=162,156) received multimodal analgesia. Multivariable models showed that multimodal analgesia versus opioids only analgesia was not associated with reductions in opioid utilization dose, while a slight increase was observed: opioid dose effect estimates for those receiving 1, 2 and >2 additional modes were +2.2% (CI +1.3% to +3.2%), +5.2% (CI +3.9% to +6.5%), and +6.5% (CI +4.4% to +8.6%), respectively; all $P < 0.0001$. This coincided with similar patterns (null-effects in terms of potential benefits of multimodal analgesia) for opioid-related adverse events and resource utilization.

CONCLUSIONS

In contrast to evidence from joint arthroplasties, multimodal analgesia was not associated with theoretically expected outcome-benefits in spine surgery. While further research is needed, contributors to this null-effect may include, high baseline opioid utilization and limited use of non-opioid analgesic modes in spine surgery.