THE ANTISECRETORY FACTOR AND PRETERM BIRTH

Gustafsson A^{1,2}, Henckel E^{1,2}, Johansson E^{3,4}, Oshalim M^{3,4}, Lange S^{3,4},
Bernhardsson AK^{1,5}, Olin A⁵, Brodin P⁵, Bohlin K^{1,2}

¹Department of Neonatal Medicine, Karolinska University Hospital, Stockholm

²Department of Clinical Science, Intervention and Technology, Karolinska Institutet

³Department of Infectious Diseases, Institute of Biomedicine, Gothenburg University, Gothenburg

⁴Clinical Microbiology, Sahlgrenska University Hospital, Gothenburg

⁵Science for Life Laboratory, Department of Women's and Children's Health, Karolinska Institutet
SWEDEN

Conclusion:

- Following preterm birth, levels of active AF appear to be lower in both cord blood and placenta.
- The regional differences of AF in placenta, in this pilot study, are in line with markers of oxidative stress previously described following preterm preeclampsia.
- Low levels of active AF may suggest ongoing inflammatory processes.
- Dietary supplementation of AF is available today and further insight into the role of AF in prematurity may serve as the basis for new treatments to prevent preterm birth and its consequences.

Introduction:

Antisecretory factor (AF) is an endogenous protein involved in the innate regulation of secretory and inflammatory processes.

Inflammation contributes to preterm labour and various complications of prematurity.

We have previously described lower levels of AF associated with increased markers of inflammation in preterm placental tissue compared to term (Gustafsson et al 2018).

The objective in this ongoing study is to further describe the involvement of AF in preterm birth.

Results:

The level of AF in cord blood was lower in the preterm (n=14) compared to the term (n= 10) group (p=0.02).

A pilot of multiple samples from 5 placentas (3 preterm and 2 term) exhibited lower AF-levels in peripheral biopsies of preterm placentas (p=0.01). In term placenta, the central and peripheral part had similar AF-levels (p=0.86).

Material and methods:

- Preterm (<30 weeks gestation) and term control infants born at Karolinska University Hospital, Sweden had umbilical cord blood collected.
- Placental tissue biopsies, central and peripheral part, were taken from a subgroup of deliveries.
- Sandwich ELISA to determine active AF in plasma and placental tissue
- Proteasome subunit AF / C3 complex ELISA

Karolinska
Institutet

KAROLINSKA
UNIVERSITY HOSPITAL