Outcomes of Embedded Palliative Care Outpatients Consults on Timing of Palliative Care Access, Symptoms, and End-of-Life Quality Indicators among Advanced Non-Small Cell Lung Cancer Patients

Sriram Yennurajalingam MD, MS, Bernard Prado MD, Zhanni Lu MPH, Syed Naqvi MD, Janet L. Williams MPH, Taekyu Lim MD, Eduardo Bruera MD.

What we learned: Embedded PC initial consults provide earlier access to PC, and more frequent and earlier advanced care planning discussions.

Introduction

- Outpatient PC improves patient-reported and healthcare utilization outcomes among advanced NSCLC patients. However, the best consultation model of outpatient PC delivery is not yet defined.
- Two types of outpatient PC consults have been described: stand-alone PC consults (OPC) referred to an independent functioning PC specialty clinic; and embedded PC consults (EPC) referred to PC professionals who are physically embedded into an oncology clinic. Limited studies have compared the outcomes of initial EPC with initial OPC.
- Between August 1, 2012, and June 30, 2013, a board-certified PC specialist and a PC-trained nurse of our team were embedded in the thoracic oncology clinic at our institution to improve advanced lung cancer patients access to PC.

Study Aim:

- To determine the timing of PC access, symptoms and end-of-life (EOL) quality care outcomes of advanced NSCLC patients referred to initial EPC as compared to those referred to initial OPC.

Methods

- Design: Retrospective study of a random sample of EPC (Aug, 2012 - Jun, 2013) and OPC (Jan 2009 – Jul 2012). Eligibility criteria: (a) diagnosis of advanced cancer (locally advanced, recurrent or metastatic); (b) ≥18 years; (c) NSCLC histology.
- EPC referral criteria were used by oncologists to adjust to the local resources: severe symptoms, distress, or an estimated prognosis of ≤ 6 months. After first consultation, patients were seen at the OPC clinic.
- OPC clinic is staffed with an interdisciplinary team (a PC specialist, a PC-trained nurse, a social worker, a pharmacists, a nutritionist, a counselor, and a chaplain). All referrals are at oncologists discretion.
- Baseline features, symptoms (Edmonton System Assessment Scale), and EOL quality care outcomes were collected: number of ICU deaths, EC visits, ICU admissions, hospitalizations and length of hospital stay within 30 days before death; cancer treatments within 14 days before death; documentation of advanced care planning (ACP), and hospice discussions, do not resuscitate (DNR) order at first follow-up, completion of hospice referrals, and advance directives
- The time from referral to first consult, ACP date to death, overall survival from first consult to death, and number of encounters/patient were also retrieved.

Statistical Analysis

- Categorical variables were summarized using percentages and frequencies and differences analyzed using a two-sided Chi-square/Fisher’s exact test. Continuous variables were summarized using median and interquartile (IQR).

Table 1: Patient baseline characteristics

Covariates	Total: (n=340, 100%)	EPC (n=147, 43.2%)	OPC (n=193, 56.8%)	p ^a
Age				
Median (IQR)	62(53.3, 69.0)	61(53, 68.5)	63(54, 70)	.13
Gender				
Male	182(53.5)	77(52.4)	105(54.4)	.71
Race				
White	241(70.9)	109(74.1)	132(68.4)	.048
Black	48(14.1)	14(9.5)	34(17.6)	
Hispanic	31(9.1)	16(10.9)	15(7.8)	
Asian	13(3.8)	3(2.0)	10(5.2)	
Other	7(2.1)	5(3.4)	2(1.0)	
CAGE				
Positive (%)	60(18)	24(16)	36(19)	.6
ESAS				
Median (IQR)				
Pain	6(3, 8)	6(4, 8)	5(3, 8)	.038
Fatigue	6(4, 8)	6(4, 8)	6(4, 8)	.95
Nausea	0(0, 3)	0(0, 3)	1(0, 4)	.41
Depression	3(0, 5)	2(0, 5)	3(1, 5)	.12
Anxiety	3(0, 6)	3(0, 5)	3(1, 6)	.41
Drowsiness	3(1, 6)	3(1, 6)	4(1, 6)	.92
Appetite	5(2, 7)	5(1, 7)	5(3, 7)	.09
Dyspnea	5(2, 7)	5(2, 7)	4(2, 7)	.50
Well Being	5(3, 7)	5(3, 7)	5(3, 7)	.79
Sleep	5(3, 8)	5(2, 8)	5(3, 8)	.42
SDS	37(27, 49)	36(27, 50)	38(2, 48)	.71
MDAS				
Median (IQR)	1(0, 2)	1(1, 2)	1(0, 3)	.49
ECOG				
Mean (SD)	2.0(0.8)	2.2(0.8)	1.9(0.8)	<.001
Comorbidities				
0-2	249(73.2)	106(72.1)	143(74.1)	.68
≥3	91(26.8)	41(27.9)	50(25.9)	
Clinical encounters ^b				
Median (IQR)	2.0(2.0, 4.8)	2(1, 4)	3(2, 5)	<.001
First follow-up				
Yes	267(78.5)	95(64.6)	172(89.1)	<.001
Second follow-up				
Yes	158(46.5)	61(41.5)	97(50.3)	.11

Abbreviations- EPC: Embedded Palliative care clinic consults, OPC- stand-alone outpatient palliative care clinic, ESAS: Edmonton Symptom Assessment Scale; SD: Standard Deviation, CAGE: Cut down, Annoyed, Guilty, and Eye-opener; MDAS: Memorial Delirium Assessment Scale; SDS: Symptom Distress Score. ECOG: performance Status; a: Independent Sample T test for means of age, ESAS, MDAS and ECOG; Mann-Whitney U for medians of age, ESAS, MDAS and ECOG; Chi-square or Fisher’s exact test for other variables; b: per patient

Table 2: Symptom Changes in EPC and OPC groups

Median (IQR); N=340	EPC				OPC				
	Baseline	Follow-up	Difference	p ^a	Baseline	Follow-up	Difference	p ^a	p ^b
Pain	6(4, 8)	5(3, 8)	-1(-3, 1)	.07	5(3, 8)	5(2, 7)	0(-2, 1)	.037	.65
Fatigue	6(4, 8)	5(3, 8)	0(-2, 1)	.34	6(4, 8)	6(4, 8)	0(-2, 2)	.93	.43
Nausea	0(0, 3)	0(0, 4)	0(-1, 1)	.74	1(0, 4)	1(0, 4)	0(-1, 1)	.76	.64
Depression	2(0, 5)	1(0, 4)	0(-2, 0)	.12	3(1, 5)	3(0, 5)	0(-2, 1)	.66	.21
Anxiety	3(0, 5)	2(0, 4.5)	0(-2, 0)	.05	3(1, 6)	3(0, 6)	0(-2, 2)	.67	.05
Drowsiness	3(1, 6)	3(0, 6)	0(-2, 1)	.37	4(1, 6)	3(1, 6)	0(-2, 2)	.85	.29
Appetite	5(1, 7)	4(2, 7)	0(-2, 1)	.86	5(3, 7)	5(3, 7)	0(-2, 2)	.72	.77
Dyspnea	5(2, 7)	3(1, 6)	-1(-2, 0.5)	.014	4(2, 7)	4(2, 6)	0(-2, 1)	.42	.039
Well Being	5(3, 7)	5(3.3, 7)	0(-2, 1)	.96	5(3, 7)	5(3, 7)	0(-2, 2)	.57	.54
Sleep	5(2, 8)	5(2.3, 7)	0(-2, 1)	.55	5(3, 8)	5(3, 7)	-1(-2.5, 1)	.08	.25
SDS	36(27, 50)	33(20, 43)	-3(-13, 3)	.05	38(27, 48)	36(24, 50)	-2(-12, 9)	.53	.22

Abbreviations- ESAS: Edmonton Symptom Assessment Scale; IQR: Interquartile; EPC: Embedded Palliative Care consults; OPC: stand-alone Outpatients Palliative Care consults a: Comparison between baseline and first follow-up; Mann-Whitney Test. b: Comparison of the changes from baseline to first follow-up between groups; Mann-Whitney Test.

Table 3: End-of-Life Quality Care Outcomes between EPC and OPC cohorts

	Total: (N=340, 100%)	EPC (N=147, 43.2%)	OPC (N=193, 56.8%)	p*
EC visits 30 days before death				
Yes	97(50.5)	38(53.5)	59(48.8)	.52
Death in ICU				
Yes	11(6.2)	4(6.0)	7(6.3)	1.0
ICU admission 30 days before death				
Yes	26(14.5)	8(12.1)	18(15.9)	.48
Hospitalizations 30 days before death				
Yes	116(57.7)	116(57.7)	116(57.7)	.42
Hospital LOS 30 days before death, Days				
Median (IQR)	3.0(0.0, 8.0)	3.0(0.0, 8.5)	1.0(0.0, 8.0)	.31
Advance directive completion				
Yes	171(50.3)	67(45.6)	104(53.9)	.12
Advanced care planning discussion				
Yes	225(66.4)	114(77.6)	111(57.8)	<.001
DNR at 1 st . follow-up				
Yes	14(4.1)	8(5.4)	6(3.1)	.28
Hospice discussion				
Yes	160(47.1)	62(42.2)	98(50.8)	.11
Hospice referral				
Yes	129(37.9)	50(34.0)	79(40.9)	.19
Cancer Treatment				
Yes	53(27.70)	23(31.50)	30(25.40)	.36
Chemotherapy 14 days before death				
Yes	6(3.1)	3(4.1)	3(2.5)	.67
Radiation therapy 14 days before death				
Yes	23(12.0)	13(17.8)	10(8.5)	.05
Surgery 14 days before death				
Yes	1(0.5)	1(1.4)	0(0.0)	.38
Immunotherapy 14 days before death				
Yes	0(0.0)	0(0.0)	0(0.0)	-
Target therapy 14 days before death				
Yes	24(12.5)	7(9.6)	17(14.3)	.33

Abbreviations-EC: Emergency Care; ICU: Intensive Care Unit; DNR: Do not resuscitate. Mo: Months. *: Mann-Whitney U; Independent Sample T test for continuous variables in End of Life Care; Chi-square or Fisher’s exact test for categorical variables.

- The variation of symptoms, and changes of ESAS items from first PC consult to first follow-up between groups were analyzed using Wilcoxon rank sum test.
- Variation of the EOL outcomes between cohorts were examined using the two-sided Chi-square/Fisher’s exact test. Length of hospital stay was tested using Wilcoxon rank sum test between the two groups.
- We described differences on time using median and IQR, and used Wilcoxon rank sum test to analyze between-group differences. Survival was analyzed with the Kaplan-Meier method. $P \leq 0.05$ was considered statistically significant.

Results

- 340 were evaluable (EPC= 147). At baseline, mean ECOG (2.2 vs 1.9, p<.001), and median pain (6 vs 5, p=.038) were higher in EPC.
- The median number of clinical encounters/patient (2 vs 3), and the proportion of patients attending their first follow-up visit (64% vs 89%) were higher in OPC (**Table 1**).
- Time from referral to first consult was shorter (median 0 vs 7 days, p<.001) and dyspnea was better in EPC at follow-up (-1 vs 0, p=.039). (**Table 2**).
- A higher proportion in EPC (77% vs 58%, p<.001) (**Table 3**) had ACP discussions and these occurred earlier (median 4 vs 1 month before death, p<.001).
- EPC group showed trends towards improved anxiety scores.

Conclusions

- As compared to OPC, EPC accessed PC earlier.
- However, no other significant differences in symptom control or EOL outcomes were found, except for improved dyspnea at follow-up, and more frequent, and earlier ACP discussion
- Further larger prospective studies are needed to compare the outcomes of various consultation models of outpatient PC delivery.

Email: syennu@mdanderson.org

Twitter Handle: @Palliativeconf