First shock delays in out of hospital cardiac arrest

García García JJ, Cuervo Menéndez JM, Fente Álvarez AI, García Castro A, Buelga Díaz C, Roza Alonso CL
SAMU-Asturias (EMS, Spain)

Purpose of the study

To determine the time from call to first shock in OHCA assisted by our ALS team.
To study the time intervals involved in it: time from call to dispatch, time from dispatch to switching on the monitor, and time from monitor switched on to first shock delivery.

Material and methods

Descriptive, observational, retrospective study.
Study period: January 1st 2014 to December 31st 2016.
Inclusion criteria: OHCA with initial shockable rhythm and event log available assisted by SAMU-Asturias.
Exclusion criterion: OHCA witnessed by ALS teams.
Data collected from clinical records, both in paper and digital, and event log from the monitor/defibrillator Lifepak 15®.

Statistical analysis with SPSS Statistics 21 software.

Results

Out of 1706 OHCA assisted by our EMS 126 (7.38%) meet inclusion criteria.
Mean interval between call and first shock is 14 m 42 s (SD=385 s). Figure 1 compares our results, using the intervals defined by Tanigawa et al, with theirs. Our findings show a delay over 13 m in 54.0% of cases, being lower than 9 m in 15.3% of cases and bigger than 20 min in 13.7%.

TABLE I: Other intervals

Call to dispatch ¹	n (%)	Dispatch to monitor on ²	n (%)	Monitor on to shock ³	n (%)
00'00" - 00'59"	19 (15.7%)	00'00" - 04'59"	17 (13.9%)	00'00" - 00'59"	19 (15.1%)
01'00" - 01'59"	20 (16.5%)	05'00" - 09'59"	67 (54.9%)	01'00" - 01'59"	45 (35.7%)
02'00" - 02'59"	36 (29.8%)	10'00" - 14'59"	20 (16.4%)	02'00" - 02'59"	43 (34.1%)
03'00" - 03'59"	22 (18.2%)	15'00" - 19'59"	13 (10.7%)	03'00" - 03'59"	11 (8.7%)
04'00" - 04'59"	8 (6.3%)	More than 20 min	5 (4.1%)	04'00" - 04'59"	5 (4.0%)
05'00" - 05'59"	6 (4.8%)			05'00" - 05'59"	2 (1.6%)
More than 6 min	10 (8.3%)			More than 6 min	1 (0.8%)
¹ mean = 02'15" (SD=78") 5 missed cases		² mean = 10'21" (SD=343") 4 missed cases		³ mean = 02'08" (SD=68")	

Table I resumes the other intervals studied.
Mean interval between call and ALS team dispatch is 2 m 15 s (SD=78 s).
Mean interval between ALS team dispatch and when the monitor is switched on is 10 m 21 s (SD=343 s).
Mean interval between when the monitor is switched on and the first shock is 2 m 8 s. (SD=68 s) (Figure II).

Conclusions

- OHCA have specific difficulties that imply delays in the first shock delivery (patients hard to reach, excess of clothes, unsafe area) that have to be studied.
- Our intervals "call to first shock" seem to be better than in other published studies.
- Our interval "call-ALS activation" shows an improvement area in order to make it closer to published standards.
- Interval variability can be due to the absence of a fixed procedure regarding the moment to turn the monitor on in our units.
- Rates of time "monitor on-shock" higher than two minutes have to be studied further.

Bibliography:

1. Perkins GD et al. ERC Guidelines for Resuscitation 2015: Section 2. Adult basic life support and automated external defibrillation. Resuscitation. 2015 Oct;95:81-99.
2. Tanigawa K, Tanaka K, Shigematsu A. Outcomes of out-of-hospital ventricular fibrillation: their association with time to defibrillation and related issues in the defibrillation program in Japan. Resuscitation. 2000 Jul;45(2):83-90.