IMPACT OF AN INTERACTIVE ONLINE MODULE FOR MEDICAL STUDENTS ON THE MANAGEMENT OF AGITATED PATIENTS IN THE PSYCHIATRIC EMERGENCY SETTING

C. Zhou¹, C. Murphy¹, M. Esliger¹, M. Kara².

¹Dalhousie University, Psychiatry, Halifax, Canada.

²Nova Scotia Health Authority- Research Methods Unit, Department of Medicine, Halifax, Canada.

OBJECTIVES

We developed an interactive video simulation of an increasingly agitated patient, followed by an online learning module as a training tool. Our hypothesis is that this new learning tool will improve medical students' confidence in management of agitated patients in the Psychiatric Emergency setting.

BACKGROUND

Adequate training in management of agitated patients is paramount for the safety of the patient and the staff. One difficulty in ensuring adequate training lies with the variability of occurrence and high-stakes nature of agitation. Many residents do not feel adequately prepared to manage agitated patients in the emergency, indicating insufficient exposure during medical clerkship. A search of databases (including MedEdPORTAL, PsycINFO, and PubMed) showed minimal simulations or teaching tools for residents regarding management of agitation, and none for medical students.

MATERIALS AND METHODS

Study design: Single group, repeated measures design. The outcome measures are self-reported scales of comfort and competence with different aspects of identifying and managing the acutely agitated patient. The change in self-reported level is the subjective outcome measure of interest. The three points of assessment are pre-module (T0), immediately post-module (T1), and two weeks post-module (T2).

ACKNOWLEDGEMENTS

Thanks to Dalhousie Psychiatry Medical Education Coordinator, Mandy Esliger, for her assistance in developing the online module

RESULTS

For subjective ratings of comfort, in pairwise comparison with Tukey adjustment for multiple comparisons, the difference (5.797, SE 0.8896, $p < 0.001$) between T0 and T1 and between T0 and T2 (6.35, SE 0.97, $p < 0.001$) were statistically significant. The difference between T1 and T2 were not statistically different (0.5564, SE 0.9952, $p = 0.8420$). For subjective ratings of competence, in pairwise comparison with Tukey adjustment for multiple comparisons, the difference between T0 and T1 (5.40, SE 0.93, $p < 0.001$) and the difference between T0 and T2 (6.15, SE 1.01, $p < 0.0001$) were statistically significant. The difference between T2 and T1 (0.75, SE 1.03) was not statistically different ($p = 0.7474$)

CONCLUSIONS

Data shows that medical students' level of comfort and subjective sense of competence with having an approach to and management of an agitated patient are significantly improved from T0 to T1, and these improvements are maintained at T2.

This study suggests that the educational modality of an interactive online module is a low-cost, significantly impactful method to help develop medical students' confidence with managing agitated patients.

LITERATURE CITED

- Dvir Y, Moniwa E, Crisp-Han H, Levy D, Coverdale JH. Survey of threats and assaults by patients on psychiatry residents. *Acad Psychiatry*. 2012;36(1):39-42.
- Finch S. Beyond Haloperidol: Teaching Emergency Medicine Residents to Manage Acute Agitation and Aggression in the Emergency Department. *Can J Psychiatry*. 2004;49(8):574
- Hirschbein LD., Gay T. Case-Based Independent Study for Medical Students in Emergency Psychiatry. *Academic Psychiatry*. 2005;29(1):96-99.
- Keltner N, Grant J, McLernon D. Use of Actors as Standardized Psychiatric Patients: Facilitating Success in Simulation Experiences. *J Psychosoc Nurs Ment Health Serv*. 2011;49(5):34-40.
- McGaghie WC, Draycott TJ, Dunn WF, Lopez CM, Stefanidis D. Evaluating the Impact of Simulation on Translational Patient Outcomes. *Simulation in healthcare: Journal of the Society for Simulation in Healthcare*. 2011;6
- Vestal H, Sowden G, Nejad S, Stoklosa J, Valcourt S, Keary C, Caminis A, Huffman J. Simulation-Based Training for Residents in the Management of Acute Agitation: A Cluster Randomized Controlled Trial. *Academic Psychiatry*. 2017;41(1):62-67.
- Zigman D, Young M, and Chalk C. Using Simulation to Train Junior Psychiatry Residents to Work With Agitated Patients: A Pilot Study. *Academic Psychiatry*. 2013;37(1):38-41

Figure 1: Subjective ratings of comfort at pre-module (T0), immediately post-module (T1), and two weeks post-module (T2).

Figure 2: Subjective ratings of competence at pre-module (T0), immediately post-module (T1), and two weeks post-module (T2).