Thrombogenesis in Atrial Fibrillation

Hanan Ahmed GalalAzzam

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/53613>

1. Introduction

Atrial fibrillation is the most common sustained cardiac arrhythmia, which is associated with a high risk of stroke and thromboembolism. Increasing evidence suggests that the thrombogenic tendency in atrial fibrillation is related to several underlying pathophysiological mechanisms. Virchow's triad, a time-honored paradigm that offers mechanistic insights for thrombus initiation and development regardless of origin, does indeed apply to atrial fibrillation thrombogenesis [1,2].

2. Mechanisms of thrombogenesis in atrial fibrillation

More than 150 years ago, Rudolf Virchow proposed a triad of events needed for thrombus formation ie, abnormal changes of the vessel wall, blood flow, and blood constituents [2]. In the 21st century, we now recognize Virchow's triad as: endothelial or endocardial damage or dysfunction (and related structural abnormal changes); abnormal blood stasis; and abnormal haemostasis, platelets, and fibrinolysis (Figure 1). Extensive abnormal changes of these variables are clearly evident in atrial fibrillation. Thus, atrial fibrillation could in fact drive a prothrombotic or hypercoagulable state, by virtue of its fulfillment of Virchow's triad for thrombogenesis [3].

Abnormal changes shown in the vessel wall (eg, atrial tissue changes, endothelial damage and dysfunction), in flow (stasis—eg, in the left atrial appendage), and in blood constituents (eg, haemoconcentration, platelets, coagulation cascade activation, inflammation); all factors contribute to propensity for thrombus formation (thrombogenesis) in atrial fibrillation. vWf=von Willebrand factor.

Figure 1. Components of Virchow's triad for thrombogenesis in atrial fibrillation [4](with permission)

2.1. Anatomical and structural considerations:

Attached to each atria is a blind-ended passage known as an appendage. The left atrial appendage (LAA) is long with a narrow inlet, thereby predisposing to blood stasis. Thus, the LAA is the most common site of intra-atrial thrombus formation, not only in atrial fibrillation, but also in patients with sinus rhythm [5,6].

Changes in the dimensions of the left atrium and LAA occur as a consequence of atrial fibrillation, with some correlation to subsequent thromboembolism. Detailed descriptions of endothelial damage in the context of atrial fibrillation are well described and can be visualised by scanning electron microscopy, especially within the appendages. Goldsmith and colleagues [7] have reported more severe endocardial changes in the LAA than in the right-atrial appendages, especially in atrial fibrillation (compared with sinus rhythm) and in mitral stenosis (compared with mitral regurgitation). Similarly, Masawa and co-workers [8] have described a "rough endocardium" with a wrinkled appearance attributable to oedema and fibrinous transformation; small areas of endothelial denudation and thrombotic aggregation have also been noted in patients with atrial fibrillation and cerebral embolism.

Extracellular matrix turnover is a dynamic structure, which continually undergoes a process of structural remodeling [9]. Structural remodelling of the atria could contribute to the hypercoagulable state, by virtue of both enhanced blood stasis and an abnormal endocardium. Structural remodeling of the left atrial appendage to include the pectinate muscles and multiple lobes of the lumen occurs in patients with permanent atrial fibrillation [10]. Morphologic studies have shown larger volumes and luminal surface areas when compared to patients without atrial fibrillation; however, both the absolute and relative surface areas of the pectinate muscles are reduced. In addition, there is significant endocardial thickening with fibrous and elastic tissue (endocardialfibroelastosis) [10].

Several studies have shown that patients with atrial fibrillation have altered amounts of collagen degradation products and impaired matrix degradation, with abnormal plasma concentrations of various matrix metalloproteinases (MMPs), their inhibitors (tissue inhibitor of MMPs [TIMPs]), and various growth factors (eg, transforming growth factor β 1) reported [11-13]. These proteins are important in the breakdown of various collagens and hence their regulation is key to ensuring healthy matrix turnover.

Evidence suggests that abnormal changes in the extracellular matrix are not related to the presence of atrial fibrillation in itself, but are probably a consequence of various coexisting comorbidities (eg, hypertension). Nevertheless, MMPs and TIMPs could have a link with the prothrombotic state, as exemplified by a correlation with prothrombin fragments 1 and 2, markers of thrombogenesis [12]. Further studies have identified disruption of other extracellular matrix components, although most have focused on these factors as a cause for the arrhythmia or explanation for remodelling and chamber dilatation [14-17]. One study suggested that some of the changes in MMPs were due to concomitant mitral valve disease [16], whereas another reported changes in the ventricular myocardium, albeit to a lesser extent [17]. Similarly, in patients with ventricular dysfunction (a potent risk factor for atrial fibrillation), various studies have also shown striking atrial structural changes [18,19].

2.2. Abnormal blood stasis

In addition to stasis consequent on the failure of atrial systole, the presence of non-valvular atrial fibrillation seems to promote progressive left atrial (LA) dilatation [20], thus amplifying the potential for stasis. In the presence of mitral stenosis, LA dilatation is increased and leads to further stasis and propensity to thrombosis [21]. The contribution of LA dilatation to thrombogenesis (at least, in non-valvular atrial fibrillation) is indicated by the finding that atrial size corrected for body surface area is an independent risk factor for stroke [22,23].

The contribution of valvular heart disease to thrombogenesis in atrial fibrillation cannot be ignored. In mitral stenosis, up to 75% of patients with cerebral emboli on computed tomography or autopsy are identified to have atrial fibrillation, presumably due to alterations in LA emptying and transmitral flow [24]. By contrast, moderate-to-severe (non-rheumatic) mitral regurgitation seems to reduce the risk of stroke with atrial fibrillation [25]. Defining patients with atrial fibrillation and mitral valve disease who are at the greatest risk of stroke has proved complex. The risk of emboli increases with age and in individuals with a lower cardiac index, but seems to correlate poorly with clinical classification or mitral valve area.

Studies assessing the degree of LA dilatation have also proven inconsistent. However, an initial embolic event is highly predictive for subsequent or recurrent thromboemboli [26].

Abnormal stasis in the LA and LAA can be visualized on TEE with spontaneous echo contrast (SEC) or pulsed-wave doppler during paroxysms of atrial fibrillation [27-30]. In sinus rhythm, aquadriphasic pattern of blood flow can be seen in the LAA, affording minimum blood stasis [31]. This pattern in blood flow is thought to be related to the intimate yet slightly delayed relations between atrial and ventricular passive and active filling. In atrial fibrillation, SEC has been shown to independently predict increased risk of thromboembolism [32].

2.3. Abnormal blood constituents

The main intravascular promoters of thrombogenesis are platelets and the various proteins of the coagulation cascade. In atrial fibrillation, abnormal changes in both these promoters and other blood constituents (eg, inflammatory cytokines, growth factors) are evident, thereby completing Virchow's triad.

2.3.1. Abnormal changes in coagulation

Abnormal haemostasis and coagulation are well described in atrial fibrillation (figure 1, table 1). In particular, increased fibrin turnover has been reported in patients with acute onset or chronic atrial fibrillation [33-39]. These changes initially seemed to be unrelated to the cause of atrial fibrillation or structural heart disease [38,39]. However, abnormal concentrations of prothrombotic indices (eg, prothrombin fragments 1 and 2 and thrombin-antithrombin complexes) are more prominent in patients with stroke who have atrial fibrillation than in those who have sinus rhythm [40], as well as in patients with atrial fibrillation and many stroke risk factors (eg, diabetes plus heart failure) compared with either risk factor alone [41-43]. Furthermore, some prothrombotic indices are abnormal in the patients with atrial fibrillation only [44,45] and in those with paroxysmal atrial fibrillation [46]. Notably, some markers have been proposed as suitable candidates to refine various stroke risk stratification schema, many of which are reasonably able to identify patients at low risk or high risk of stroke, but poor at identifying patients at moderate risk [47].

An association between various prothrombotic indices, stasis, and intracardiac thrombus has been described [48,49]. In one study, congestive cardiac failure, a history of recent embolus, and fibrin D-dimer were shown to independently predict the presence of LAA thrombi on TEE, leading the researchers to conclude that D-dimer could be useful in predicting the absence of LAA thrombi [49].

The prothrombotic state also correlates with the degree of LAA dysfunction [50,51]. Furthermore, a relation to TEE indices of stroke risk has been described. For example, SEC that is visible during TEE shows a significant correlation to prothrombin fragments 1 and 2, fibrinopeptide A, and thrombin-antithrombin III complex in non-valvular atrial fibrillation [52,53]. Patients with atrial flutter and impaired LAA function (shown by pulsed-wave doppler) have increased amounts of D-dimer and α -thromboglobulin [53]. In accordance with clinical data suggesting that mitral regurgitation protects against stroke in atrial fibrillation, a greater degree of mitral regurgitation is associated with reduced coagulation activity as estimated by fibrin D-dimer amounts [54], highlighting the important contribution of stasis.

	Study design	Comment
Gustafsson et al., [55]	20 AF with stroke; 20 AF without stroke; 20 stroke without AF; 40 healthy controls	↑D-dimer, vWF in NV AF with and without stroke
Kumagai et al., [39]	73 AF; 73 controls	↑D-dimer
Asakura et al., [56]	83 AF vs healthy controls	↑PF1+2, TATIII
Sohara&Miyahara [57]	13 paroxysmal AF vs healthy controls	NS in D-dimer, TATIII
Lip et al., [38]	87 AF; 158 controls	↑D-dimer, vWF
Lip et al., [58]	51 AF; 26 healthy controls	↑D-dimer
Kahn et al., [36]	75 NV AF with or without previous embolic events; 42 controls with or without previous thrombotic stroke	vWF higher in AF after stroke than controls without stroke and similar to controls after stroke
Heppell et al., [48]	109 AF with or without thrombus in left atrium	↑D-dimer, vWF, TATIII in patients with left atrial thrombus compared with patients without thrombus
Shinohara et al., [51]	45 NV AF	↑D-dimer, TATIII in patients with low LAA velocity vs patients with high LAA velocity
Feinberg (SPAF III) et al., [59]	1531 AF	PF1+2 not associated with thromboembolism
Mondillo et al., [44]	45 AF; 35 healthy controls	↑D-dimer, vWF, s-thrombomodulin
Fukuchi et al., [60]	AF vs without AF	↑vWF in atrial appendage tissue
Conway et al., [61]	1321 AF	↑vWF in high-risk group for stroke
Kamath et al., [62]	93 AF; 50 healthy controls	↑D-dimer
Vene et al., [63]	113 AF	↑D-dimer in AF with cardiovascular events vs no events
Nakamura et al., [64]	LAA tissue samples of 7 NV AF vs 4 without AF	↑vWF, TF expression
Conway et al., [65]	994 AF	vWF not a significant predictor of stroke and vascular events
Kamath et al., [66]	31 acute onset AF; 93 permanent AF; 31 healthy controls	Haematocrit raised in acute AF; ↑D-dimer in permanent AF, but not in acute AF
Sakurai et al., [67]	28 AFL; 27 controls	↑D-dimer in patients with impaired LAA function
Inoue et al., [42]	246 NV AF; 111 healthy controls	↑D-dimer in NV AF with risk factors, NS in PF1+2
Kumagai et al., [68]	16 AF post mortem	↑vWF mRNA and protein in AF with enlarged atriums
Marin et al., [33]	24 acute onset AF; 24 chronic AF vs 24 coronary artery disease in sinus rhythm; 24 healthy controls	↑D-dimer, vWF, s-thrombomodulin in all AF groups with no significant after cardioversion
Nozawa et al., [69]	509 AF; 111 healthy controls	↑D-dimer, NS in PF1+2
Freestone et al., [70]	59 AF; 40 healthy controls	↑Vwf
Nozawa et al., [71]	509 NV AF	↑D-dimer but not PF1+2 with predictive significance for thromboembolic events
Ohara et al., [43]	591 NV AF; 129 controls	↑D-dimer, PF1+2, platelet factor 4, and ð-thromboglobulin in NV AF; D-dimer, prothrombin fragments correlated with accumulation of clinical risk factors for stroke

AF=atrial fibrillation. NV=non-valvular. TATIII=thrombin-antithrombin III complex. AFL=atrial flutter. vWf=von Willebrand factor. LAA=left atrial appendage. TF=tissue factor. NS=non-significant. PF1+2=prothrombin fragments 1 and 2. s-thrombomodulin=soluble thrombomodulin

Table 1. Coagulation abnormal changes in atrial fibrillation [4] (with permission).

2.3.2. *Von Willebrand factor (vWf)*

Further insight into the hypercoagulable state in atrial fibrillation is provided by studies of vWf, which is a well-established index of endothelial damage and dysfunction. Raised vWf concentrations independently predict presence of LAA thrombus in atrial fibrillation [48]. Furthermore, increased LAA endocardial expression of vWf has been described [60], especially in those with an overloaded appendage, which seems to correlate with the presence of adherent platelet thrombus. Furthermore, increased expression of vWf in the endocardium has been shown to associate with enlarged LA dimensions in mitral valve disease and increased myocyte diameter [68].

Both vWf and tissue factor are overexpressed in the atrial endothelium in patients with atrial fibrillation who have a history of cardiogenic thromboembolism—specifically in the endothelial sites containing inflammatory cells and denuded endocardium, which indicate features of persistent myocarditis [64]. Plasma vWf and D-dimer are also positively correlated in patients receiving either aspirin or no antithrombotic treatment, but not in those receiving warfarin [38], further indicating the ability of warfarin to modulate the thrombogenic process.

Furthermore, a positive association between atrial fibrillation and plasma vWf was seen in the Rotterdam study [72]. This relation was most apparent in female patients, which could explain the excess risk of stroke due to atrial fibrillation in women compared with men. Furthermore, plasma vWf amounts were associated with the presence of four independent risk factors for stroke (heart failure, previous stroke, age, and diabetes) and stroke risk stratification schema [61,65]. Follow-up data from this study suggests that vWf concentrations might independently predict subsequent stroke and vascular events [65,73]. However, such applications will probably be hampered by the non-specificity of vWf, concentrations of which are also increased in various other disorders [74,75].

2.3.3. *Tissue factor*

An understanding of left atrial—left atrial appendage thrombogenesis may have its roots in distinguishing hemostatic and thrombotic clotting. Studies performed by Hoffman and Monroe [76] offer potential mechanistic insight. In a series of wounding experiments, skin punch biopsy tissue was placed on the dorsal skin of C57 black mice. Samples containing the wound specimens were then collected. For comparison, thrombus was provoked in saphenous veins by application of 10% ferric chloride. After complete occlusion, tissue blocks containing the clotted vessels were collected. Histologic evaluation revealed extensive tissue factor staining within saphenous vein thrombi. In distinct contrast, tissue factor staining in hemostatic clots was localized to squamous endothelial cells at the wounds edges—not within the thrombus itself.

The experimental findings suggest that a large volume of blood must flow over an injured surface, such as the left atrial—left atrial appendage endocardium in a person with atrial fibrillation for significant tissue factor, derived from both circulating cells and microparticles [77,78], to accumulate in high concentrations. Further, and of fundamental teleological rele-

vance, hemostasis occurs rapidly, with tissue factor of local origin determining the rate of thrombus development.

The cell-based model of coagulation translates well to left atrial-left atrial appendage thrombogenesis and supports

a primary role for tissue factor-based thrombin generation, with a secondary role being played by platelets. While the results of clinical trials [79,80], and meta-analyses are consistent with this hypothesis, several biological constructs potentially provide a mechanistic platform as well.

The integrated complexity of coagulation in general and platelet-dependent thrombin generation in particular is becoming evident. One of the most interesting and clinically relevant observations over the past decade is the concomitant interdependence and independence of platelet activation and thrombin generation. The former is best considered in the context of primary hemostasis and possibly arterial thrombosis-both highly dependent on platelet

activation, platelet aggregation and thrombin generation (in concentrations sufficient to provoke further platelet activation). In the latter instance, platelet subpopulations with distinct intracellular calcium signaling properties yield procoagulant domains [81]. The down regulation of platelet $\alpha\text{IIb}/\text{b3}$, in turn, attenuates proaggregatory potential.

2.3.4. Platelets

Many studies indicate a potential role for platelets in the hypercoagulable state (table 2). However, the results of many of these studies have been conflicting, representing the diverse aspects of platelet physiology that have been measured and possibly confounding from interlaboratory assay variability. The available data support the notion that abnormal changes of platelets in atrial fibrillation do exist, but the relation between these measures and increased thrombotic risk remains uncertain, and many of such abnormal changes could simply indicate underlying vascular comorbidities.

Choudhury and colleagues [82] recently showed that patients with atrial fibrillation had far higher amounts of platelet microparticles and soluble P-selectin than healthy controls in sinus rhythm, but no difference was seen between patients with atrial fibrillation and disease-matched controls, implying that the abnormal changes detected were a consequence of the underlying comorbidities rather than atrial fibrillation itself. Increased amounts of β -thromboglobulin, a platelet-specific protein that indicates platelet activation and is released from α -granules during platelet aggregation and subsequent thrombus formation, have been shown in patients with both valvular and non-valvular atrial fibrillation compared with controls in sinus rhythm [51,58,62,83-86]. Substantially higher β -thromboglobulin amounts have been measured in patients with the lowest LAA flow velocities, who had greater left-atrial dimensions [51], suggesting that platelet activation could be enhanced in patients with a greater degree of intra-atrial stasis.

Despite the presence of enhanced platelet activation in atrial fibrillation, any firm clinical evidence indicating that it directly enhances thrombotic risk is lacking. A substudy from the

Stroke Prevention in Atrial Fibrillation III (SPAF-III) trial [59] recorded no association between plasma β -thromboglobulin amounts and subsequent thromboembolic events. By contrast, the population-based Rotterdam study [87] showed that plasma concentrations of soluble P-selectin were predictive of adverse clinical outcomes in elderly patients with atrial fibrillation.

	Patient group	Results
Yamauchi et al., [87]	26 V AF; 73 NV AF; 57 healthy controls	↑ BTG in AF, NS in PF4
Furui et al., [95]	20 AF; 15 healthy controls	BTG increase greater in AF than in controls after treadmill exercise
Gustafsson et al., [55]	20 AF with stroke; 20 AF without stroke; 20 stroke with sinus rhythm; 40 healthy controls	↑BTG, PF4 in NV AF with and without stroke
Sohara et al., [57]	13 paroxysmal AF vs healthy controls	NS in BTG, PF4
Lip et al., [58]	51 AF; 26 healthy controls	↑BTG
Heppell [48]	109 AF with or without thrombus in left atrium	↑BTG, PF4 in patients with left atrial thrombus compared with patients without thrombus
Minamino et al., [86]	25 AF vs healthy controls	↑mP-sel
Shinohara et al., [51]	45 NV AF	↑BTG, PF4 in patients with low LAA velocity vs patients high LAA velocity
Feinberg et al., SPAF III) [59]	1531 AF	BTG not associated with thromboembolism
Minamino et al., [93]	28 AF	↑BTG, mP-sel
Mondillo et al., [44]	45 AF; 35 healthy controls	↑BTG, PF4
Kamath et al., [45]	93 AF; 50 healthy controls	↑BTG, soluble GPV; NS in platelet aggregation
Conway et al., [61]	1321 AF	sP-sel was independent of risk for stroke
Conway et al., [65]	994 AF	sP-sel not a significant predictor of stroke and vascular events
Atalar et al., [96]	15 paroxysmal AF; 25 chronic AF; 22 healthy controls	Higher BTG, PF4 in chronic AF than in paroxysmal AF and after conversion to sinus rhythm
Nozawa et al., [69]	509 AF; 111 healthy controls	↑BTG, PF4
Sakurai et al., [53]	28 AFL; 27 controls	↑BTG in patients with impaired LAA function
Inoue et al., [42]	246 NV AF; 111 controls	↑BTG, NS in PF4
Nozawa et al., [69]	509 NV AF	BTG, PF4 did not predict thromboembolic events
Choudhury et al., [97]	121 NV AF; 65 healthy controls; 78 disease-matched controls	Similar but raised levels of CD62P, CD63 and sP-sel in AF and disease-matched controls compared with healthy controls
Choudhury et al., [82]	70 NV AF; 46 disease controls; 33 healthy controls	↑PMP and sP-sel in AF and disease controls, but AF not an independent determinant

Atrial fibrillation. V=valvular. NV=non-valvular. AFL=atrial flutter. BTG= β -thromboglobulin. PF4=platelet factor 4. sPsel=soluble P-selectin. mP-sel=matrix P-selectin. GPV=glycoprotein V. LAA=left atrial appendage. NS=non-significant. PMP=platelet microparticles.

Table 2. Studies of platelet function in atrial fibrillation [4] (with permission)

However, Nagao et al., [88] suggests that thromboembolism in atrial fibrillation is probably due to enhancement of various components of the coagulation system due to stasis of blood in the inordinate and irregular atria, rather than to platelet activation *per se*. The relative role of coagulation versus platelet activation in the pathogenesis of thrombogenesis in patients with atrial fibrillation can roughly be inferred from the results of antithrombotic drug interventions that have been tested in randomized clinical trials [89-91]. These results indicate that inhibition of coagulation remains the mainstay in preventing atrial fibrillation –related thrombogenesis. The lesser but significant role of platelets- best inhibited by a combined antiplatelet drug regimen – is presumably related to the platelet activation seen in this arrhythmia could contribute to thrombogenesis indirectly. For example, increased expression of P-selectin on platelets associated with reduced concentrations of nitric oxide has also been shown to be a risk factor for silent cerebral infarction in patients with atrial fibrillation [92]. Moreover, raised amounts of P-selectin and CD63 have both been associated with the embolic and pre-embolic status of patients with non-rheumatic atrial fibrillation [93]. Or the prominent involvement of platelets in the pathogenesis of atherothrombotic (that is, non-cardiembolic) events [94].

2.3.5. *Abnormal changes in fibrinolysis*

Few studies have focused on fibrinolytic function in atrial fibrillation. Enhanced fibrinolysis, shown by increased concentrations of tissue-plasminogen activator (t-PA) antigen and t-PA inhibitor (PAI)-1 and reduced amounts of plasmin-antiplasmin complex can be attributable to a pathophysiological response to the prothrombotic state [95,66]. However, the available data are not consistent and conflicting results have also been reported [35]. In the Stroke Prevention in Atrial Fibrillation (SPAF) III study [99], increased concentrations of plasmin-antiplasmin complexes were independently associated with thromboembolic risk factors such as older age (>75 years), recent congestive heart failure, decreased fractional shortening, and recent onset of atrial fibrillation. A significant correlation can be also shown between t-PA amounts and left-atrial diameter in atrial fibrillation [35]. Predictably, anticoagulation leads to some improvement in fibrinolytic markers in rheumatic atrial fibrillation [98].

Increased amounts of t-PA and PAI-1 can indicate the coexistence of confounders, such as hypertension, heart failure, or ischaemic heart disease, all of which can cause endothelial dysfunction, damage, and inflammation. However, studies in patients with atrial fibrillation only confirm that presence of the disorder does modulate these markers. [35,95,99]. Thus, the high amounts of t-PA and PAI-1 in atrial fibrillation could be a consequence of endothelial damage and dysfunction or represent systemic inflammation [100,101]. PAI-1 concentrations are also predictive of successful cardioversion [102], and are independent predictors of the development of atrial fibrillation after cardiopulmonary bypass [103].

It is unclear whether increased amounts of t-PA or PAI-1 in atrial fibrillation are due to endothelial dysfunction, inflammation, fibrinolysis, or vascular disease, or a combination. Nevertheless, abnormal changes in the fibrinolytic system might relate not only to thrombogenesis but also to structural remodelling of the atria, in view of the strong links to extracellular matrix turnover.

2.3.6. Restoration of sinus rhythm

Some evidence suggests that activation of the coagulation system could be adversely affected by cardioversion of atrial fibrillation [104]. Electrical cardioversion has been associated with more prominent activation of the coagulation system than a pharmacological strategy [105]. One study found a positive correlation between the energy delivered for cardioversion to sinus rhythm and plasma D-dimer values on day 7 [105]. Additionally, an extended duration of atrial fibrillation could lead to a more prominent hypercoagulable state (estimated by D-dimer value) after cardioversion[106]. The hypercoagulable state after cardioversion has been seen despite optimum anticoagulation with warfarin [107]. Nevertheless, patients receiving therapeutic low-molecular-weight heparin (LMWH) before cardioversion seem to have reduced hypercoagulability [108].

2.4. What drives the prothrombotic state in atrial fibrillation?

Several mechanisms have been purported to drive then prothrombotic state in atrial fibrillation (figure 2), but recent evidence has focused on the potential role of inflammation and the release of various growth factors.

Figure 2. Abnormal changes in coagulation during atrial fibrillation [4] (with permission)

2.4.1. *Inflammation*

In atrial fibrillation, inflammation might not only result in endothelial damage, dysfunction, or activation, but also be linked directly to thrombogenesis. Increasing evidence has supported a link between inflammation and the initiation and perpetuation of atrial fibrillation [109-113]. Furthermore, abnormal changes in systemic inflammation have been related to prothrombotic indices in atrial fibrillation, suggesting that inflammation could drive the prothrombotic state in atrial fibrillation [109].

Although most cases of atrial fibrillation are associated with various comorbidities, many of which could also enhance the baseline inflammatory state, there may be an underlying direct link between atrial fibrillation and inflammation. Interleukin-6 concentrations are abnormal in atrial fibrillation, with some prognostic implications shown in one study [114]. Many studies have also shown that amounts of high-sensitivity C-reactive protein (hs-CRP) are greater in patients with atrial fibrillation than in controls in sinus rhythm, with a step-wise increase in hs-CRP with the transition from patient groups with an increasing burden (sinus rhythm to paroxysmal then persistent) in atrial fibrillation [115]. Raised hs-CRP amounts consistently correlate with cardiovascular risk, although not with future atrial fibrillation [109]. More recently, high hs-CRP amounts were shown to be predictive of mortality and vascular death in atrial fibrillation, but not stroke itself [116].

How is inflammation linked to thrombogenesis in atrial fibrillation? Both CRP and interleukin 6 stimulate tissue factor production from monocytes in vitro [117, 118]. Furthermore, interleukin 6 increases platelet production and sensitivity to thrombin [119], stimulates transcription of fibrinogen [120], and is linked to both endothelial activation and damage [121,122]. However, no link seems to exist between hs-CRP and thrombin-antithrombin complexes [123]. Tissue factor and high stroke risk are also independent associates of interleukin 6, whereas fibrinogen and plasma viscosity are independent associates of hs-CRP amounts [124].

2.4.2. *Growth factors*

Another potential driver for thrombogenesis could be growth factors. Various pro-angiogenic factors have been identified; concentrations of some of these factors have been shown to alter in atrial fibrillation [70,125,126]. Vascular endothelial growth factor (VEGF) is largely produced by activated platelets [127], and results in upregulation of tissue factor mRNA production and subsequent expression of this compound on the endothelial membrane [128]. VEGF amounts are substantially increased in both persistent and permanent atrial fibrillation, with a corresponding increase in tissue factor [125]. Additionally, raised serum concentrations of transforming growth factor- β 1 [126] and angiopoietin 2 (but not angiopoietin 1) [70] are also recorded in atrial fibrillation, showing the depth and complexity of modulation of growth factor amounts.

Although the requirements for enhanced angiogenesis in atrial fibrillation are unknown, in view of the intimate association between VEGF and tissue factor, enhanced growth factors could be a crucial driving force behind the hypercoagulable state. Notably, tissue factor acts

as a cofactor to factor VIIa and is widely regarded as the physiological trigger to thrombin formation [129]. But, why are factors such as the angiopoietins involved? Angiopoietin 1 and 2 are natural co-antagonists and both compete for the same binding site on Tie-2, an endothelial tyrosine kinase receptor. With an excess of angiopoietin 1, stability of the endothelium is favoured, whereas the converse is true with an excess of angiopoietin 2 [70]. In these circumstances, the balance could ultimately favour endothelial destabilisation and therefore the action of cytokines such as VEGF.

2.4.3. Extracellular matrix turnover

The relationship between atrial fibrillation and remodeling of the left atrium/arterial appendage is traditionally explained by the absence of contractility and altered flow dynamics. This hypothesis is not entirely fulfilling for several reasons, not the least of which is its inability to substantiate the mechanism of progressive structural change. A contemporary view considers the contribution of coagulation factors and thrombus substrate itself as both initiators and perpetuators of the prothrombotic environment that includes remodeling. Thrombin, a serine protease, beyond its widely recognized role in hemostasis and thrombosis, is directly involved in tissue repair and remodeling through an endothelial mesenchymal transdifferentiation process [130]. Thrombin also exerts an effect on endothelial cell junctions (reviewed in [131], endothelial cell and smooth muscle cell migration and smooth muscle cell proliferation via protease activated receptor (PAR)-1 [132].

Thrombin-induced membrane-type matrix metalloproteinase (MMP)-2 gene transcription and activity [133] may also contribute to structural changes in the atrium/atrial appendage, as may thrombin-augmented fibroblast-mediated collagen gel contraction [134]. Locally generated thrombin has been shown in tissue culture to upregulate tissue factor expression and activity [135].

While thrombin is known to possess a variety of cell regulating capabilities, one must not overlook the contribution

of other coagulation proteases in the remodeling process. Indeed, factor Xa has been shown to promote fibroblast proliferation, migration and differentiation into myofibroblasts through a PAR-2 specific mechanism [136]. Accordingly, the development of oral/direct factor Xa and thrombin inhibitors provides an unprecedented opportunity to investigate fundamental pathological mechanisms in atrial fibrillation.

2.4.4. Nitric oxide

Nitric oxide is synthesized by nitric oxide synthase, which is present in large concentrations in the endothelium. The expression of nitric oxide synthase is regulated by flow-mediated shear stress and is consequently downregulated at sites with low flow velocity [137]. Nitric oxide shows potent antithrombotic effects in arterial endothelium, and nitric oxide released from activated platelets inhibits platelet recruitment to the growing thrombus [138], while also inhibiting expression of PAI-1 [139].

In animal models of atrial fibrillation, the loss of atrial contraction and consequent reduction in shear stress seems to reduce LA expression of nitric oxide synthase with a corresponding decrease in nitric oxide bioavailability and increase in PAI-1 expression [140]. In the LAA, nitric oxide concentrations were also significantly reduced compared with control animals, but this finding did not indicate decreased expression of nitric oxide synthase at this site. Since atrial thrombus is frequently formed in the LAA, this finding still has no adequate explanation.

2.4.5. Renin-Angiotensin-Aldosterone System (RAAS)

The RAAS is now appreciated as key to the pathophysiology of various cardiovascular disease states. The extent of these changes seems to relate predominantly to the reduction in angiotensin-II amounts. Atrial tissue has the capacity to produce and use this hormone with local expression of acetylcholinesterase and angiotensin-II receptors, both of which could be upregulated in atrial fibrillation [141]. RAAS could be mechanistically implicated in initiation and perpetuation of atrial fibrillation [141-143], as well as providing the link to other mechanisms promoting the prothrombotic state in atrial fibrillation.

Angiotensin II has been shown to possess several proinflammatory properties and increases the production of proinflammatory cytokines (eg, interleukin 6 and tumour necrosis factor α [TNF α]), adhesion molecules (eg, vascular-cell adhesion molecule 1), monocyte chemoattractant protein 1, and selectins (eg, P-selectin) [144-146]. Similarly, through release of various chemokines (eg, cytokine-induced neutrophil chemoattractant), angiotensin II can initiate neutrophil recruitment [146]. Expression of angiotensin-II receptors has also been linked with increased atrial cell death and leucocyte infiltration [147]. These data potentially support a complex relation between RAAS, inflammation, and atrial fibrillation.

Additionally, RAAS has been implicated in the activation of various MMPs and thromboxane A2 (a prothrombotic signalling molecule produced by activated platelets). These processes could occur both as a direct effect of angiotensin II and also through induction of interleukin 6 [148]. Furthermore, angiotensin II could accelerate degradation of nitric oxide through production of reactive oxygen species and thereby impair endothelium dependent vasodilatation [149]. Likewise, activation of RAAS increases synthesis of PAI-1, possibly indicating either enhanced endothelial damage or impaired fibrinolysis in atrial fibrillation [150].

3. Conclusion

The mechanisms underlying thrombogenesis in atrial fibrillation are clearly complex and remain only partly understood. Abnormal changes in flow, vessel wall, and blood constituents in atrial fibrillation fulfil Virchow's triad for thrombogenesis, and accord with a prothrombotic or hypercoagulable state in this arrhythmia. That this process is related purely to blood stasis is no longer accepted. Various abnormal changes related both to atrial fibrillation and its comorbidities impart a synergistic effect in maintaining a hypercoagulable state in this condition.

Author details

Hanan Ahmed GalalAzzam

Mansoura university/ Faculty of Medicine, Egypt

References

- [1] Lip GYH. Does atrial fi brillation confer a hypercoagulable state? *Lancet* 1995; 346: 1313–14.
- [2] Brotman DJ, Deitcher SR, Lip GY, Martzdorff AC. Virchow's triad revisited. *South Med J* 2004; 97: 213–14.
- [3] Choudhury A, Lip GY. Atrial fibrillation and the hypercoagulable state: from basic science to clinical practice. *PathophysiolHaemostThromb* 2003–2004; 33: 282–89.
- [4] Watson T, Shantsila E, Lip YHG. Mechanisms of thrombogenesis in atrial fibrillation: Virchow's triad revisited. *Lancet* 2009;373(9658):155-66.
- [5] Blackshear JL, Odell JA. Appendage obliteration to reduce stroke incardiac surgical patients with atrial fi brillation.*Ann ThoracSurg*1996; 61: 755–59.
- [6] Pollick C, Taylor D. Assessment of left atrial appendage function bytransesophageal echocardiography. Implications for thedevelopment of thrombus.*Circulation* 1991; 84: 223–31.
- [7] Goldsmith IR, Blann AD, Patel RL, Lip GY. Von Willebrand factor,fibrinogen, and soluble P-selectin levels after mitral valve replacementversus mitral valve repair. *Am J Cardiol* 2000; 85: 1218–22.
- [8] Masawa N, Yoshida Y, Yamada T, Joshita T, Ooneda G. Diagnosis ofcardiac thrombosis in patients with atrial fi brillation in the absenceof macroscopically visible thrombi. *Virchows Arch A PatholAnatHistopathol* 1993; 422: 67-71..
- [9] Dollery CM, McEwan JR, Henney AM. Matrix metalloproteinasesand cardiovascular disease.*Circ Res* 1995; 77: 863–68.
- [10] Shirani J, Alaeddini J. Structural remodeling of the left atrial appendage in patients with chronic non-valvular atrial fibrillation: implications for thrombus formation, systemic embolism, and assessment by transesophageal echocardiography. *Cardio-vascPathol* 2000; 9(2):95-101.
- [11] Tziakas DN, Chalikias GK, Papanas N, Stakos DA, Chatzikyriakou SV, Maltezos E. Circulating levels of collagen type I degradation marker depend on the type of atrial fi brillation. *Europace* 2007; 9: 589–96.

- [12] Marin F, Roldan V, Climent V, Garcia A, Marco P, Lip GY. Is thrombogenesis in atrial fibrillation related to matrix metalloproteinase-1 and its inhibitor, TIMP-1? *Stroke* 2003; 34: 1181–86.
- [13] Li X, Ma C, Dong J, et al. The fibrosis and atrial fibrillation: is the transforming growth factor-beta(1) a candidate etiology of atrial fibrillation. *Med Hypotheses* 2008; 70: 317–19.
- [14] Xu J, Cui G, Esmailian F, et al. Atrial extracellular matrix remodeling and the maintenance of atrial fibrillation. *Circulation* 2004; 109: 363–68.
- [15] Nakano Y, Niida S, Dote K, et al. Matrix metalloproteinase-9 contributes to human atrial remodeling during atrial fibrillation. *J Am Coll Cardiol* 2004; 43: 818–25.
- [16] Anne W, Willems R, Roskams T, et al. Matrix metalloproteinases and atrial remodeling in patients with mitral valve disease and atrial fibrillation. *Cardiovasc Res* 2005; 67: 655–66.
- [17] Mukherjee R, Herron AR, Lowry AS, et al. Selective induction of matrix metalloproteinases and tissue inhibitor of metalloproteinases in atrial and ventricular myocardium in patients with atrial fibrillation. *Am J Cardiol* 2006; 97: 532–37.
- [18] Sanders P, Morton JB, Davidson NC, et al. Electrical remodeling of the atria in congestive heart failure: electrophysiological and electroanatomic mapping in humans. *Circulation* 2003; 108: 1461–68.
- [19] Li D, Fareh S, Leung TK, Nattel S. Promotion of atrial fibrillation by heart failure in dogs: atrial remodeling of a different sort. *Circulation* 1999; 100: 87–95.
- [20] Sanfilippo AJ, Abascal VM, Sheehan M, et al. Atrial enlargement as a consequence of atrial fibrillation. A prospective echocardiographic study. *Circulation* 1990; 82: 792–97.
- [21] Keren G, Etzion T, Sherez J, et al. Atrial fibrillation and atrial enlargement in patients with mitral stenosis. *Am Heart J* 1987; 114: 1146–55.
- [22] The Stroke Prevention in Atrial Fibrillation Investigators. Predictors of thromboembolism in atrial fibrillation: II Echocardiographic features of patients at risk. *Ann Intern Med* 1992; 116: 6–12.
- [23] Di Tullio MR, Sacco RL, Sciacca RR, Homma S. Left atrial size and the risk of ischemic stroke in an ethnically mixed population. *Stroke* 1999; 30: 2019–24.
- [24] Salem DN, Stein PD, Al Ahmad A, et al. Antithrombotic therapy in valvular heart disease—native and prosthetic: the seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126: 457–82S.
- [25] Nakagami H, Yamamoto K, Ikeda U, Mitsuhashi T, Goto T, Shimada K. Mitral regurgitation reduces the risk of stroke in patients with nonrheumatic atrial fibrillation. *Am Heart J* 1998; 136: 528–32.

- [26] Chiang CW, Lo SK, Ko YS, Cheng NJ, Lin PJ, Chang CH. Predictors of systemic embolism in patients with mitral stenosis. A prospective study. *Ann Intern Med* 1998; 128: 885–89.
- [27] Pollick C, Taylor D. Assessment of left atrial appendage function by transesophageal echocardiography. Implications for the development of thrombus. *Circulation* 1991; 84: 223–31.
- [28] Asinger RW, Koehler J, Pearce LA, et al. Pathophysiologic correlates of thromboembolism in nonvalvular atrial fibrillation: II Dense spontaneous echocardiographic contrast (The Stroke Prevention in Atrial Fibrillation [SPAF-III] study). *J Am Soc Echocardiogr* 1999; 12: 1088–96.
- [29] Handke M, Harloff A, Hetzel A, Olschewski M, Bode C, Geibel A. Left atrial appendage flow velocity as a quantitative surrogate parameter for thromboembolic risk: determinants and relationship to spontaneous echocontrast and thrombus formation—a transesophageal echocardiographic study in 500 patients with cerebral ischemia. *J Am Soc Echocardiogr* 2005; 18: 1366–72.
- [30] Obarski TP, Salcedo EE, Castle LW, Stewart WJ. Spontaneous echo contrast in the left atrium during paroxysmal atrial fibrillation. *Am Heart J* 1990; 120: 988–90.
- [31] Jue J, Winslow T, Fazio G, Redberg RF, Foster E, Schiller NB. Pulsed doppler characterization of left atrial appendage flow. *J Am Soc Echocardiogr* 1993; 6: 237–44.
- [32] The Stroke Prevention In Atrial Fibrillation Investigators committee on echocardiography. Transesophageal echocardiographic correlates of thromboembolism in high-risk patients with nonvalvular atrial fibrillation. *Ann Intern Med* 1998; 128: 639–47.
- [33] Marin F, Roldan V, Climent VE, et al. Plasma von Willebrand factor, soluble thrombomodulin, and fibrin D-dimer concentrations in acute onset non-rheumatic atrial fibrillation. *Heart* 2004; 90: 1162–66.
- [34] Mahe I, Drouet L, Chassany O, et al. D-dimer: a characteristic of the coagulation state of each patient with chronic atrial fibrillation. *Thromb Res* 2002; 107: 1–6.
- [35] Roldan V, Marin F, Marco P, Martinez JG, Calatayud R, Sogorb F. Hypofibrinolysis in atrial fibrillation. *Am Heart J* 1998; 136: 956–60.
- [36] Kahn SR, Solymoss S, Flegel KM. Nonvalvular atrial fibrillation: evidence for a prothrombotic state. *CMAJ* 1997; 157: 673–81.
- [37] Mitusch R, Siemens HJ, Garbe M, Wagner T, Sheikhzadeh A, Diederich KW. Detection of a hypercoagulable state in nonvalvular atrial fibrillation and the effect of anticoagulant therapy. *Thromb Haemost* 1996; 75: 219–23.
- [38] Lip GY, Lowe GD, Rumley A, Dunn FG. Increased markers of thrombogenesis in chronic atrial fibrillation: effects of warfarin treatment. *Br Heart J* 1995; 73: 527–33.

- [39] Kumagai K, Fukunami M, Ohmori M, Kitabatake A, Kamada T, Hoki N. Increased intracardiovascular clotting in patients with chronic atrial fibrillation. *J Am CollCardiol* 1990; 16: 377–80.
- [40] Turgut N, Akdemir O, Turgut B, et al. Hypercoagulopathy in stroke patients with nonvalvular atrial fibrillation: hematologic and cardiologic investigations. *ClinAppl-ThrombHemost* 2006; 12: 15–20.
- [41] Varughese GI, Patel JV, Tomson J, Lip GY. The prothrombotic risk of diabetes mellitus in atrial fibrillation and heart failure. *J ThrombHaemost* 2005; 3: 2811–13.
- [42] Inoue H, Nozawa T, Okumura K, Jong-Dae L, Shimizu A, Yano K. Prothrombotic activity is increased in patients with nonvalvular atrial fibrillation and risk factors for embolism. *Chest* 2004; 126: 687–92.
- [43] Ohara K, Inoue H, Nozawa T, et al. Accumulation of risk factors enhances the prothrombotic state in atrial fibrillation. *Int J Cardiol* 2008; 126: 316–21.
- [44] Mondillo S, Sabatini L, Agricola E, et al. Correlation between left atrial size, prothrombotic state and markers of endothelial dysfunction in patients with lone chronic non-rheumatic atrial fibrillation. *Int J Cardiol* 2000; 75: 227–32.
- [45] Kamath S, Blann AD, Chin BS, Lip GY. A prospective randomized trial of aspirin-clopidogrel combination therapy and dose-adjusted warfarin on indices of thrombogenesis and platelet activation in atrial fibrillation. *J Am CollCardiol* 2002; 40: 484–90.
- [46] Li-Saw-Hee FL, Blann AD, Gurney D, Lip GY. Plasma von Willebrand factor, fibrinogen and soluble P-selectin levels in paroxysmal, persistent and permanent atrial fibrillation. Effects of cardioversion and return of left atrial function. *Eur Heart J* 2001; 22: 1741–47.
- [47] Roldan V, Marin F, Garcia-Herola A, Lip GY. Correlation of plasma von Willebrand factor levels, an index of endothelial damage/ dysfunction, with two point-based stroke risk stratification scores in atrial fibrillation. *Thromb Res* 2005; 116: 321–25.
- [48] Heppell RM, Berkin KE, McLenachan JM, Davies JA. Haemostatic and haemodynamic abnormalities associated with left atrial thrombosis in non-rheumatic atrial fibrillation. *Heart* 1997; 77: 407–11.
- [49] Habara S, Dote K, Kato M, et al. Prediction of left atrial appendage thrombi in nonvalvular atrial fibrillation. *Eur Heart J* 2007; 28: 2217–22.
- [50] Igarashi Y, Kashimura K, Makiyama Y, Sato T, Ojima K, Aizawa Y. Left atrial appendage dysfunction in chronic nonvalvular atrial fibrillation is significantly associated with an elevated level of brain natriuretic peptide and a prothrombotic state. *JpnCirc J* 2001; 65: 788–92.
- [51] Shinohara H, Fukuda N, Soeki T, et al. Relationship between flow dynamics in the left atrium and hemostatic abnormalities in patients with nonvalvular atrial fibrillation. *Jpn Heart J* 1998; 39: 721–30.

- [52] Tsai LM, Chen JH, Tsao CJ. Relation of left atrial spontaneous echo contrast with pre-thrombotic state in atrial fibrillation associated with systemic hypertension, idiopathic dilated cardiomyopathy, or no identifiable cause (lone). *Am J Cardiol* 1998; 81: 1249–52.
- [53] Sakurai K, Hirai T, Nakagawa K, et al. Left atrial appendage function and abnormal hypercoagulability in patients with atrial fibrillation. *Chest* 2003; 124: 1670–74.
- [54] Lip GY, Rumley A, Dunn FG, Lowe GD. Thrombogenesis in mitral regurgitation and aortic stenosis. *Angiology* 1996; 47: 1117–25.
- [55] Gustafsson C, Blomback M, Britton M, Hamsten A, Svensson J. Coagulation factors and the increased risk of stroke in nonvalvular atrial fibrillation. *Stroke* 1990; 21: 47–51.
- [56] Asakura H, Hifumi S, Jokaji H, et al. Prothrombin fragment F1 + 2 and thrombin-antithrombin III complex are useful markers of the hypercoagulable state in atrial fibrillation. *Blood Coagul Fibrinolysis* 1992; 3: 469–73.
- [57] Sohara H, Miyahara K. Effect of atrial fibrillation on the fibrino-coagulation system-study in patients with paroxysmal atrial fibrillation. *Jpn Circ J* 1994; 58: 821–26.
- [58] Lip GY, Lip PL, Zarifis J, et al. Fibrin D-dimer and beta-thromboglobulin as markers of thrombogenesis and platelet activation in atrial fibrillation. Effects of introducing ultra-low-dose warfarin and aspirin. *Circulation* 1996; 94: 425–31.
- [59] Feinberg WM, Pearce LA, Hart RG, et al. Markers of thrombin and platelet activity in patients with atrial fibrillation: correlation with stroke among 1531 participants in the stroke prevention in atrial fibrillation III study. *Stroke* 1999; 30: 2547–53.
- [60] Fukuchi M, Watanabe J, Kumagai K, et al. Increased von Willebrand factor in the endocardium as a local predisposing factor for thrombogenesis in overloaded human atrial appendage. *J Am Coll Cardiol* 2001; 37: 1436–42.
- [61] Conway DS, Pearce LA, Chin BS, Hart RG, Lip GY. Plasma von Willebrand factor and soluble P-selectin as indices of endothelial damage and platelet activation in 1321 patients with nonvalvular atrial fibrillation: relationship to stroke risk factors. *Circulation* 2002; 106: 1962–67.
- [62] Kamath S, Blann AD, Chin BS, et al. A study of platelet activation in atrial fibrillation and the effects of antithrombotic therapy. *Eur Heart J* 2002; 23: 1788–95.
- [63] Vene N, Mavri A, Kosmelj K, Stegnar M. High D-dimer levels predict cardiovascular events in patients with chronic atrial fibrillation during oral anticoagulant therapy. *Thromb Haemost* 2003; 90: 1163–72.
- [64] Nakamura Y, Nakamura K, Fukushima-Kusano K, et al. Tissue factor expression in atrial endothelia associated with nonvalvular atrial fibrillation: possible involvement in intracardiac thrombogenesis. *Thromb Res* 2003; 111: 137–42.

- [65] Conway DS, Pearce LA, Chin BS, Hart RG, Lip GY. Prognostic value of plasma von Willebrand factor and soluble P-selectin as indices of endothelial damage and platelet activation in 994 patients with nonvalvular atrial fibrillation. *Circulation* 2003; 107: 3141–45.
- [66] Kamath S, Blann AD, Chin BS, Lip GY. Platelet activation, haemorheology and thrombogenesis in acute atrial fibrillation: a comparison with permanent atrial fibrillation. *Heart* 2003; 89: 1093–95.
- [67] Sakurai K, Hirai T, Nakagawa K, et al. Prolonged activation of hemostatic markers following conversion of atrial flutter to sinus rhythm. *Circ J* 2004; 68: 1041–44
- [68] Kumagai K, Fukuchi M, Ohta J, et al. Expression of the vonWillebrand factor in atrial endocardium is increased in atrial fibrillation depending on the extent of structural remodeling. *Circ J* 2004; 68: 321–27.
- [69] Nozawa T, Inoue H, Iwasa A, et al. Effects of anticoagulation intensity on hemostatic markers in patients with non-valvular atrial fibrillation. *Circ J* 2004; 68: 29–34.
- [70] Freestone B, Chong AY, Lim HS, Blann A, Lip GYH. Angiogenic factors in atrial fibrillation: a possible role in thrombogenesis? *Ann Med* 2005; 37: 365–72.
- [71] Nozawa T, Inoue H, Hirai T, et al. D-dimer level influences thromboembolic events in patients with atrial fibrillation. *Int J Cardiol* 2006; 109: 59–65.
- [72] Conway DS, Heeringa J, Van Der Kuip DA, et al. Atrial fibrillation and the prothrombotic state in the elderly: the Rotterdam Study. *Stroke* 2003; 34: 413–17.
- [73] Lip GY, Lane D, Van Walraven C, Hart RG. Additive role of plasma von Willebrand factor levels to clinical factors for risk stratification of patients with atrial fibrillation. *Stroke* 2006; 37: 2294–300.
- [74] Varughese GI, Patel JV, Tomson J, Lip GY. Effects of blood pressure on the prothrombotic risk in 1235 patients with non-valvular atrial fibrillation. *Heart* 2007; 93: 495–99.
- [75] Lip GY, Pearce LA, Chin BS, Conway DS, Hart RG. Effects of congestive heart failure on plasma von Willebrand factor and soluble P-selectin concentrations in patients with non-valvular atrial fibrillation. *Heart* 2005; 91: 759–63.
- [76] Hoffman M, Whinna HC, Monroe DM. Circulating tissue factor accumulates in thrombi, but not in hemostatic plugs. *J ThrombHaemost* 2006; 4:2092–2093.
- [77] Lechner D, Weltermann A. Circulating tissue factor-exposing microparticles. *Thromb Res* 2008; 122:S47–S54.
- [78] George FD. Microparticles in vascular diseases. *Thromb Res* 2008; 122:S55–S59.
- [79] Connolly S, Pogue J, Hart R et al., Clopidogrel plus aspirin versus oral anticoagulation for atrial fibrillation in the Atrial fibrillation Clopidogrel Trial with Irbesartan

- for prevention of Vascular Events (ACTIVE W): a randomised controlled trial. *Lancet* 2006; 367:1903–1912.
- [80] Healey JS, Hart RG, Pogue J et al., Risks and benefits of oral anticoagulation compared with clopidogrel plus aspirin in patients with atrial fibrillation according to stroke risk: the atrial fibrillation clopidogrel trial with irbesartan for prevention of vascular events (ACTIVE-W). *Stroke* 2008; 39:1482–1486.
 - [81] Munnix IC, Kuijpers MJ, Auger J et al., Segregation of platelet aggregatory and pro-coagulant microdomains in thrombus formation: regulation by transient integrin activation. *ArteriosclerThrombVascBiol* 2007; 27:2484–2490.
 - [82] Choudhury A, Chung I, Blann AD, Lip GY. Elevated platelet microparticle levels in nonvalvular atrial fibrillation: relationship to P-selectin and antithrombotic therapy. *Chest* 2007; 131: 809–15
 - [83] Kaplan KL, Nossel HL, Drillings M, Lesznik G. Radioimmunoassay of platelet factor 4 and beta-thromboglobulin: development and application to studies of platelet release in relation to fibrinopeptide A generation. *Br J Haematol* 1978; 39: 129–46.
 - [84] Kunishima S, Hattori M, Kobayashi S, et al. Activation and destruction of platelets in patients with rheumatic heart disease. *Eur Heart J* 1994; 15: 335–38.
 - [85] Minamino T, Kitakaze M, Asanuma H, et al. Plasma adenosine levels and platelet activation in patients with atrial fibrillation. *Am J Cardiol* 1999; 83: 194–98.
 - [86] Yamauchi K, Furui H, Taniguchi N, Sotobata I. Plasma beta-thromboglobulin and platelet factor 4 concentrations in patients with atrial fibrillation. *Jpn Heart J* 1986; 27: 481–87.
 - [87] Heeringa J, Conway DS, van der Kuip DA, et al. A longitudinal population-based study of prothrombotic factors in elderly subjects with atrial fibrillation: the Rotterdam Study 1990–1999. *J ThrombHaemost* 2006; 4: 1944–49.
 - [88] Nagao, T.; Hamamoto, M.; Kanda, A.; Tsuganesawa, T.; Ueda, M.; Kobayashi, K.; Miyazaki, T. & Terashi, A. (1995). Platelet activation is not involved in acceleration of the coagulation system in acute cardioembolic stroke with nonvalvular atrial fibrillation. *Stroke*, Vol.26:pp.1365-8.
 - [89] Hart, RG.; Pearce, LA. & Aguilar, MI. Meta-analysis: antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann internal Med* 2007; 146:857-67.
 - [90] Lip, GY. & Lim, BS. Atrial fibrillation and stroke prevention. *Lancet Neurol* 2007; 6: 981-93.
 - [91] Connolly, SJ., Pogue, J., Hart, RG., Hart, RG., Hohnloser, SH., Pfeffer, M., Chrolavicius, S. & Yusuf, S. Effect of clopidogrel added to aspirin in patients with atrial fibrillation. *ACTIVE Investigators, N Engl J Med* 2009; 360: 2066-78.

- [92] Minamino T, Kitakaze M, Sanada S, et al. Increased expression of P-selectin on platelets is a risk factor for silent cerebral infarction in patients with atrial fibrillation: role of nitric oxide. *Circulation* 1998;98: 1721–27.
- [93] Pongratz G, Brandt-Pohlmann M, Henneke KH, et al. Platelet activation in embolic and preembolic status of patients with nonrheumatic atrial fibrillation. *Chest* 1997; 111: 929–33.
- [94] Lip, GYH.; Huber, K.; Andreotti, F.; Arnesen, H.; Airaksinen KJ; Cuisset, T.; Kirchhof, P. & Marin, F. Management of antithrombotic therapy in atrial fibrillation patients presenting with acute coronary syndrome and/or undergoing percutaneous coronary intervention/stenting. *ThrombHaemost* 2009; 102:1-15.
- [95] Furui H, Taniguchi N, Yamauchi K, Sotobata I, Saito H, Inagaki H. Effects of treadmill exercise on platelet function, blood coagulability and fibrinolytic activity in patients with atrial fibrillation. *Jpn Heart J* 1987; 28: 177–84.
- [96] Atalar E, Haznedaroglu IC, Acil T, et al. Patients with paroxysmal atrial fibrillation but not paroxysmal supraventricular tachycardia display evidence of platelet activation during arrhythmia. *Platelets* 2003; 14: 407–11.
- [97] Choudhury A, Chung I, Blann AD, Lip GY. Platelet surface CD62P and CD63, mean platelet volume, and soluble/platelet P-selectin as indexes of platelet function in atrial fibrillation: a comparison of “healthy control subjects” and “disease control subjects” in sinus rhythm. *J Am CollCardiol* 2007; 49: 1957–64.
- [98] Feinberg WM, Macy E, Cornell ES, et al. Plasmin–a2-antiplasmin complex in patients with atrial fibrillation. *ThrombHaemost* 1999; 82: 100–03.
- [99] Marin F, Roldan V, Marco P, et al. Improvement of fibrinolytical function in chronic rheumatic atrial fibrillation after anticoagulation. *Rev EspCardiol* 1999; 52: 25–30.
- [100] Chung MK, Martin DO, Sprecher D, et al. C-reactive protein elevation in patients with atrial arrhythmias. Inflammatory mechanisms and persistence of atrial fibrillation. *Circulation* 2001; 104: 2886–91.
- [101] Dernellis J, Panaretou M. C-reactive protein and paroxysmal atrial fibrillation: evidence of the implication of an inflammatory process in paroxysmal atrial fibrillation. *ActaCardiol* 2001; 56: 375–80.
- [102] Tveit A, Seljeflot I, Grundvold I, Abdelnoor M, Smith P, Arnesen H. Levels of PAI-1 and outcome after electrical cardioversion for atrial fibrillation. *Thromb Res* 2008; 121: 447–53.
- [103] Pretorius M, Donahue BS, Yu C, Greelish JP, Roden DM, Brown NJ. Plasminogen activator inhibitor-1 as a predictor of postoperative atrial fibrillation after cardiopulmonary bypass. *Circulation* 2007; 116: I1–7.

- [104] Oltrona L, Broccolino M, Merlini PA, Spinola A, Pezzano A, Mannucci PM. Activation of the hemostatic mechanism after pharmacological cardioversion of acute non-valvular atrial Fibrillation. *Circulation* 1997; 95: 2003–06.
- [105] Giansante C, Fiotti N, Miccio M, Altamura N, Salvi R, Guarnieri G. Coagulation indicators in patients with paroxysmal atrial fibrillation: effects of electric and pharmacologic cardioversion. *Am Heart J* 2000; 140: 423–29.
- [106] Hatzinikolaou-Kotsakou E, Kartasis Z, Tziakas D, et al. Clotting state after cardioversion of atrial fibrillation: a haemostasis index could detect the relationship with the arrhythmia duration. *Thromb J* 2005; 3: 2.
- [107] Jacob K, Talwar S, Copplestone A, Gilbert TJ, Haywood GA. Activation of coagulation occurs after electrical cardioversion in patients with chronic atrial fibrillation despite optimal anticoagulation with warfarin. *Int J Cardiol* 2004; 95: 83–88.
- [108] Zeuthen EL, Lassen JF, Husted SE. Haemostatic activity in patients with atrial fibrillation treated with low-molecular-weight heparin before and after electrical cardioversion. *J Thromb Thrombolysis* 2004; 17: 185–89.
- [109] Boos CJ, Anderson RA, Lip GY. Is atrial fibrillation an inflammatory disorder? *Eur Heart J* 2006; 27: 136–49.
- [110] Hernandez Madrid A, Moro C. Atrial fibrillation and C-reactive protein: searching for local inflammation. *J Am Coll Cardiol* 2007; 49: 1649–50.
- [111] Engelmann MD, Svendsen JH. Inflammation in the genesis and perpetuation of atrial fibrillation. *Eur Heart J* 2005; 26: 2083–92.
- [112] Gedikli O, Dogan A, Altuntas I, et al. Inflammatory markers according to types of atrial fibrillation. *Int J Cardiol* 2007; 120: 193–97.
- [113] Bruins P, teVelthuis H, Yazdanbakhsh AP, et al. Activation of the complement system during and after cardiopulmonary bypass surgery: postsurgery activation involves C-reactive protein and is associated with postoperative arrhythmia. *Circulation* 1997; 96: 3542–48.
- [114] Conway DS, Buggins P, Hughes E, Lip GY. Prognostic significance of raised plasma levels of interleukin-6 and C-reactive protein in atrial fibrillation. *Am Heart J* 2004; 148: 462–66.
- [115] Chung MK, Martin DO, Sprecher D, et al. C-reactive protein elevation in patients with atrial arrhythmias: inflammatory mechanisms and persistence of atrial fibrillation. *Circulation* 2001; 104: 2886–91.
- [116] Lip GY, Patel JV, Hughes E, Hart RG. High-sensitivity C-reactive protein and soluble CD40 ligand as indices of inflammation and platelet activation in 880 patients with nonvalvular atrial fibrillation: relationship to stroke risk factors, stroke risk stratification schema, and prognosis. *Stroke* 2007; 38: 1229–37.

- [117] Cermak J, Key NS, Bach RR, Balla J, Jacob HS, Vercellotti GM. C-reactive protein induces human peripheral blood monocytes to synthesize tissue factor. *Blood* 1993; 82: 513–20.
- [118] Neumann FJ, Ott I, Marx N, et al. Effect of human recombinant interleukin-6 and interleukin-8 on monocyte procoagulant activity. *ArteriosclerThrombVascBiol* 1997; 17: 3399–405.
- [119] Burstein SA. Cytokines, platelet production and hemostasis. *Platelets* 1997; 8: 93–104.
- [120] Amrani DL. Regulation of fibrinogen biosynthesis: glucocorticoid and interleukin-6 control. *Blood Coagul Fibrinolysis* 1990; 1: 443–46.
- [121] Burstein SA, Peng J, Friese P, et al. Cytokine-induced alteration of platelet and hemostatic function. *Stem Cells* 1996; 14: 154–62.
- [122] Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW. C-reactive protein in healthy subjects: associations with obesity, insulin resistance and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *ArteriosclerThrombVascBiol* 1999; 19: 972–78.
- [123] Acevedo M, Corbalan R, Braun S, Pereira J, Navarrete C, Gonzalez I. C-reactive protein and atrial fibrillation: “evidence for the presence of inflammation in the perpetuation of the arrhythmia”. *Int J Cardiol* 2006; 108: 326–31.
- [124] Conway DS, Buggins P, Hughes E, Lip GY. Relationship of interleukin-6 and C-reactive protein to the prothrombotic state in chronic atrial fibrillation. *J Am CollCardiol* 2004; 43: 2075–82.
- [125] Chung NA, Belgore F, Li-Saw-Hee FL, Conway DS, Blann AD, Lip GY. Is the hypercoagulable state in atrial fibrillation mediated by vascular endothelial growth factor? *Stroke* 2002; 33: 2187–91.
- [126] Seko Y, Nishimura H, Takanashi N, Ashida T, Nagai R. Serum levels of vascular endothelial growth factor and transforming growth factor- β 1 in patients with atrial fibrillation undergoing defibrillation therapy. *Jpn Heart J* 2000; 41: 27–32.
- [127] Webb NJ, Bottomley MJ, Watson CJ, Brenchley PE. Vascular endothelial growth factor (VEGF) is released from platelets during blood clotting: implications for measurement of circulating VEGF levels in clinical disease. *Clin Sci (Lond)* 1998; 94: 395–404.
- [128] Armesilla AL, Lorenzo E, Gomez del Arco P, Martinez-Martinez S, Alfranca A, Redondo JM. Vascular endothelial growth factor activates nuclear factor of activated T cells in human endothelial cells: a role for tissue factor gene expression. *Mol Cell Biol* 1999; 19: 2032–43.
- [129] Morrissey JH. Tissue factor: an enzyme cofactor and true receptor. *ThrombHaemost* 2001; 86: 66–74.

- [130] Archiniegas E, Neves CY, Candelle D, Cardier JE. Thrombin and its protease-activated receptor-1 (PAR1) participate in the endothelial-mesenchymal transdifferentiation process. *DNA Cell Biol* 2004; 23:815–825.
- [131] Vandenbroucke E, Mehta D, Minshall R, Malik AB. Regulation of endothelial junctional permeability. *Ann N Y Acad Sci* 2008; 1123:134–145.
- [132] Gluck N, Schwob O, Krinsky M, Yedgar S. Activation of cytosolic phospholipase A2 and fatty acid transacylase is essential but not sufficient for thrombin-induced smooth muscle cell proliferation. *Am J Physiol Cell Physiol* 2008; 294:C1597–C1603.
- [133] Dahi S, Lee JG, Lovett DH, Sarkar R. Differential transcriptional activation of matrix metalloproteinase-2 and membrane type-1 matrix metalloproteinase by experimental deep venous thrombosis and thrombin. *J Vasc Surg* 2005; 42:539–545.
- [134] Fang Q, Mao L, Kobayashi T et al., PKC δ mediates thrombin-augmented fibroblast-mediated collagen gel contraction. *Biochem Biophys Res Commun* 2008; 369:1199–1203.
- [135] BelAiba RS, Djordjevic T, Bonello S et al., The serum- and glucocorticoid-inducible kinase Sgk-1 is involved in pulmonary vascular remodeling: role in redox-sensitive regulation of tissue factor by thrombin. *Circ Res* 2006; 98:828–836.
- [136] Borensztajn K, Stiekema J, Nijmeijer S, Reitsma PH, Peppelenbosch MP, Spek CA. Factor Xa stimulates proinflammatory and profibrotic responses in fibroblasts via protease-activated receptor-2 activation. *Am J Pathol* 2008; 172:309–320.
- [137] Davis ME, Cai H, Drummond GR, Harrison DG. Shear stress regulates endothelial nitric oxide synthase expression through c-Src by divergent signaling pathways. *Circ Res* 2001; 89: 1073–80.
- [138] Freedman JE, Loscalzo J, Barnard MR, Alpert C, Keaney JF, Michelson AD. Nitric oxide released from activated platelets inhibits platelet recruitment. *J Clin Invest* 1997; 100: 350–56.
- [139] Swiatkowska M, Cierniewska-Cieslak A, Pawlowska Z, Cierniewski CS. Dual regulatory effects of nitric oxide on plasminogen activator inhibitor type 1 expression in endothelial cells. *Eur J Biochem* 2000; 267: 1001–07.
- [140] Cai H, Li Z, Goette A, et al. Downregulation of endocardial nitric oxide synthase expression and nitric oxide production in atrial fibrillation: potential mechanisms for atrial thrombosis and stroke. *Circulation* 2002; 106: 2854–58.
- [141] Goette A, Staack T, Rocken C, et al. Increased expression of extracellular signal-regulated kinase and angiotensin-converting enzyme in human atria during atrial fibrillation. *J Am Coll Cardiol* 2000; 35: 1669–77.
- [142] Choudhury A, Varughese GI, Lip GY. Targeting the renin-angiotensin-aldosterone system in atrial fibrillation: a shift from electrical to structural therapy? *Expert Opin Pharmacother* 2005; 6: 2193–207.

- [143] Healey JS, Baranchuk A, Crystal E, et al. Prevention of atrial fibrillation with angiotensin-converting enzyme inhibitors and angiotensin receptor blockers: a meta-analysis. *J Am CollCardiol* 2005; 45: 1832–39.
- [144] Das UN. Is angiotensin-II an endogenous pro-inflammatory molecule? *Med Sci Monit* 2005; 11: 155–62.
- [145] Tamarat R, Silvestre JS, Durie M, Levy BI. Angiotensin II angiogenic effect in vivo involves vascular endothelial growth factor and inflammation-related pathways. *Lab Invest* 2002; 82: 747–56.
- [146] Suzuki Y, Ruiz-Ortega M, Lorenzo O, Ruperez M, Esteban V, Egido J. Inflammation and angiotensin II. *Int J Biochem Cell Biol* 2003; 35: 881–900.
- [147] Cardin S, Li D, Thorin-Trescases N, Leung TK, Thorin E, Nattel S. Evolution of the atrial fibrillation substrate in experimental congestive heart failure: angiotensin-dependent and -independent pathways. *Cardiovasc Res* 2003; 60: 315–25.
- [148] Takagishi T, Murahashi N, Azagami S, Morimatsu M, Sasaguri Y. Effect of angiotensin II and thromboxane A₂ on the production of matrix metalloproteinase by human aortic smooth muscle cells. *Biochem Mol Biol Int* 1995; 35: 265–73.
- [149] Griending KK, Alexander RW. Oxidative stress and cardiovascular disease. *Circulation* 1997; 96: 3264–65.
- [150] Dzau VJ, Bernstein K, Celermajer D, et al; working group on tissue angiotensin-converting enzyme, International Society of Cardiovascular Pharmacotherapy. The relevance of tissue angiotensin-converting enzyme: manifestations in mechanistic and endpoint data. *Am J Cardiol* 2001; 88: 1–20L.