TALEN-mediated gene editing in epidermolytic ichthyosis patient-derived keratinocytes

Oliver March^{1,2}, Magomet Aushev², Thomas Kocher¹, Ulrich Koller¹ and Julia Reichelt^{1,2}

¹EB House Austria, Research Program for Molecular Therapy of Genodermatoses, Department of Dermatology, University Hospital of the Paracelsus Medical University Salzburg, Austria ² Institute of Cellular Medicine, Framlington Place, Newcastle University, Newcastle, NE2 4HH, UK

Introduction

Epidermolytic ichthyosis (EI) is an inherited skin fragility disorder resulting from dominant-negative mutations in either the keratin 1 (*KRT1*) or keratin 10 (*KRT10*) genes.

Keratins are expressed in pairs, specifically polymerising to build the intermediate filament cytoskeleton of epithelial cells. Dominant-negative mutant keratins integrate into the cytoskeleton, resulting in fragility and collapse upon mild stress. As is the case in EI, this leads to cytolysis and blistering of the skin. Elimination of these mutant keratins is essential for curation of the disease.

Heterozygous parents of patients with recessive EI, express only one *KRT10* allele. Their normal phenotype demonstrates that this is sufficient to support normal skin function.

We are developing an *ex vivo* gene therapy for EI using transcription activator-like effector nuclease (TALEN) technology to knockout mutant *KRT10* alleles in keratinocyte stem cells (KSCs). TALENs are sequence specific nucleases which can be transfected transiently, so as not to persist in treated cells. The specific genome modifications persist and are passed on to the KSC progeny.

Once proof-of-principle is demonstrated, TALEN technology can be applied to a range of genetic skin diseases.

Approach

Figure 1. Schematic of TALEN-mediated allele disruption. TALENs can be designed to bind to specific DNA sequences and induce DSBs. These may be repaired via non-homologous end-joining (NHEJ) which is error-prone and may introduce frameshift mutations. This may induce premature stop codons (PTC) in the target. These mRNAs are targeted for nonsense-mediated decay (NMD), preventing their expression. TALEN 6.21 was designed and engineered to specifically target exon 6 of *KRT10*.

Results

TALEN activity

The activity of TALEN 6.21 was confirmed in an EI cell line (EH31) via T7E1 assay. This relies upon the detection and cleavage of DNA mismatches by T7 endonuclease I in denatured/renatured PCR products of the target site. These mismatches are due to the mutational insertions or deletions introduced by the error-prone NHEJ repair pathway (Figures 1 and 2).

Screening for Modified Clones

Keratinocytes were transfected without selection. They were subsequently seeded and grown clonally before being screened for the correct modification using restriction digestion of the PCR products and Long-Range Allele-Specific PCR.

Acknowledgements

We would like to thank Hans Törmä for providing both the EH31 and NKc11 cell lines. We would also like to thank Thomas Lahaye for enabling the efficient design of our TALENs using the Golden Gate platform that he established. Toni Cathomen and Claudio Mussolino helped to establish TALEN technology in our lab and continue to provide support. This project is funded by Debra Austria and the British Skin Foundation

Figure 2. TALEN activity via T7E1 assay and clonal screening of modified cells. A) TALEN target site, PCR product and restriction sites. B) The presence of two restriction fragments (arrows) of the correct size confirmed activity of TALEN 6.21 at the target site. C) Target site amplified from clonal DNA was subjected to restriction digestion. Loss of a restriction site within the TALEN target site (green arrow) indicates modification of that clone. Approximately 29% of clones were modified (data not shown). Modifications were confirmed and characterised via sequencing. Correctly modified clone AH2 and K10 knockout clone AE5 were isolated for further study.

Figure 3. Analysis of K10 RNA and protein for reversion of disease. A) Sequencing of mutation site (exon 1) and target site (exon 6) in gDNA and cDNA from differentiated monolayers showed loss of mutant and modified RNA in corrected clone AH2. K10 RNA cannot be isolated from knockout clone AE5. B) Western blot analysis of differentiated monolayers reveals K10 protein expression to be closer to both wild-type (NKc11) and skin control in corrected clone AH2, without truncated products. Knockout of K10 was confirmed in clone AE5, α-Actinin included as loading control.

Figure 4. Reversion of disease phenotype. Immunofluorescence staining of K10 (green) and desmoplakin (red) in differentiated cultures following heat shock. Formation of K10 aggregates was demonstrated in diseased cells (arrows) but not in corrected clone AH2. Knockdown of K10 in clone AE5 was further confirmed.

Clone AH2 will be xenografted onto mice to confirm correction *in vivo*. We aim to develop a permanent cure for epidermolytic ichthyosis. We will achieve this by using TALEN 6.21 to phenotypically correct KSCs before transplantation of these cells back into the skin. This technique could then be applied to a wide range of skin diseases.