A Qualitative Analysis of Barriers to Preoperative Aerobic Conditioning in the Thoracic Surgery Population

Hylton D, Hanna WC, Finley CJ, Shargall Y, Agzarian J, and Fahim C

Background

- Postoperative Pulmonary Complications (PPC) are common, yet serious complications of thoracic surgery.¹
- Preoperative Aerobic Conditioning (PAC) through exercise may reduce the incidence of PPC in the thoracic surgery population.²
- Characteristics of the thoracic population have not been considered in the design of existing PAC trials, leading to high attrition rates and non-informative conclusions.
- We sought to identify patient barriers to participation in PAC programs, in preparation for a preconditioning trial.

Methods

- Semi-structured interviews were conducted with patients who underwent lung resection between January and September 2016.
- A 14-item interview guide was developed to evaluate trends in aerobic exercise and assess barriers to participation in PAC programs. Interviews were audiotaped and transcribed verbatim.
- Two researchers analyzed the transcripts using a grounded theory approach to identify emergent themes (Table 1). Descriptive statistics were generated.

PAC Interest Amongst Surgical Lung Cancer Patients

Figure 1. Graphical display of PAC interest in surgical lung cancer patients post-operation.

Results

- Over 200 minutes were transcribed from 21 participant interviews.
- Common barriers to PAC included pain and limited mobility (11/21, 52.4%) and lack of motivation (14/21, 66.7%).
- Qualitative content analysis indicated that patients who exercise periodically demonstrate more interest in a PAC program, as compared to those who do not exercise, or those who exercise regularly.
- When asked, 85.7% (18/21) of participants were agreeable to using a wearable activity tracker as part of a home-based PAC program.

Table 1. Qualitative Content Analysis: PAC Interest & Emergent Themes

Emergent Themes	Highlighted Participant Quotes
Participants did not exercise due to pain, limited mobility, and lack of motivation.	“I have no motivation to exercise, I just can’t do it.”
Strong family and surgeon support cultivated feelings of empowerment.	“Well, I was certainly put in good hands and so I felt that I was going to be fine.”
Participants believed setting post-operative goals provided additional motivation to recover. Common goals were: returning to ‘baseline’ health and exercising more.	“I started pushing a lot more towards the end of the season, trying to get back into normal activity.”
Participants that did not exercise pre-surgery were unlikely to be interested in PAC, those that exercised periodically pre-surgery were interested in PAC, those that exercised regularly pre-surgery were unsure of any additional PAC benefit.	“I was already doing a lot so participating in that program - I don’t know how much that would have necessarily benefited me.”
Participants that expressed personality traits relating to optimism and confidence were more likely to express an interest in PAC.	“If you prepare for it physically and mentally obviously you’ll be better off.”

Conclusion

- A successful PAC program should be tailored to participants’ baseline level of exercise and should be designed to address barriers related to lack of motivation.
- These findings will be used to design a pragmatic, patient-informed, prospective trial titled “Move for Surgery” to assess the impact of PAC using wearable technology in the thoracic population.

References

1. Agostini, P., Cieslik, H., Rathinam, S., Bishay, E., Kalkat, M., Rajesh, P., Steyn, R., Singh, S., & Naidu, B. (2010). Postoperative pulmonary complications following thoracic surgery: are there any modifiable risk factors. *BMJ*, 65(9): 815-818.

2. Hoogeboom, T., Dronkers, J., Hulzebos, E., van Meeteren, L. (2014). Merits of exercise therapy before and after major surgery. *Curr Opin Anaesthesiol.*, 27(2):161-166.