Influence effect of breastfeeding commitment to quality of life in nursing mothers

A. Aminova, Z. Gumbatova , S. Lahova , S. Guseva , G. Abdullaeva
Sechenov First Moscow State Medical University, pediatric faculty, Moscow, Russia.

Background and Aims: Data on the impact of breastfeeding on the quality of life of mothers are not available. The aim of the work is to study the impact of commitment to breastfeeding on the quality of life of nursing mothers.

Method: A survey was conducted of 510 women with infants. SF-36 questionnaires were used to assess the quality Of all patients were divided into two groups: 1 - women who breastfed (260 women) and 2 - children who are not breastfed (250).

Results: The average income of families of non-lactating women is almost 2 times more than in non-lactating families ($p<0.01$). Indicators of physical functioning (PF) of women in the main group and the comparison group did not different statistically ($p>0.05$). At the same time, role functioning due to physical condition (RP) in lactating women was significantly lower than in the comparison group ($p<0.05$), that is, they make less effort to perform physical work and, probably, the physical condition of lactating women has less impact on their daily activities. Pain intensity factor (BP) and its impact on the ability to engage in daily activities, including work at home and outside the home, was almost 2 times lower in nursing mothers than in non-lactating ($p<0.05$). The physical and emotional state significantly less affected of social activity and communication (SF) in the women of the main group than in the comparison group ($p<0.05$). According to the questionnaire, the mental health index (MH) of nursing mothers showed that they are less prone to depression and have a higher rate of positive emotions than women who are not breastfeeding ($p<0.01$).

The RE index in the comparison group was significantly higher than in the main group ($p<0001$). As for vital activity (VT) and subjective assessment (GH) of one's own health, we have not found statistically significant differences in the compared groups ($p>0.05$). Women, regardless of their commitment to breastfeeding, did not report ill health after childbirth.

Conclusion: It is established that the higher the material well-being of the family, the less committed women are to breastfeeding. Subjective feelings of physical condition (functioning), indicators of vital activity (ME) and subjective assessment (PR) of own health were identical both at nursing, and at non-lactating mothers. Nursing women were characterized by high endurance in terms of RP and BP, rarely experienced difficulties in daily activities due to their physical condition and pain, physical and emotional state did not interfere with communication and social activity, they are less prone to depression and have a higher rate of positive emotions.

Characteristics of groups

Groups	Nursing women	Non-lactating women	p
Average age of women (years)	30,16+2,25	32,4+1,86	>0,05
Material welfare (rub)	22500,0+3,800	43958,33+4,240	<0,05