The time for extubation after stopping infusion of propofol depends on the amount of bolus administration of fentanyl at the end of surgery, but dose not depend on ketamine administered in a surgery with AOPRfK(air-oxygen-propofol-remifentanil-low dose ketamine) anaesthesia.

MASASHI UCHIDA, M. D.

Department of Anaesthesiology and Pain Clinic, Saitamaseikeikai Hospital, Higashimatsuyama, Saitama, Japan

Background and goal of study

It is well known that AOPFK anaesthesia has an advantage in postoperative analgesia. However, AOPFK anaesthesia sometimes sets up a delay of awareness from anaesthesia. Nowadays, continuous infusion of remifentanil(Rf), instead of fentanyl(F), is very common. However, in AOPRfK anaesthesia, the same phenomenon sometimes sets up. Therefore, we investigated what the main reason of the phenomenon in spinal surgery.

Materials and methods

The study style is retrospective. 11 patients (male/female=9/2), ASAII-II were included in the study. We deleted one patient in the analysis on this e-poster due to massive bleeding during the operation and added another patient in the analysis on this e-poster. They were 58 ± 15 (mean±standard deviation) years old. The operation time was 192 ± 51 min. The induction of anaesthesia was performed with bolus infusion of 1mg midazolam, 2mg/kg propofol(PO), 50mg rocuronium (Rb), and $0.2\mu\text{g/kg}$ Rf. In 9 patients, bolus Ketamine(Ket) was administered (0.34 ± 0.27 mg/kg) at the induction. The maintenance of anaesthesia was performed with continuous infusion of PO, Ket (0.42 ± 0.12 mg/kg/h), Rf($0.1\text{--}0.3\mu\text{g/kg/min}$), and Rb. PO and Rf infusion rate were adjusted to maintain bispectral index between 40 and 60. When a surgery ended, continuous infusion of PO, Ket, Rf, and Rb were stopped and bolus $100\text{--}200\mu\text{g}$ F and 50mg flurbiprofen were administered. 10–20 mL ropivacaine or revobupivacaine were given as infiltration anaesthesia when the operative wound was closed. The time between stopping PO infusion and the extubation (Time)(minutes) was recorded. Regression lines were drawn in the amount of F per body weight vs. Time (RL1) (Fig. 2), the amount of F per body weight vs. effect-site concentration change of PO from stopping infusion of PO to extubation (RL2) (Fig. 3), the total amount of Ket administered per body weight (mg/kg) vs. Time(RL3) by using SigmaPlot 11.0. (Sistat Software Inc., San Jose, CA, USA). Effect-site concentration of PO and F were simulated by TIVAtainer (ver 8) (EuroSIVA).

Diagnosis and operative method

Lumbar spinal stenosis: 4 patients, operative methods: lamynectomy + spinal fusion surgery 1 patients, enlarged window surgery 3 patient, lumber vertebrae herniated disk: 4 patients, operative method: nucleotomy 3 patient, enlarged window operation 1 patient, cervical myelopathy: 1 patients, operative method: laminaplasty + posterior diskectomy 1 patient, L4 spondylolistesis 1 patient, operative method: spinal fusion surgery 1 patient.

Results and discussion

Time was 21 ± 6.8 minutes. The total amount of Ket was 1.32 ± 0.58 mg/kg. The coefficient of correlation value of RL1, RL2, and RL3 is 0.7078, 0.4259, and 0.0563 respectively. According to RL1, it is suggested that F less than $2\mu\text{g/kg}$ dose not set a prolongation of Time. The total amount of Ket administered per body weight dose not contribute to the phenomenon with given dose in the study.

conclusion

The amount of F given at the end of surgery prolongs Time in a dose dependent manner in AOPRfK (low dose ket) anaesthesia.

Fig. 1 fCe vs pCe at extubation

Fig. 2 total F / body weight vs. Time

Fig. 3 total F / body weight vs. pCe change from stopping infusion of PO to extubation