PROTOTHECA ZOPFII
CAUSES BLOODSTREAM INFECTION IN
IMMUNOCOMPROMISED PATIENT

P211

Fernández Norma¹, Taverna Constanza², Farías Luciana¹, Paravano Lucia¹, Sierra Mariela¹,
Córdoba Susana ², Tiraboschi INora¹, Stecher Daniel ·

¹ Infectious Disease Division. Hospital de Clínicas “José de San Martín”. Universidad de Buenos Aires.

² INEI- ANLIS “Dr. C G Malbrán”.

Buenos Aires. Argentina

Introduction

Prototheca are common in nature and human are undoubtedly often exposed to them but appear to have relatively little ability to infect the human host. A revision of 160 protothecosis human cases reported from 1966 to 2011 showed that *Prototheca wickerhamii* was responsible of 115 cases, only 11 cases were due to *Prototheca zopfii* and 34 were not speciated (1). Disseminated protothecosis is a rare disease that occurs in patients with primary immunodeficiency, hematological neoplasia, organ transplantation, diabetes mellitus and autoimmune diseases with variable therapeutic success.

Objective

To describe disseminated protothecosis in a patient with acute lymphoblastic leukemia with favorable outcome.

Methods

Clinical case: a 19-year-old man no medical history background

Results

Patient received amphotericin B 10 days and outcome was favorable. He remained hospitalized until he received a new treatment regimen for his underlying disease. The strains grew well at 37 °C and 40 °C. On SDA after 7 days at 28°C: white colonies 10 mm of diameter with irregular margin, rough surface. Fermentation(+) Glucose, Galactose, Sucrose, Maltose (-) Trehalose. Assimilation (+)Glucose (-)Galactose, L-Sorbose, Sucrose, Maltose, Cellobiose, Trehalose, Lactose Melibiose, Raffinose, Melezitose, D -Xylose, L-Arabinose, D-Ribose, D-Mannitol, Inositol, L-Rhamnose, Citrate and Nitrate. Urease test negative.

MIC Fluconazole 128 mg/L Voriconazole 0.13 mg/L Itraconazole 0.06mg/L AmphotericinB 0.25 mg/L Posaconazole >32 mg/L

Neighbor Joining method and Tamura-3 parameter model. Numbers at nodes are percentage bootstrap from 1000 replicates.

Prototheca zopfii* var. *hydrocarborea

Conclusions

Optimal antifungal treatment of protothecosis is controversial as in vitro MIC are difficult to interpret due to established clinical cut-off points lack. Sequence analysis showed 100 % identity with the strain *Prototheca zopfii* var. *hydrocarborea* RND16, a strain isolated from a hot spring (2). Only one human case of meningitis due to this *P. zopfii* variety has been described(3). Here we described the first case of human blood infection due to *P. zopfii* var. *hydrocarborea* in an immunocompromised host with a favorable outcome after amphotericin B antifungal treatment.

References: 1-Todd J, King J, Oberle A, Matsumoto T, Odaka Y, Fowler M, Pore RS, Shahan TA, Yin Y, Sanusi I. Protothecosis: report of a case with 20-year follow-up and review of previously published cases. Medical Mycology 2012; 50:673-689. 2-Ueno R, Urano N, Suzuki M, Kimura S. Isolation, characterization, and fermentative pattern of a novel thermotolerant *Prototheca zopfii* var. *hydrocarborea* strain producing ethanol and CO₂ from glucose at 40 °C. Arch Microbiol 2002;177 :244-250. 3-Liu S, Ran Y, He X et al. Identification of a strain *Prototheca zopfii* var. *hydrocarborea* by analyzing the sequence of ribosome RNA gene. Chin J Zoon 2007; 23: 656-660.