IS EPIDURAL CATHETER THE BEST CHOICE TO ANAESTHETIZE A PARTIALLY OBSTRUCTED

CENTRO
HOSPITALAR
DE LISBOA
CENTRAL, EPE

AIRWAY FOR AN AWAKE FIBEROPTIC TRACHEAL INTUBATION?

Pimenta, Marta; Andrade, Diogo; Rodrigues, Marta; Gonçalves, Margarida; Canas, Margarida; Matos, Francisco
Centro Hospitalar Lisboa Central, Portugal

Euroanaesthesia 2017
THE EUROPEAN ANAESTHESIOLOGY CONGRESS
GENEVA, 03-05 JUNE

BACKGROUND

A common cause of death in patients with deep neck infections is acute loss of the airway during interventions to control it.³

Tracheostomy using local anaesthesia has been considered the gold standard of airway management in these patients. However, tracheostomy site may be involved in the infection and this procedure also conveys substantial risk of infection spread to the mediastinum and tracheal stenosis as a late complication.³

Nowadays fiberoptic intubation using topical anaesthesia is the first choice for airway control in adult patients with deep neck infections.³

The successful conduct of awake fiberoptic intubation is dependent upon effective local anaesthesia. Sprayed lidocaine onto the larynx and trachea during fiberoptic intubation through an epidural catheter produces excellent local anaesthesia.⁴

CASE REPORT

- ♂, 71 years

Past Medical History:

- Hypertension
- Hypercholesterolemia

Emergency Department:

- Altered mental status
- Mild hypoxemia (oxygen saturation 89%) with face mask oxygen therapy 15 L.min⁻¹

Diagnosis:

Retropharyngeal Abscess

Proposed Therapy:

Emergent surgical drainage of retropharyngeal abscess

Preoperative Evaluation:

ASA II

Anticipated difficult airway with partial obstruction

Airway approach:

Nose and pharynx were anaesthetized with topical lidocaine

Fiberscope was introduced through the nostril and progressed until the oropharynx.

An epidural catheter was advanced through the working channel of the fiberscope

40 mg of lidocaine 2% were injected in the epidural catheter to atomize the local anaesthetic into the larynx and trachea

DISCUSSION

Tracheal intubation in patients with deep neck infections is challenging. The distorted airway anatomy, tissue immobility, and limited access to the mouth make orotracheal intubation with rigid laryngoscopy difficult.³ The greatest risk with oral or nasal intubation is rupture of the abscess with aspiration of purulent fluid, which can be devastating, therefore awake intubation is preferable.^{1,3}

Deep neck infections cause hyperemia and edema of the pharyngolaryngeal mucosa, which may enhance transmucosal systemic absorption of local anaesthetic, thus increasing risk of lidocaine toxicity.

Among the various techniques described to anaesthetize the larynx and trachea, local anaesthetic spray through epidural catheter provides several advantages: availability, non-invasive, reliability and safety reaching areas beyond airway obstruction, as well as reduced need of local anaesthetic doses.

Local anaesthetic nebulization directly through the fiberscope working channel would probably be ineffective due to airway obstruction.

In case of infection, invasive procedures, such as nerve blocks or trans-tracheal injection, pose the risk of significant complications: in the event of abscess perforation or leakage of pus to the airway, which may have catastrophic consequences.

CONCLUSION

Patients with deep neck infections present challenging airways for an anaesthesiologist. Airway compromise should always be considered and the potential for abscess rupture is ever-present.³

Adequate preparation with local anaesthesia of the upper airway in combination with a safe sedation scheme is essential to reducing airway irritation, cough, and laryngeal spasm.^{3,5}

This case reflects anaesthesia of the airway through an epidural catheter which proved to be a successful and safe method to anaesthetize the larynx and trachea for awake fiberoptic tracheal intubation in a quite critical situation.

References: 1. Heindel, D. (1987). Deep Neck Abscesses in Adults. *Anaesthesia & Analgesia*, 66(8), pp.774:776. 2. Long, T. and Wass, C. (2004). An Alternative to Transtracheal Injection for Fiberoptic Intubation in Awake Patients: A Novel Noninvasive Technique Using a Standard Multiorifice Epidural Catheter through the Bronchoscope Suction Port. *Anesthesiology*, 101(5), p.1253. 3. Ovassapian, A., Tuncbilek, M., Weitzel, E. and Joshi, C. (2005). Airway Management in Adult Patients with Deep Neck Infections: A Case Series and Review of the Literature. *Anaesthesia & Analgesia*, 100(2), pp.585-589. 4. Sethi, N., Tarneja, V., Madhusudanan, T. and Shouche, S. (2005). Local Anaesthesia for Fiberoptic Intubation : A Comparison of Three Techniques. *Medical Journal Armed Forces India*, 61(1), pp.22-25. 5. Simmons, S. and Schleich, A. (2002). Airway Regional Anaesthesia for Awake Fiberoptic Intubation. *Regional Anesthesia and Pain Medicine*, 27(2), pp.180-192.