Impact of resuscitation duration on long-term functional outcome after out-of-hospital cardiac arrest

Z. Nehme^{1,2}, R. Nair¹, E. Andrew^{1,2}, S. Bernard^{1,2}, K. Smith^{1,2,3}

¹ Ambulance Victoria, Melbourne, Australia; ² Monash University, Melbourne, Australia; ³ University of Western Australia, Perth, Australia.

Purpose

There is increasing community interest in the long-term prognosis after out-of-hospital cardiac arrest (OHCA). We examine the impact of cardiopulmonary resuscitation (CPR) duration on 12-month survival with good functional recovery.

Methods

Between 2010 and 2016, consecutive adult non-traumatic OHCA patients were identified from the Victorian Ambulance Cardiac Arrest Registry. Survivors to hospital discharge underwent follow-up at 12 months post-arrest to determine functional outcome using the Glasgow Outcome Scale–Extended (GOSE).

CPR duration was defined as the total emergency medical service CPR time either before return of spontaneous circulation or termination of resuscitation. Adjusted logistic regression was used to assess the effect of CPR duration on 12-month survival with good functional outcome.

Results

Of the 13,898 attempted resuscitations, 710 (5.1%) survived to 12 months with good functional outcome. The median CPR duration was 17 minutes overall (interquartile range: 9, 33), but was lower in 12-month survivors with good functional outcome (7 vs. 20 minutes; $p < 0.001$).

The 99th cumulative proportion of 12-month survivors with good functional outcome was reached at a CPR duration of 41 minutes overall (95% CI: 34, 47), 42 minutes (95% CI: 35, 48) in patients with initial shockable rhythms, and 26 minutes (95% CI: 20, 26) in patients with initial non-shockable rhythms (see Figure 1).

Every minute increase in CPR duration was associated with a 10.9% reduction in the risk-adjusted odds of good functional recovery at 12 months (OR 0.89, 95% CI: 0.88, 0.90, $p < 0.001$). Figure 2 shows the predicted probability of 12 month survival with good functional outcome according to CPR duration, holding other variables at their mean.

“Increasing CPR duration is associated with poorer long-term functional recovery after OHCA. However, long-term favourable outcomes are possible with CPR durations >40 mins.”

Figure 1: Cumulative proportion of CPR duration attempts in 12 month survivors with good functional outcome, stratified by initial rhythm.

Figure 2: Predicted probability of 12 month survival with good functional outcome according to CPR duration and initial rhythm, holding other variables at their mean.