The impact of blood group on survival following critical illness

Slade, R¹., Alikhan, R²., Wise, MP¹., Germain, L¹., Morgan, MPG¹.
University Hospital of Wales Critical Care Directorate¹; Department of Haematology²

Introduction

Predicting patient outcomes following critical illness is challenging. Although genetic variation is known to be one factor in determining survival in intensive care, it is difficult to measure in clinical practice (1). Recent evidence has suggested that patients with blood group AB are more likely to survive following major cardiac surgery and this is associated with a reduced number of blood transfusions (2). There is a body of evidence which suggests that blood group AB has higher levels of Von-Willebrand factor and factor VIII (3,4), making this group more pro-thrombotic, which may be beneficial in patient groups where bleeding risks are high. However, there is no current data to indicate whether a patient's blood group affects general intensive care outcomes.

Objectives

The objective of the study was to determine if ABO blood group affects survival in intensive care. The primary outcome measure was 90-day mortality with a secondary outcome measure of the percentage of patients receiving a blood transfusion.

Materials and Methods

We performed a retrospective cohort analysis of all patients admitted to the Intensive Care Unit of a major tertiary University Hospital between 2006 and 2016. Baseline characteristics for each blood group were collected including, age, gender, BMI, surgery during admission, days in hospital, ethnicity and APACHE II score. 90-day survival was estimated by Kaplan-Meier plots. ABO blood groups were individually compared and then pooled to compare AB vs non-AB blood groups.

Figure 1: ICU survival by Blood Group

Figure 2: ICU survival blood group AB vs non-AB

Results

7340 patients were included in the study. Blood group AB accounted for 3% (221), A 41% (3008), B 10.6% (775) and O 45.4% (3336). These values are similar to UK averages (5).

Blood group AB had the greatest survival benefit when assessing per individual blood group **Figure 1** (blood group AB 90-day survival estimate 76.75, 95% CI 72.89- 80.61 with the overall estimate 72.07, 95% CI 71.31-72.82) (Log rank Chi square 16.128, p=0.001). Transfusion requirements were similar in all groups with no significant difference between the percentages of patients transfused (AB 23.1%, A 21.5%, B 18.7%, O 19.9%, Pearson chi square 5.060 p=0.167).

When pooling blood group data, Kaplan-Meier analysis showed blood group AB (n=221) to have improved survival compared to non-AB blood groups **Figure 2** (n=7119) (blood group AB 90-day survival estimate 76.75, 95% CI 72.89- 80.61 non-AB groups 71.92, 95% CI 71.15-72.69) (log rank Chi Square 3.890, p=0.049).

Conclusion

- Intensive care patients with blood group AB have a higher 90-day survival compared with other blood groups.
- There was no correlation between blood group and percentage of patients receiving transfusion. Blood group AB is relatively rare accounting for 3% of the study population and therefore our study may have been underpowered to determine this bleeding risk.
- Improved survival outcomes in blood group AB are a promising area for further study including any mechanistic basis for this finding.