USEFULNESS OF N-TERMINAL PRO B-TYPE NATRIURETIC PEPTIDE FOR ATRIAL FIBRILLATION SCREENING IN PRIMARY STROKE PREVENTION

Alejandro Bustamante, Josep Luis Clua, Juan Acosta, Felipe González, Juan Ballesta-Ors, Andrea Caballero, Anna Penalba, Eulalia Muria-Subirats, Elena Pala, Pilar Delgado, Jorge Pagola, Alonso Pedrote, Miguel Ángel Muñoz, Joan Montaner
The Atrial Fibrillation Research In CATalonia (AFRICAT) investigators

INTRODUCTION

- Atrial fibrillation (AF) screening programs have not shown a clear usefulness in stroke prevention
- Blood biomarkers such as N-terminal pro B-type natriuretic peptide (NT-proBNP) have been proposed as a useful element to identify subjects at high risk
- Given its temporal profile after stroke and after AF paroxysms¹, NT-proBNP might act as a signal of the paroxysm standing for longer than the “electric” sign(biological memory)
- In this study we aimed to test whether NT-proBNP differentiates between different patterns of AF in a high risk population

METHODS

- The Atrial Fibrillation Research In CATalonia (AFRICAT) aims at developing a screening program for AF in subjects aged 65-75 with previous hypertension and diabetes by combining clinical information, blood biomarkers and electronic devices
- For the pilot phase, 100 subjects were randomly selected
- ECG and blood samples were obtained at a single visit
- Additionally, all patients were monitored for four weeks with a wearable Holter device (Nuubo TM)
- NT-proBNP was measured by immunoassay (Roche Laboratories)

RESULTS

DESCRIPTION OF THE COHORT

	All (N=100)	AF+ (20)	AF- (80)	p
Age	70 (68-73)	69 (66-71.5)	70 (68-73)	0.273
Sex (% male)	67	13	54	0.832
Alcohol	11	1	10	0.456
Tobacco	20	4	16	1.000
Dyslipidemia	81	16	65	1.000
Coronary disease	18	8	10	0.008**
CHF	3	3	0	0.007**
Valvulopathy	4	3	1	0.024**
Previous stroke	6	2	4	0.597

* PMH: past medical history

CONCLUSIONS

- Measurement of NT-proBNP is a useful tool to be included in AF screening programs in primary care
- The recommended cut-offs for CHF lack sensitivity. The proposed cut-off of 95 mg/mL is promising but requires further validation
- In patients with NT-proBNP levels > 95pg/mL and no AF in ECG, wearable Holters might have a role in paroxysmal AF detection

REFERENCES

- Wozakowska-Kaplon et al. Am J Cardiol. 2004;93:1555-8
- Verdú et al. Rev Esp Cardiol. 2012;65:613-9
- ESC. Guidelines CHF 2012

CONTACT

alejandro.bustamante@vhir.org
www.lin-bcn.com
www.stroke-biomarkers.com

The study is supported by
Fundació La Marató de TV3,
Grant number: 201528-30-31-3