Cardiac Arrhythmias and In-hospital Mortality in 63 Cases of Ischemic Stroke Involving the Insula

Maria Mirabela Manea^{1,2} D. Dragoș^{2,3} C. Popa^{1,2} S. Tuță^{1,2}

1. National Institute of Neurology and Neurovascular Diseases Bucharest
2. UMF Carol Davila, Bucharest
3. Emergency University Hospital Bucharest

OBJECTIVE

Several studies on acute ischemic stroke patients sustain that insula involvement correlates with both electrocardiographic abnormalities and mortality, the former mediating the latter. However there is not universal agreement upon this conclusion, other studies invalidating it (1),(2).

To test whether insula involvement and insula lateralization in ischemic stroke correlates with ST displacement and cardiac dysrhythmias (as recorded by continuous 24 hours electrocardiographic monitoring) and with in-hospital mortality.

METHODS:

Prospective study on 110 patients admitted to stroke unit with acute (<24 hours from onset) ischemic stroke with no history of arrhythmias, including atrial fibrillation (AF), or severe heart failure. We analyzed the data provided by continuous 24 hours electrocardiographic monitoring performed within 48 hours of stroke onset. For the detection of AF we extended the surveillance period to the first 5 days after stroke onset.

RESULTS (1)

The average age was 67.2 +/-12.93 years. The proportion male: female was 64.55%:35.55% (fig.1). The etiology of ischemic stroke was classified based on the TOAST (Trial of Org 10172 in Acute Stroke Treatment) criteria as: cardioembolic (29,1%), large artery atherosclerosis including atheroembolic (50,9%), small-vessel lacunar (3,6%), other causes (1,8%) and undetermined / cryptogenic (14,5%) - fig.2.

Insular lesions were detected in 63 of 110 (57.27 %) patients (fig.3). New AF was present in 35 of 110 (31.81%) - patients - fig.4 (29.58% in males and 35.90% in females, the difference being statistically insignificant). Significantly more VPC couplets (P 0.045) - fig.5, bigeminism (p 0.049) - fig.6 were detected in patients with cardioembolic stroke. AF detection was not associated with insula infarction.

RESULTS (2)

Regarding ST displacement, only ST depression was correlated with insula lateralization, being greater for patients with left insular injury (p 0.02) - fig.7.

There was no relationship between insular lesion and other ECG changes (SPVC, VPC, VPC couplets, bigeminism, trigeminism, VPC run, SVPC run) or pulse rate.

There was a 12.72% in-hospital mortality rate (14 out of 110 patients). All deaths occurred 72 hours after stroke onset or later. There was a tendency for the mortality to be correlated with the number of VPC runs - fig.8, but this correlation failed to reach statistical significance (p = 0.08).

Insula involvement was not a risk factor for death, nor were insula lateralization and AF.

CONCLUSIONS

These findings support only the relationship between ST depression and insula lesion, but do not confirm that insular infarction predicts electrocardiographic abnormalities with potential prognostic implications during hospitalization.