Proximal region of carotid atherosclerotic plaque shows more intraplaque hemorrhage: the Plaque At RISK (PARISK) study

Geneviève AJC Crombag^{1,2}, Floris HBM Schreuder³, Faysal Benali¹, Dianne HK van Dam-Nolen⁴, Madieke I Liem⁵, Carlo Lucci⁶, Anton F van der Steen⁷, Mat JAP Daemen⁸, Werner H Mess⁹, Aad van der Lugt⁴, Paul J Nederkoorn⁵, Jeroen Hendrikse⁶, Paul AM Hofman¹, Robert J van Oostenbrugge^{2,10}, Joachim E Wildberger^{1,2}, M Eline Kooi^{1,2}

Background and aims:

Intraplaque haemorrhage (IPH) contributes to enlargement of the lipid core and plaque progression, leading to plaque destabilization. The mechanisms that contribute to IPH development are not completely understood. We hypothesise that development of IPH is affected by biomechanical and hemodynamic variables and that fissuring or rupture of the fibrous cap can contribute to IPH. The biomechanical load and wall shear stress are highest in the proximal region of the plaque. We aim to study non-invasively a possible difference in volume of IPH in the proximal region of the plaque.

Methods:

88 symptomatic stroke patients with carotid stenosis included in the Plaque At RISK (PARISK) study demonstrated ipsilateral carotid IPH on multi-sequence MR images. Inner and outer vessel wall contours and IPH area (hyperintense signal on MPRAGE/3D-T1-weighted fat suppressed spoiled gradient echo images in the bulk of the plaque) were delineated on the MR images by trained observers. IPH area was calculated with respect to the total vessel wall area. Fibrous cap status was scored by comparing pre- and post-contrast black blood TSE images. Differences in mean IPH percentages between proximal and distal regions were calculated using a paired samples t-test, while a McNemar test was used to reveal differences in proportions of a thin and/or ruptured fibrous cap.

Conclusion:

We demonstrated that IPH is more prevalent on the proximal side of the plaque compared to the distal side. This may indicate that wall stress and wall shear stress play an important role in the development of IPH.

Results:

We found significantly larger areas of IPH in the proximal part of the plaque at 2, 4, and 6 mm from the position of maximal luminal narrowing, respectively: 14.4% vs. 9.6% ($p=0.04$), 14.7% vs. 5.4% ($p < 0.001$), and 11.1% vs. 2.2% ($p=0.001$). Additionally, we found an increased proximal prevalence of a thin and/or ruptured fibrous cap on MRI at 2, 4, 6, and 8 mm from the MRI slice with maximal luminal narrowing, respectively; 33.7% vs. 18.1%, $p=0.007$; 36.1% vs. 7.2%, $p < 0.001$; 33.7% vs. 2.4%, $p=0.001$; 30.1% vs. 3.6%, $p=0.022$.

¹ Department of Radiology and Nuclear Medicine, MUMC+, Maastricht; ² CARIM School for Cardiovascular Diseases, Maastricht University, Maastricht; ³ Department of Neurology & Donders Institute for Brain Cognition & Behaviour, Radboud UMC, Nijmegen; ⁴ Department of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center Rotterdam, Rotterdam; ⁵ Department of Neurology, Amsterdam University Medical Centres, University of Amsterdam, Amsterdam; ⁶ Department of Radiology, Department of Radiology, University Medical Center Utrecht, Utrecht; ⁷ Department of Biomedical Engineering, Erasmus MC, University Medical Center Rotterdam, Rotterdam; ⁸ Department of Pathology, Amsterdam University Medical Centres, University of Amsterdam; ⁹ Department of Clinical Neurophysiology, Maastricht University Medical Center, Maastricht; ¹⁰ Department of Neurology, Maastricht University Medical Center, Maastricht, The Netherlands