Sleep Characteristics of Pediatric Burn Patients

Esra ARDAHAN AKGÜL, Hatice YILDIRIM SARI
İzmir Kâtip Çelebi University, Faculty of Health Sciences, Department of Pediatric Nursing

Pediatric burns are the third most common causes of accident-induced mortality following motor vehicle accidents and drowning. After burn, patients are exposed to many stressors that lower sleep quality and interrupt sleep. Treatment-associated factors such as fixation methods, physiological factors, therapeutic interventions, diagnostic procedures, mechanical ventilation, sedatives, analgesic and anesthetic drugs are known to affect sleep. Thermal injuries cause increases in sleeplessness, and significant reductions in non-REM sleep and REM sleep.

Aim: This present study was planned to determine sleep characteristics, sleeplessness and sleep habits in pediatric burn patients.

MATERIALS METHODS

Settings:The study was designed as a descriptive one and carried out with 96 children who were 2-6year-old in Pediatric Surgery Department/Burn Unit for 2-7 days. For data collection; Demografic Data Collection Form, Sleep Problems Characteristics and Identification Survey which has been created by researchers are used. Data were collected by face to face interviews with mothers. Questions were asked to compare pre-post burn sleep quality. At the age of 0-2; there are multiple phase sleep patterns and an equal number of sleeping and awakening periods during the day and night. Babies are polyphasic, preschool children sleep biphasic. Physiological biphasic sleep pattern becomes a monophasic night sleep with the necessity of school life. For this reason, children aged 2 to 6 years with biphasic sleep were studied.

The data obtained from the study were analyzed using the SPSS version 21.Statistical significance was established at a p value <0.05.

RESULTS

The participating children’s mean age was 3.36 ± 1.39 and of the children 53.1% were male The part of the body exposed to burns most in children (10.4%) was the "hands". Of the children in the study, 79.2% (n = 76) had second-degree burns. The mean percentage of burns was 8.12% ± 5.61 (min: 1%, max 30%). Of the causes of burns in children, 47.9% were hot water.

Pre- and Post-Burn Sleep Characteristics-I

		Mean	SD	Min.-Max. Values	t	p
Wake up time in the morning	Pre-burn	8:48 a.m.	1.14	Min:5	10.023	<0.001
				Max: 11		
	Post-burn	6:56 a.m.	1.09	Min:1		
				Max: 9		
Duration of night sleep	Pre-burn	10.83 hours	1.13	Min:6 hours	13.644	<0.001
				Max: 12 hours		
	Post-burn	7.44 hours	1.79	Min:3 hours		
				Max: 12 hours		
Duration of daytime sleep	Pre-burn	1.46 hours	1.07	Min: 0 hours	-1.307	.194
				Max: 3 hours		
	Post-burn	1.67 hours	1.01	Min: 0 hours		
				Max: 6 hours		
Duration of daily sleep	Pre-burn	11.56 hours	1.40	Min: 8 hours	8.715	<0.001
				Max: 15 hours		
	Post-burn	9.13 hours	2.07	Min: 4 hours		
				Max: 15 hours		

Pre-Post Burn Sleep Characteristics –II

		Pre-burn		Post-burn		p
		n	%	n	%	
What time does the child to go to sleep?	7:00 p.m. - 9:59 p.m.	8	8.3	12	12.5	12.938**
	10:00 p.m. - 11:59 p.m.	78	81.3	43	44.8	
	After 12 midnight	10	10.4	41	42.7	
Does the child have interruptions of sleep?	Yes	61	63.5	92	95.8	<0.001
	No	35	36.5	4	4.2	
How many times does the child have interruptions of sleep?	Once	32	32.5	5	5.2	55.000**
	Twice	16	26.2	13	14.1	
	3 times	13	21.3	22	23.9	
	4 or more times	-	-	52	56.5	
Does the child have trouble falling asleep?	Yes	52	54.2	79	82.3	<0.001
	No	44	45.8	17	17.7	
How long does it take the child to fall asleep?	Less than 15 minutes	24	25.0	17	20.7	22.349**
	15-30 minutes	25	26.0	13	15.9	
	30-45 minutes	4	4.2	31	37.8	
	45-60 minutes	4	4.2	19	23.2	
	More than 60 minutes	-	-	2	2.4	
Does the child wake up tired in the morning?	Yes	12	12.5	58	60.4	<0.001
	No	84	87.5	36	38.3	
Does the mother have trouble awakening the child in the morning?	Yes	33	34.4	55	57.3	<0.001
	No	63	65.6	41	42.7	
How long does it take to awaken the child?	Less than 15 minutes	22	66.7	16	16.7	15.000**
	15-30 minutes	9	27.3	34	35.4	
	30-45 minutes	2	6.1	6	6.3	
	45-60 minutes	-	-	2	2.1	
Dos the mother think that the child has sleep problems?	Yes	35	36.5	78	81.3	<0.001
	No	61	63.5	18	18.8	

* Analysis was performed with the McNemar's test. ** Analysis was performed with the Friedman test.

Comparison of Pre-Post Burn Sleep Habits

	Pre burn		Post burn		p	
	Never	Frequently	Never	Frequently		
	Occasionally	Every night	Occasionally	Every night		
The child;	n	%	n	%	n	%
has trouble falling asleep (needs a parent)	69	71.9	27	28.1	12	12.5
has trouble falling asleep at night again after waking up	67	69.8	29	30.2	16	16.7
wants a pacifier when he/she wakes up at night; wants his/her parent to put the pacifier back to his/her mouth	66	68.8	30	31.3	53	55.2
wants to drink something overnight (Sucking Mother's Breast or bottle)	86	89.6	10	10.4	68	70.8
is afraid of sleeping alone	75	78.1	21	21.9	30	31.3
is afraid of sleeping in the dark	79	82.3	17	17.7	46	47.9
hugs an object while falling asleep (blanket, toy, bottle etc.)	70	72.9	26	27.1	71	74.0
is anxious and restless during the day because of not having enough sleep at the previous night	80	83.3	16	16.7	30	31.3
wants to be read books or listen to a lullaby before going to sleep	74	77.1	22	22.9	54	56.3
feels restless while sleeping	95	99.0	1	1.0	20	20.8
wakes up screaming, crying, wakes up due to a nightmare	95	99.0	1	1.0	23	24.0

There was a significant differences between pre and post burn periods on children’s sleeping and waking hour, total and night sleep duration, total daily sleep duration, existence of sleep interruption and number of interruptions, trouble of falling asleep and sleep latency, being tired in the morning, forcing the children to wake up, duration of waking up.

With the knowledge about the importance of sleep, attention should be paid to sleep of children that we are providing care. Supportive environmental regulations should be made to improve the quality of sleep in hospitals.

CONCLUSION