Angiopoietin-1 mimetic preserves microcirculatory perfusion during cardiopulmonary bypass in a rat model

Dekker N.A.M.^{1,3}, van den Brom C.E.^{1,3}, van Leeuwen A.L.I.^{1,3}, Koning N.J.¹, Vonk A.B.A.², Boer C.¹

Departments of Anesthesiology¹, Cardio-thoracic Surgery² and Physiology³, VU University Medical Center, Amsterdam, The Netherlands.

Background

Cardiopulmonary bypass (CPB) during cardiac surgery impairs microcirculatory perfusion and is paralleled by microvascular leakage which is associated with postoperative organ failure and poor patient outcome. Interfering in the Angiopoietin/Tie2 system might preserve endothelial barrier function and attenuate microvascular leakage. This study investigated the effects of an Angiopoietin-1 mimetic on microcirculatory perfusion in a rat CPB model.

Materials and methods

Male rats were anesthetized and subjected to 75 minutes of CPB after treatment with PBS (CPB; n=10), an angiopoietin-1 mimetic (CPB + Ang1mim; n=9) or underwent sham procedure (SHAM; n=9). Microcirculatory perfusion was measured in the cremaster muscle using intravital microscopy at baseline and five time points until 60 minutes after CPB.

Results

Unaltered hemodynamics

Fig. 2: Onset of CPB resulted in comparable blood pressures, hematocrit levels and HCO_3^- levels in the CPB and CPB + Ang1mim groups, respectively.

Intravital microscopy

Fig. 3: Isolated and prepared cremaster muscle (A) to perform microcirculatory measurements (B) using intravital microscopy.

Preserved microvascular perfusion

Fig. 4: Pretreatment with an angiopoietin-1 mimetic preserved microvascular perfusion during cardiopulmonary bypass. CPB = cardiopulmonary bypass, CPB + Ang1mim = cardiopulmonary bypass pretreated with an ang1-mimetic. Two-way ANOVA, * $P < 0.0001$ CPB versus CPB + Ang1mim.

Conclusions

Intervening with an **Angiopoietin-1 mimetic** prevents CPB-induced microcirculatory perfusion disturbances suggesting that therapeutic targeting the Angiopoietin/Tie2 system may contribute to **preserved microcirculatory function during extracorporeal circulation**.