Bewilderment & dismay at lack of staff competence

Introduction

Vascular access devices (VADs), although essential, are associated with many potential complications. These complications include failure to cannulate; ecchymosis, haematoma; vaso-vagal reaction; syncope; nerve injury; arterial puncture; mechanical, chemical and infective phlebitis; infiltration; extravasation and infection (*Ray-Barruel et al., 2013*). Although some of these complications might be viewed as minor, according to *Zhang et al., (2016)* there is potential for significant economic consequences as a result of infections and in extreme cases of mortality related to systemic infections. To prevent such complications, it is imperative that healthcare professionals (HCPs) dealing with VADs possess that necessary knowledge and skills to manage them safely. However, it is often decided at a local level what specific training and education is required. As a result, many HCP do not routinely receive education and training for VADs care and maintenance. When patients present with these devices, HCPs are often unable to use the devices and therefore resort to performing venepuncture which we know can be traumatic for some patients. This issue was highlighted in a recent study focusing on the patient's experience of vascular access.

Study aim

To gain an insight into the experiences of patients living with VADs.

Method

This study followed an Interpretive Phenomenological Analysis (IPA) approach. A purposive sampling technique was used to identify eleven patients who had a VAD in situ and were willing to share their experiences. One-to-one semi-structured interviews were performed. The interviews were digitally recorded and transcribed by the principle investigator and analysed using IPA.

Findings

Although three super-ordinate themes were identified, this poster will focus on the theme Bewilderment at lack of staff ability/competence to care for device. The patients in this study highlighted issues with competence and confidence of practitioners caring for their VAD. Patients either voiced their fears and concerns to the practitioner, refused the practitioner access to their devices, resorted to self-care or had to have treatment delivered through peripheral veins. The patients appeared to be bewildered that nurses and doctors were unfamiliar and at times unwilling to use their devices in certain clinical settings. This also led to feelings of anxiety and fear of device complications. One relative discussing her husband's experience explained how he refused to let staff touch his device:

"He was in the hospital with a chest infection and he wouldn't let the doctors touch it... He said you can't touch it unless you're qualified" (John, PORT)

This issue was highlighted when the patients themselves could access and care for devices themselves when healthcare professionals could not.

"It didn't take long to train him, I said to the nurses my husband learned in 10 mins and he's not medically minded and yet you are nurses and you're not willing to get trained... My 12 year old used to come and clean it." (Amaya, PORT)

The patients in the study could not understand why certain individuals could access their device while others could not. They also described the disappointment of having to revert to peripheral cannulation which they had previously found to be traumatic:

"The only issue is that everyone is trained to use them because I can get my district nurse to come out and take a sample but if I go to my GP he has to do it through a cannula or a conventional syringe and I find that strange particularly as I've got this in place specifically for that purpose and you've got medics that aren't able to use it. There's differences. Everybody's trained up, but my GP can't use it and the nurses at my GP practice can't use it, but my district nurses can use it." (Sam, PICC)

Most of the participants had experience of variations in practice between the hospital setting and within the community setting.

"The care is different in the community, yes, and you try to be loyal and you try not to be critical and you say, well if that's all you've got. You shouldn't be caught in the middle like that." (Tina, PICC)

Finally, some patients became so disheartened about the care received that they chose to look after the devices themselves.

"There was always a complication somewhere in the chain, for the simple thing of getting your dressing changed to the point I ended up doing it myself." (Ruby, TCVC)

Discussion

The findings from this study suggest that there remains a lack of consistent and reliable education and training for some practitioners responsible for caring and maintaining VADs. This was more evident within community care. This lack of competence and consistency leaves patients feeling bewildered and dismayed. More disturbingly it means that patients often have to endure repeated peripheral cannulations rather than have the devices used for the specific reason they were placed. Patients in this current study offered insight into practice variations which echoes finding from *Hainey, Green and Kelly (2017)* and in the findings of a study by *Ritchie, et al., (2015)* who discussed patients taking a 'defensive stance' when approached by healthcare professionals who did not appear to be competent. Similar findings were uncovered during a study by *Alpenberg, Joelsson and Rosengren (2015)* where patients described feelings of insecurity and concerns about potential complications resulting from incompetence of staff caring for their devices. Finally, an earlier study by *Oakley, Wright and Ream (2000)* demonstrated the importance of ongoing training in the care of patients with VADs. It was stated that such training resulted in a reduction of catheter – related complications.

Conclusion

When staff are unable to competently and confidently access and use long-term devices, patients must return to the previous painful procedure of peripheral cannulation which they thought they had left behind. We are aware that care and maintenance training and education in the form of dedicated theoretical and practical workshops can improve the confidence and competence of staff *Kelly, Green and Hainey (2015)* and this is further evidenced when coupled with an eLearning element *Hainey, Green, & Kelly (2016)*. Initiatives such as these would ensure staff competence and thus increase satisfaction and outcomes for patients. Finally, we should consider making education and training in VAD care and maintenance mandatory and this should be performed within pre-registration nursing and at medical schools. An alternative would be dedicated vascular access teams who could provide necessary care and offer training required.

Authors: Linda Kelly PhD student, RN, BSc, MSc, PgC TLHE, FHEA, TCH
Professor Austyn Snowden PhD, Napier University & Karen Campbell MSc, Napier University

References: Alpenberg, S., Joelsson, G. and Rosengren, K. (2015) 'Feeling Confident in Using PICC Lines: Patients' Experiences of Living With a PICC Line During Chemotherapy Treatment', *Home Health Care Management and Practice*, 27(3), pp. 119–125. doi: 10.1177/1084822314566300.

Hainey, K., Green, A. and Kelly, L. J. (no date) 'A blended learning approach to teaching CVAD care and maintenance'.

Kelly, L. J., Green, A. and Hainey, K. (2015) 'Implementing a new teaching and learning strategy for CVAD care.', *British journal of nursing* (Mark Allen Publishing), 24(8), pp. S4-8, S9, S12. doi: 10.12968/bjon.2015.24.Sup8.S4.

Oakley, C., Wright, E. and Ream, E. (2000) 'The experiences of patients and nurses with a nurse-led peripherally inserted central venous catheter line service', *European Journal of Oncology Nursing*. Harcourt Publishers Ltd, 4(4). doi: 10.1054.

Ray-Barruel, G., Polit F., D., Murfield, J. and Rickard M., C. (2013) 'Infusion Phlebitis Assessment Scales: A Systematic Review of Their Use and Psychometric Adequacy', *Journal of the Association for Vascular Access*, 18(4), p. 217. doi: 10.1016/j.jva.2013.10.006.

Ritchie, M., Kelly, L. J., Moss, J., Paul, J. and Shaw, R. (2015) 'Exploring attitudes towards a randomised controlled trial of venous access devices – a nested pre-trial qualitative study.', *The journal of vascular access*, 16(5), pp. 407–12. doi: 10.5301/jva.5000447.

Zhang, L., Cao, S., Marsh, N., Ray-Barruel, G., Flynn, J., Larsen, E. and Rickard, C. M. (2016) 'Infection risks associated with peripheral vascular catheters', *Journal of Infection Prevention*. SAGE PublicationsSage UK: London, England, 17(5), pp. 207–213. doi: 10.1177/1757177416655472.

For more information please contact **Linda Kelly** on linda.kelly@vygon.co.uk