Obtaining informed consent for anaesthesia in elective surgery at a tertiary-care hospital: practices and ethical-legal context

Casimiro L.G.¹, Pereira S.², Azevedo B.¹, Pires S.³, Abelha F.^{1,2}, Mourão J.^{1,2}
¹Centro Hospitalar São João E.P.E., Dept of Anaesthesiology, Porto, Portugal,
²Faculty of Medicine, University of Porto, Dept of Anaesthesiology, Porto, Portugal
³Public Ministry

BACKGROUND AND GOUL OF STUDY

Currently there’s a growing ethical and legal concern regarding the practice of informed consent (IC) for anaesthesia. This practice has not been adopted on a regular basis in all institutions, despite our general health institution recommending it. The anaesthetic practice has inherent risks, aside those from the surgery specific-ones, which makes the anaesthetic consent of great importance. Anaesthetic IC should never be neglected. The aim of this study was to evaluate the degree of implementation of specific informed consent for anaesthesia in the context of elective surgery at a tertiary hospital.

MATERIALS AND METHODS

It was conducted an observational prospective study.

Inclusion criteria	Exclusion criteria
Patients aged >60 years old	Inability to give informed consent
Submitted to elective surgery	ICU admission after surgery
May – July 2017	

Demographic data, the Charlson Comorbidity Score, previous medical history, clinical records and Clinical Frailty Score were collected. The patients were divided in two groups, those who had informed consent signed by the anaesthesiologist (IC) and those who had not (NIC). The Chi-square, Fisher’s exact or Mann-Whitney U tests were used for comparison.

RESULTS AND DISCUSSION

- It was obtained a sample of 230 patients.
- Only 41.7% of the informed consent were signed by the anaesthesiologist, while 97.8% were signed by the surgeon.

Variable	IC	NIC	P value
Female n (%)	54 (43.5)	70 (56.5)	0.547
Male n (%)	42 (39.6)	64 (60.4)	
Age n (p25-p75)	69 (65.0-77.0)	70 (65.0-77.0)	0.863
ASA Score n (%)			0.046
1	6 (60.0)	4 (40.0)	
2	56 (44.8)	69 (55.2)	
3	32 (37.6)	53 (62.4)	
4	2 (20.0)	8 (80.0)	
Frailty Clinical Scale			0.844
Very Fit	11 (45.8)	13 (54.2)	
Well	21 (45.7)	25(54.3)	
Managing Well	27 (38.0)	44 (62.0)	
Vulnerable	14 (35.9)	25 (64.1)	
Midly Frail	9 (56.2)	7 (43.8)	
Moderately Frail	9 (42.9)	12 (57.1)	
Severely Frail	5 (38.5)	8 (61.5)	
Charlson score n (p25-p75)	5.0 (3.0 – 9.0)	7.0 (4.0 – 10.0)	0.017
Previous Stroke n (%)			0.021
Yes	2 (13.3)	13 (86.7)	
No	94 (43.7)	121 (56.3)	

ASA - American Society of Anesthesiologists

Variable	IC	NIC	P value
Dementia n (%)			0.508
Yes	2 (33.3)	4 (66.7)	
No	94 (42.0)	130 (58.0)	
Psychiatric disease n (%)			0.640
Yes	15 (45.5)	18 (54.5)	
No	81 (41.1)	116 (58.9)	
Anemia n (%)			0.028
Yes	13 (27.7)	34 (72.3)	
No	83 (45.4)	100 (54.6)	
Heart failure n (%)			0.798
Yes	7 (38.9)	11 (61.1)	
No	89 (42.0)	123 (58.0)	
Isquemic heart disease n (%)			0.973
Yes	8 (42.1)	11 (57.9)	
No	88 (41.7)	123 (58.3)	
Miocardial infarction n (%)			0.721
Yes	6 (37.5)	10 (62.5)	
No	90 (42.1)	124 (57.9)	
Chronic obstructive pulmonarydisease n (%)			0.484
Yes	4 (36.4)	7 (63.6)	
No	92 (42.0)	127 (58.0)	

CONCLUSION

- It was identified a low practice of informed consent for anaesthesia procedures.
- There seems to be an association regarding an increased absence of informed anesthetic consent and:
 - Higher ASA score and higher Charlson score.
 - Previous history of stroke
 - Anemia
- The informed consent is essential for the patient clarification about the anaesthesia practice as a unique medical speciality.
- It is a ethical e legal duty of great importance that cannot be neglected.