PATIENT COMPLEXITY AND RISK FACTORS ASSOCIATED WITH PHLEBITIS

Adamuz J^{1,2}, Jiménez-Martínez E³, López-Jiménez MM¹, Tapia-Pérez M¹, Rodríguez-Fernández H¹, González-Samartino M^{1,2} and Juvé-Udina ME².

¹Nursing Information Systems Department Support, Hospital Universitari de Bellvitge (GRIN-IDIBELL). Barcelona, Spain. ²Nursing, Faculty of Medicine and Health Sciences, University of Barcelona (GRIN-IDIBELL). Barcelona, Spain. ³Department of Infectious Disease, Hospital Universitari de Bellvitge (GRIN-IDIBELL). Barcelona, Spain.

INTRODUCTION

- Phlebitis is a common adverse event of peripheral venous catheter¹.
- Prevention of complications must focus on optimizing the use of these devices and identify the risk factors.
- Complex patients have more frequent adverse events and are at higher risk for poor health outcomes².
- Patients can be complex not only due to multiple co-occurring medical conditions, but also due to their socioeconomic, cultural, behavioral, and environmental circumstances (*Vector Model of Complexity*)³.
- Care complexity individual factors on in-patients are included in the five domains⁴:

MENTAL-COGNITIVE, PSYCHO-EMOTIONAL, SOCIOCULTURAL, DEVELOPMENTAL, AND COMORBIDITY/COMPLICATIONS

¹Göransson K, et al. Measurement of peripheral venous catheter-related phlebitis: a cross-sectional study. *Lancet Haematol*. 2017.
²Adamuz J, et al. Care complexity individual factors associated with hospital readmission. *Journal of Nursing Scholarship*. 2018.
³Safford MM, et al. Patient complexity: more than comorbidity. The vector model of complexity. *J Gen Intern Med*. 2007.
⁴Juvé-Udina ME, et al. Intensity of nursing care: workloads or individual complexity? *Metas de Enferm*. 2010.

OBJECTIVE

To identify care complexity individual factors associated with peripheral venous catheter (PVC)-related phlebitis.

FINDINGS

Of 16,925 episodes of hospitalized patients, 1,440 (8.5%) had PVC-related phlebitis and 15,190 (89.7%) had one or more individual care complexity factors recorded in the nursing assessment data.

MAIN CLINICAL CHARACTERISTICS

	Phlebitis n=1,440 No. (%)	Non-phlebitis n=15,485 No. (%)	P-value
Demographic characteristics			
Age (years), median (IQR)	69 (57-78)	68 (55-77)	0.03
Age ≥ 75 years	36 (2.5)	342 (2.2)	0.46
Male sex	826 (57.4)	9,138 (59)	0.23
Continuity of care (other center)	175 (12.2)	1,427 (9.2)	<0.001
Medical ward	830 (57.6)	7,907 (51.1)	<0.001
ICU admission	82 (5.7)	1,092 (7.1)	0.06
Underlying disease			
Chronic respiratory disease	136 (9.4)	1,451 (9.4)	0.93
Chronic heart disease	160 (11.1)	1,937 (12.5)	0.13
Chronic liver disease	134 (9.3)	779 (5)	<0.001
Chronic renal disease	172 (11.9)	2,097 (13.5)	0.09
Neurodegenerative disease	5 (0.3)	169 (1.1)	0.007
Cancer	237 (16.5)	2,667 (17.2)	0.49
Immunosuppression	24 (1.7)	179 (1.2)	0.09
Duration of first hospitalization (days), median (IQR)	12 (7-21)	6 (3-11)	<0.001

Abbreviations: IQR, interquartile range; ICU, intensive care unit

MULTIVARIATE ANALYSIS

Factor	Odds ratio	95% confidence interval	P-value
Agitation	1.39	0.89-2.17	0.15
Consciousness disorders	0.99	0.78-1.27	0.95
Perception of reality disorders	1.13	0.79-1.60	0.49
Impaired adaptation	1.18	0.94-1.48	0.16
Fear/anxiety	1.03	0.85-1.24	0.77
Lack of caregiver support	1.01	0.74-1.38	0.96
Old age (≥75 years)	1.09	0.96-1.56	0.63
Hemodynamic instability	1.92	1.68-2.18	<0.001
High-risk of hemorrhage	2.27	1.76-2.92	<0.001
Communication disorders	1.19	0.95-1.51	0.13
Mixed incontinence	1.19	0.96-1.49	0.11
Uncontrolled pain	1.59	1.42-1.78	<0.001
Transmissible infection	2.24	1.85-2.70	<0.001

INDIVIDUAL CARE COMPLEXITY FACTORS OF ADULT PATIENTS ACCORDING PVC-RELATED PHLEBITIS

DOMAIN/Complexity factor	Phlebitis n=1,440 No. (%)	Non-phlebitis n=15,485 No. (%)	P-value
MENTAL-COGNITIVE			
Agitation	27 (1.9)	160 (1)	0.006
Consciousness disorders	116 (8.1)	954 (6.2)	0.006
Impaired cognitive functions	5 (0.3)	37 (0.2)	0.40
Perception of reality disorders	44 (3.1)	337 (2.2)	0.04
PSYCHO-EMOTIONAL			
Aggressive behaviour	7 (0.5)	57 (0.4)	0.50
Fear/anxiety	155 (10.8)	1,383 (8.9)	0.02
Impaired adaptation	114 (7.9)	877 (5.7)	<0.001
SOCIOCULTURAL			
Language barriers	11 (0.8)	97 (0.6)	0.49
Social rejection	1 (0.1)	15 (0.1)	-
DEVELOPMENTAL			
Cultural shock	4 (0.3)	40 (0.3)	0.79
Lack of caregiver support	54 (3.8)	438 (2.8)	0.046
Old age (≥75 years)	36 (2.5)	342 (2.2)	0.46
Adolescence (13-19 years)	2 (0.1)	53 (0.3)	0.32
COMORBIDITY/COMPLICATIONS			
Major chronic disease	663 (46)	7,168 (46.3)	0.87
Hemodynamic instability	1,117 (77.6)	9,622 (62.1)	<0.001
High-risk of hemorrhage	80 (5.6)	375 (2.4)	<0.001
Communication disorders	107 (7.4)	827 (5.3)	<0.001
Mixed incontinence	110 (7.6)	867 (5.6)	0.002
Vascular fragility	12 (0.8)	127 (0.8)	0.88
Functional impotence	53 (3.7)	473 (3.1)	0.20
Dyskinesias	12 (0.8)	79 (0.5)	0.13
Extreme weight	128 (8.9)	1,219 (7.9)	0.69
Dehydration	0 (0)	2 (0)	-
Edema	19 (1.3)	251 (1.6)	0.44
Uncontrolled pain	641 (44.5)	4,855 (31.4)	<0.001
Transmissible infection	151 (10.5)	693 (4.5)	<0.001
Immunosuppression	24 (1.7)	179 (1.2)	0.09
Anatomical and functional disorders	159 (11)	1,469 (9.5)	0.06

After adjustment for potential confounders in the multivariate analysis, we found that hemodynamic instability, high-risk of hemorrhage, uncontrolled pain and transmissible infection were independent factors associated with PVC-related phlebitis. The goodness-of-fit of the model was 0.917.

CONCLUSIONS

- Identify care complexity individual factors during the admission can predict adverse outcomes such as PVC-related phlebitis.
- Independent factors associated with PVC-related phlebitis were hemodynamic instability, high-risk of hemorrhage, uncontrolled pain and transmissible infection.
- Our study demonstrated the need for vigilance in ensuring efficacy in device use among at-risk patients.
- This findings can help to implement strategies for optimize use and type of intravenous catheter on hospitalized patients.