The Uptake of Laparoscopic Colon Cancer Surgery in Canada: 2004-2014

C. Hoogerboord MB ChB, MMed, FCS(SA), FRCSC, FACS (marius.hoogerboord@dal.ca),
M. Hu MA, G. Flowerdew DSc, A. Levy PhD, G. Porter MD, MSc, FRCSC, FACS

Introduction

- Compared to open colectomy, laparoscopic colectomy for cancer (LAC) has equivalent oncologic outcomes and enhances early postoperative recovery¹⁻⁴
- No published data examine the uptake of LAC in Canada

Objectives

- To describe the uptake of LAC in Canada from April 01, 2004 to March 31, 2015
- To estimate the impact of demographic (province, rural/urban residence, year of surgery), patient (age, sex, comorbidities, location of tumor), and system (surgeon and hospital volume) factors on use of LAC

Methods

- 01/04/2004-31/03/2015: All non-emergent colectomies for colon cancer performed in all Canadian provinces, except Quebec, were identified from the Discharge Abstract Database (DAD) of the Canadian Institute for Health Information (CIHI)
- Pregnant patients and those who required multivisceral resection were excluded
- Comorbid status was defined according to the Elixhauser comorbidity index
- Average annual surgeon and hospital volumes were categorized as quartiles and dichotomized into high and low, defined as above and below the upper quartile
- Multiple logistic regression was used to estimate the association between predictor variables and the use of LAC

Results

Figure 1. Proportion of laparoscopic colectomy in Canada, 2004-2014

Figure 2. Proportion of laparoscopic colectomy by province, 2004-2014

References:
1. COST Study Group (2004) *Lancet Oncol.* 6: 477-84
2. COLOR Study Group (2009) *Lancet Oncol.* 10: 44-52
3. MRC CLASICC Trial Group (2007) *J Clin Oncol.* 25: 3061-8
4. Lacy et al. (2002) *Lancet* 359:2224-9

Table 1. Logistic regression of factors associated with LAC

Variable	Multivariate Adjusted OR (95% CI)	Multivariate p-value
Odds ratio estimate, 2014 relative to 2004	9.31 (8.60-10.09)	*
Sex		
Female (reference)	1.00	
Male	0.94 (0.90-0.98)	0.001
Age		
18-50 years	1.15 (1.07-1.24)	*
51-65	1.14 (1.09-1.19)	*
66-80 (reference)	1.00	
>80	0.87 (0.83-0.92)	*
Province		
Newfoundland	0.14 (0.12-0.16)	*
New Brunswick	0.16 (0.14-0.19)	*
Prince Edward Island	0.23 (0.17-0.32)	*
Manitoba	0.39 (0.36-0.43)	*
Saskatchewan	0.54 (0.49-0.60)	*
Nova Scotia	0.53 (0.48-0.59)	*
Alberta	0.57 (0.53-0.60)	*
British Columbia	0.89 (0.85-0.93)	*
Ontario (reference)	1.00	
Residence		
Rural (reference)	1.00	
Urban	1.24 (1.18-1.30)	*
Elixhauser score		
0 (reference)	1.00	
1-3	0.90 (0.87-0.94)	*
4-8	0.79 (0.63-0.98)	0.035
Hospital volume		
Low (reference)	1.00	
High	2.04 (1.96-2.13)	*
Surgeon volume		
Low (reference)	1.00	
High	1.29 (1.24-1.35)	*
Resection type		
Right (reference)	1.00	
Left	0.91 (0.87-0.95)	*
Anterior	0.58 (0.55-0.62)	*
Other	0.71 (0.66-0.76)	*

*P<0.001

Summary of Results

- Among 63,504 cases overall, 19,691 (31%) involved LAC
- The proportion of LAC increased from 9% in 2004 to 52% in 2014 in relatively constant fashion
- 78% of colectomies were performed in 3 provinces: Ontario (50%), British Columbia (18%) and Alberta (10%)
- Provincial uptake ranged from 8% in Newfoundland to 36% in Ontario
- By 2014, the majority of colectomies were being performed laparoscopically in Ontario, British Columbia and Alberta
- Predictor variables of LAC included later year of study, female gender, urban residence, right hemicolectomy, high provincial uptake and high hospital and surgeon volume of colectomies.
- Older patients and those with higher level of comorbidities were less likely to undergo LAC

Conclusion

- While the pan-Canadian use of laparoscopic colectomy increased rapidly between 2004 and 2014, substantial interprovincial variation exists that is not attributable to demographic, clinical or system factors