DETERMINING VIRAL ETIOLOGY: TRAIL BIOMARKER LEVELS AND MOLECULAR VIRAL DETECTION IN THE AUTOPILOT-DX STUDY

Niv Mastboim¹, Edoardo Farinelli², Cihan Papan³, Liat Etshtein¹, Asi Cohen¹, Einav Simon¹, Olga Boico¹, Liran Shani¹, Tanya Gottlieb¹, Kfir Oved¹, Eran Eden¹, Ilaria Testa², Alberto Argentiero², Maria Bruna Pasticci², Daniele Mezzetti², Katia Perruccio², Marian Porwoll³, Ummaya Hakim³, Arne Simon⁴, Johannes G. Liese⁵, Markus Knuf⁶, Sven Schneider⁷, Tobias Tenenbaum^{3*}, Susanna Esposito²

¹MeMed Diagnostics, Tirat Carmel, Israel, ²Pediatric Clinic, Università degli Studi di Perugia, Perugia, Italy, ³University Children's Hospital Mannheim, Mannheim, Germany, ⁴Pediatric Oncology and Haematology, Saarland University Hospital, Homburg, Germany, ⁵Department of Pediatrics, University of Würzburg, Würzburg, Germany, ⁶Children's Hospital, Dr. Horst Schmidt Klinik, Wiesbaden, Germany, ⁷Institute for Clinical Chemistry, University of Heidelberg Medical Faculty Mannheim, Mannheim, Germany. *shared senior author

Background

TRAIL serves as a useful biomarker for distinguishing between bacterial and viral infections when combined with IP-10 and CRP (ImmunoXpert™)

Name	Curiosity	Pathfinder	Opportunity
Target population	Adults and pediatric with acute infection	Pediatric with acute infection	Pediatric with FWS or RTI
Potentially eligible patients (n)	1002	597	777

* Oved et al., PLoS ONE, 2015
* Srugo et al., Pediatrics, 2017
* Van Houten et al., Lancet ID, 2016

AutoPilot-Dx study primary objective

To validate the diagnostic accuracy and potential clinical utility of the host-immune signature (ImmunoXpert™).

Design

Sub-analysis goal

To evaluate TRAIL levels in children with acute bacterial or viral expert diagnosis and nasopharyngeal PCR virus detection

Results and conclusions

TRAIL levels has potential to correctly classify febrile children with viral detection

These are the three most frequently detected viruses, accounting for a total of 113 out of the 159 viruses detected.

Bacterial expert diagnosis was assigned to 29%, 29% and 25% of patients with adenovirus, rhinovirus and respiratory-syncytial virus detection, respectively, highlighting that viral detection may not necessarily indicate underlying etiology.

TRAIL levels were significantly increased in viral patients as compared to bacterial patients, irrespective of virus detection.

Conclusion: The differential expression of TRAIL in response to viral versus bacterial infections can complement molecular viral detection in the classification of febrile children.